

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
2 August 2001 (02.08.2001)

PCT

(10) International Publication Number
WO 01/55085 A1

(51) International Patent Classification⁷: C07C 69/708,
59/64, C07D 213/55, 307/80, 317/54, 215/14, 307/40,
A61K 31/435, 31/34, 31/357, A61P 3/04, 3/10

(DK). PETTERSSON, Ingrid; Grundtvigsvej 18, 2th,
DK-1864 Frederiksberg (DK).

(21) International Application Number: PCT/DK01/00058

(22) International Filing Date: 26 January 2001 (26.01.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
PA 2000 00136 28 January 2000 (28.01.2000) DK
PA 2000 01071 7 July 2000 (07.07.2000) DK
PA 2000 01594 25 October 2000 (25.10.2000) DK

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:


- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(71) Applicant: NOVO NORDISK A/S [DK/DK]; Corporate
Patents, Novo Allé, DK-2880 Bagsværd (DK).

(72) Inventors: MOGENSEN, John, Patrick; Rørløkken
96, DK-2730 Herlev (DK). SAUERBERG, Per; Syren-
vænget 27, DK-3520 Farum (DK). BURY, Paul, Stanley;
Hjortholms Allé 48, DK-2400 København NV (DK).
JEPPESEN, Lone; Malmosevej 121, DK-2830 Virum

(54) Title: PROPIONIC ACID DERIVATIVES AND THEIR USE IN THE TREATMENT OF DIABETES AND OBESITY



(57) Abstract: The present invention relates to compounds of general formula (I). The compounds are useful in the treatment and/or prevention of conditions mediated by nuclear receptors, in particular the Peroxisome Proliferator-Activated Receptors (PPAR).

(I)

WO 01/55085 A1

NEW COMPOUNDS, THEIR PREPARATION AND USE

FIELD OF INVENTION

5

The present invention relates to novel compounds, pharmaceutical compositions containing them, methods for preparing the compounds and their use as medicaments. More specifically, compounds of the invention can be utilised in the treatment and/or prevention of conditions mediated by nuclear receptors, in particular the Peroxisome Proliferator-Activated Re-

10

BACKGROUND OF THE INVENTION

Coronary artery disease (CAD) is the major cause of death in Type 2 diabetic and metabolic syndrome patients (i.e. patients that fall within the 'deadly quartet' category of impaired glucose tolerance, insulin resistance, hypertriglyceridaemia and/or obesity).

15

The hypolipidaemic fibrates and antidiabetic thiazolidinediones separately display moderately effective triglyceride-lowering activities although they are neither potent nor efficacious enough to be a single therapy of choice for the dyslipidaemia often observed in Type 2 diabetic or metabolic syndrome patients. The thiazolidinediones also potently lower circulating glucose levels of Type 2 diabetic animal models and humans. However, the fibrate class of compounds are without beneficial effects on glycaemia. Studies on the molecular actions of these compounds indicate that thiazolidinediones and fibrates exert their action by activating distinct transcription factors of the peroxisome proliferator activated receptor (PPAR) family, resulting in increased and decreased expression of specific enzymes and apolipoproteins respectively, both key-players in regulation of plasma triglyceride content. Fibrates, on the one hand, are PPAR α activators, acting primarily in the liver. Thiazolidinediones, on the other hand, are high affinity ligands for PPAR γ acting primarily on adipose tissue.

20

25

30

Adipose tissue plays a central role in lipid homeostasis and the maintenance of energy balance in vertebrates. Adipocytes store energy in the form of triglycerides during periods of nutritional affluence and release it in the form of free fatty acids at times of nutritional deprivation. The development of white adipose tissue is the result of a continuous

differentiation process throughout life. Much evidence points to the central role of PPAR γ activation in initiating and regulating this cell differentiation. Several highly specialised proteins are induced during adipocyte differentiation, most of them being involved in lipid storage and metabolism. The exact link from activation of PPAR γ to changes in glucose metabolism, most notably a decrease in insulin resistance in muscle, has not yet been clarified. A possible link is via free fatty acids such that activation of PPAR γ induces Lipoprotein Lipase (LPL), Fatty Acid Transport Protein (FATP) and Acyl-CoA Synthetase (ACS) in adipose tissue but not in muscle tissue. This, in turn, reduces the concentration of free fatty acids in plasma dramatically, and due to substrate competition at the cellular level, skeletal muscle and other tissues with high metabolic rates eventually switch from fatty acid oxidation to glucose oxidation with decreased insulin resistance as a consequence.

PPAR α is involved in stimulating β -oxidation of fatty acids. In rodents, a PPAR α -mediated change in the expression of genes involved in fatty acid metabolism lies at the basis of the phenomenon of peroxisome proliferation, a pleiotropic cellular response, mainly limited to liver and kidney and which can lead to hepatocarcinogenesis in rodents. The phenomenon of peroxisome proliferation is not seen in man. In addition to its role in peroxisome proliferation in rodents, PPAR α is also involved in the control of HDL cholesterol levels in rodents and humans. This effect is, at least partially, based on a PPAR α -mediated transcriptional regulation of the major HDL apolipoproteins, apo A-I and apo A-II. The hypotriglyceridemic action of fibrates and fatty acids also involves PPAR α and can be summarised as follows: (I) an increased lipolysis and clearance of remnant particles, due to changes in lipoprotein lipase and apo C-III levels, (II) a stimulation of cellular fatty acid uptake and their subsequent conversion to acyl-CoA derivatives by the induction of fatty acid binding protein and acyl-CoA synthase, (III) an induction of fatty acid β -oxidation pathways, (IV) a reduction in fatty acid and triglyceride synthesis, and finally (V) a decrease in VLDL production. Hence, both enhanced catabolism of triglyceride-rich particles as well as reduced secretion of VLDL particles constitutes mechanisms that contribute to the hypolipidemic effect of fibrates.

30

A number of compounds have been reported to be useful in the treatment of hyperglycemia, hyperlipidemia and hypercholesterolemia (U.S. Pat. 5,306,726, PCT Publications nos.

W091/19702, WO 95/03038, WO 96/04260, WO 94/13650, WO 94/01420, WO 97/36579, WO 97/25042, WO 95/17394, WO 99/08501, WO 99/19313 and WO 99/16758).

SUMMARY OF THE INVENTION

5

Glucose lowering as a single approach does not overcome the macrovascular complications associated with Type 2 diabetes and metabolic syndrome. Novel treatments of Type 2 diabetes and metabolic syndrome must therefore aim at lowering both the overt hypertriglyceridaemia associated with these syndromes as well as alleviation of hyperglycaemia.


10

The clinical activity of fibrates and thiazolidinediones indicates that research for compounds displaying combined PPAR α and PPAR γ activation should lead to the discovery of efficacious glucose and triglyceride lowering drugs that have great potential in the treatment of Type 2 diabetes and the metabolic syndrome (i.e. impaired glucose tolerance, insulin resistance, hypertriglyceridaemia and/or obesity).

15

DETAILED DESCRIPTION OF THE INVENTION

Accordingly, the present invention relates to compounds of the general formula (I):



20

wherein A is aryl or heteroaryl and wherein A is optionally substituted with one or more substituents selected from hydroxy, halogen, perhalomethyl, perhalomethoxy, acyl, cyano, amino, C₁₋₆-alkylamino, C₁₋₆-dialkylamino, methylenedioxy, aralkenyl, aralkynyl, heteroaryloxy, heteroaralkoxy, aralkyl, heteroaralkyl, arylamino, or

- 5 A is optionally substituted with C₁₋₆-alkyl, C₂₋₆-alkenyl or C₂₋₆-alkynyl each of which is optionally substituted with one or more substituents selected from C₁₋₆-alkoxycarbonyl or carboxy, or

A is optionally substituted with C₁₋₆-alkoxy, C₁₋₆-alkylthio or C₂₋₆-alkenyloxy each of which is optionally substituted with one or more halogens, or

- 10 A is optionally substituted with aryloxy, arylthio or aralkoxy each of which is optionally substituted with one or more substituents selected from C₁₋₆-alkoxy, nitro, carboxy or C₁₋₆-alkoxycarbonyl; and

X₁ and X₂ independently are
hydrogen,

- 15 aryl or heteroaryl each of which is optionally substituted with one or more substituents selected from hydroxy, aryloxy, arylthio, aralkoxy, heteroaryloxy, aralkoxy, C₁₋₆-alkoxy, C₁₋₆-alkylthio, halogen, perhalomethyl, perhalomethoxy, acyl, aryl, heteroaryl, aralkyl, heteroaralkyl, cyano, amino, C₁₋₆-alkylamino, C₁₋₆-dialkylamino, arylamino or methylenedioxy, or
20 aryl or heteroaryl each of which is optionally substituted with one or more substituents selected from C₁₋₆-alkyl, C₂₋₆-alkenyl or C₂₋₆-alkynyl each of which is optionally substituted with hydroxy; or

A is selected from the ring systems consisting of




Figure 1

wherein the attachment point of A to the remaining part of the structure of formula (I) is as indicated on the chemical structures in Figure 1, and wherein A is optionally substituted with one or more substituents selected from C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, hydroxy, aryloxy, arylthio, aralkoxy, C₁₋₆-alkoxy, C₁₋₆-alkylthio, heteroaryloxy, heteroaralkoxy, halogen, perhalomethyl, perhalomethoxy, acyl, aralkyl, heteroaralkyl, cyano, amino, C₁₋₆-alkylamino, C₁₋₆-dialkylamino, arylamino or methylenedioxy; and

wherein X₁ and X₂ are hydrogen; and

R₅ is hydrogen or C₁₋₆-alkyl; and

Y is hydrogen, or

Y is C₁₋₁₂-alkyl, C₂₋₁₂-alkenyl, C₂₋₁₂-alkynyl, C₄₋₁₂-alkenynyl, aralkyl or heteroaralkyl each of which is optionally substituted with one or more substituents selected from halogen, C₁₋₆-alkyl, perhalomethyl, hydroxy, aryl, heteroaryl, carboxy or amino; and

Z is hydrogen, halogen, hydroxy, or

Z is C₁₋₆-alkyl or C₁₋₆-alkoxy each of which is optionally substituted with one or more substituents selected from halogen, hydroxy, carboxy, amino, cyano or C₁₋₆-alkoxy; and

Q is O, S or NR₆, wherein R₆ is hydrogen, C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, C₄₋₆-alkenynyl, aralkyl, heteroaralkyl and wherein R₆ is optionally substituted with one or more substituents selected from halogen, hydroxy, C₁₋₆-alkoxy, amino or carboxy; and

- 5 Ar is arylene, heteroarylene or a divalent heterocyclic group each of which can be optionally substituted with one or more substituents selected from C₁₋₆-alkyl, aryl or C₁₋₆-alkoxy each of which can be optionally substituted with halogen, hydroxy, carboxy, cyano or heterocyclyl; and

R₁ is hydrogen, hydroxy or halogen; or R₁ forms a bond together with R₂; and

10

R₂ is hydrogen or C₁₋₆-alkyl; or R₂ forms a bond together with R₁; and

R₃ is hydrogen, or

- 15 R₃ is C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, C₄₋₆-alkenynyl, aryl, aralkyl, C₁₋₆-alkoxyC₁₋₆alkyl, acyl, heterocyclyl, heteroaryl or heteroaralkyl each of which is optionally substituted with one or more substituents selected from halogen, perhalomethyl, hydroxy or cyano; and


R₄ is hydrogen, C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, C₄₋₆-alkenynyl or aryl; and

- 20 n is an integer ranging from 0 to 3; and

m is an integer ranging from 0 to 1;

- 25 or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate thereof, or any tautomeric forms, stereoisomers, mixture of stereoisomers including a racemic mixture, or polymorphs.

In a preferred embodiment, the present invention is concerned with compounds of formula (I)



wherein A is aryl or heteroaryl and wherein A is optionally substituted with one or more substituents selected from hydroxy, halogen, perhalomethyl, perhalomethoxy, acyl, cyano, amino, C₁₋₆-alkylamino, C₁₋₆-dialkylamino, methylenedioxy, aralkenyl, aralkynyl, heteroaryloxy, heteroaralkoxy, aralkyl, heteroaralkyl, arylamino, or

A is optionally substituted with C₁₋₆-alkyl, C₂₋₆-alkenyl or C₂₋₆-alkynyl each of which is optionally substituted with one or more substituents selected from C₁₋₆-alkoxycarbonyl or carboxy, or

A is optionally substituted with C₁₋₆-alkoxy, C₁₋₆-alkylthio or C₂₋₆-alkenyloxy each of which is optionally substituted with one or more halogens, or

A is optionally substituted with aryloxy, arylthio or aralkoxy each of which is optionally substituted with one or more substituents selected from C₁₋₆-alkoxy, nitro, carboxy or C₁₋₆-alkoxycarbonyl; and

X₁ and X₂ independently are hydrogen,

aryl or heteroaryl each of which is optionally substituted with one or more substituents selected from hydroxy, aryloxy, arylthio, aralkoxy, heteroaryloxy, aralkoxy, C₁₋₆-alkoxy, C₁₋₆-alkylthio, halogen, perhalomethyl, perhalomethoxy, acyl, aryl, heteroaryl, aralkyl, heteroaralkyl, cyano, amino, C₁₋₆-alkylamino, C₁₋₆-dialkylamino, arylamino or methylenedioxy, or

aryl or heteroaryl each of which is optionally substituted with one or more substituents selected from C₁₋₆-alkyl, C₂₋₆-alkenyl or C₂₋₆-alkynyl each of which is optionally substituted with hydroxy; or

A is selected from the ring systems consisting of

5




Figure 1

- 10 wherein the attachment point of A to the remaining part of the structure of formula (I) is as indicated on the chemical structures in Figure 1, and wherein A is optionally substituted with one or more substituents selected from C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, hydroxy, aryloxy, arylthio, aralkoxy, C₁₋₆-alkoxy, C₁₋₆-alkylthio, heteroaryloxy, heteroaralkoxy, halogen, perhalomethyl, perhalomethoxy, acyl, aralkyl, heteroaralkyl, cyano, amino, C₁₋₆-alkylamino, C₁₋₆-dialkylamino, arylamino or methylenedioxy; and
- 15 wherein X₁ and X₂ are hydrogen; and

R₅ is hydrogen or C₁₋₆-alkyl; and

- 20 Y is hydrogen, or

Y is C₁₋₁₂-alkyl, C₂₋₁₂-alkenyl, C₂₋₁₂-alkynyl, C₄₋₁₂-alkenynyl, aralkyl or heteroaralkyl each of which is optionally substituted with one or more substituents selected from halogen, C₁₋₆-alkyl, perhalomethyl, hydroxy, aryl, heteroaryl, carboxy or amino; and

Z is hydrogen, halogen, hydroxy, or

Z is C₁₋₆-alkyl or C₁₋₆-alkoxy each of which is optionally substituted with one or more substituents selected from halogen, hydroxy, carboxy, amino, cyano or C₁₋₆-alkoxy; and

5

Q is O, S or NR₆, wherein R₆ is hydrogen, C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, C₄₋₆-alkenynyl, aralkyl, heteroaralkyl and wherein R₆ is optionally substituted with one or more substituents selected from halogen, hydroxy, C₁₋₆-alkoxy, amino or carboxy; and

10 Ar is arylene, heteroarylene or a divalent heterocyclic group each of which can be optionally substituted with one or more substituents selected from C₁₋₆-alkyl, aryl or C₁₋₆-alkoxy each of which can be optionally substituted with halogen, hydroxy, carboxy, cyano or heterocyclyl; and

R₁ is hydrogen, hydroxy or halogen; or R₁ forms a bond together with R₂; and

15

R₂ is hydrogen or C₁₋₆-alkyl; or R₂ forms a bond together with R₁; and

R₃ is hydrogen, or

20 R₃ is C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, C₄₋₆-alkenynyl, aryl, aralkyl, C₁₋₆-alkoxyC₁₋₆alkyl, acyl, heterocyclyl, heteroaryl or heteroaralkyl each of which is optionally substituted with one or more substituents selected from halogen, perhalomethyl, hydroxy or cyano; and


R₄ is hydrogen, C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, C₄₋₆-alkenynyl or aryl; and

25 n is an integer ranging from 1 to 3; and

m is 1;

30 or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate thereof, or any tautomeric forms, stereoisomers, mixture of stereoisomers including a racemic mixture, or polymorphs.

In another preferred embodiment, the present invention is concerned with compounds of formula (I)



5

wherein A is aryl or heteroaryl and wherein A is optionally substituted with one or more substituents selected from C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, hydroxy, aryloxy, arylthio, aralkoxy, C₁₋₆-alkoxy, C₁₋₆-alkylthio, heteroaryloxy, heteroaralkoxy, halogen, perhalomethyl, perhalomethoxy, acyl, aralkyl, heteroaralkyl, cyano, amino, C₁₋₆-alkylamino, C₁₋₆-dialkylamino, arylamino or methylenedioxy, and

10

X₁ and X₂ independently are hydrogen, aryl or heteroaryl optionally substituted with one or more substituents selected from C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, hydroxy, aryloxy, arylthio, aralkoxy, heteroaryloxy, aralkoxy, C₁₋₆-alkoxy, C₁₋₆-alkylthio, halogen, perhalomethyl, perhalomethoxy, acyl, aryl, heteroaryl, aralkyl, heteroaralkyl, cyano, amino, C₁₋₆-alkylamino, C₁₋₆-dialkylamino, arylamino or methylenedioxy;

15

or

wherein A is selected from the ring systems consisting of

20




Figure 1

- 5 wherein the attachment point of A to the remaining part of the structure of formula (I) is as indicated on the chemical structures in Figure 1, and
- wherein A is optionally substituted with one or more substituents selected from C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, hydroxy, aryloxy, arylthio, aralkoxy, C₁₋₆-alkoxy, C₁₋₆-alkylthio, heteroaryloxy, heteroaralkoxy, halogen, perhalomethyl, perhalomethoxy, acyl, aralkyl, heteroaralkyl, cyano, amino, C₁₋₆-alkylamino, C₁₋₆-dialkylamino, arylamino or methylenedioxy,
- 10 and wherein X₁ and X₂ are hydrogen; and

R₅ is hydrogen or C₁₋₆-alkyl; and

15

Y is hydrogen, C₁₋₁₂-alkyl, C₂₋₁₂-alkenyl, C₂₋₁₂-alkynyl, C₄₋₁₂-alkenynyl, aralkyl or heteroaralkyl optionally substituted with one or more substituents selected from halogen, C₁₋₆-alkyl, perhalomethyl, hydroxy, aryl, heteroaryl, carboxy or amino; and

- 20 Z is hydrogen, halogen, hydroxy, C₁₋₆-alkyl or C₁₋₆-alkoxy optionally substituted with one or more substituents selected from halogen, hydroxy, carboxy, amino, cyano or C₁₋₆-alkoxy; and

Q is O, S or NR₆, wherein R₆ is hydrogen, C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, C₄₋₆-alkenynyl, aralkyl, heteroaralkyl and wherein R₆ is optionally substituted with one or more substituents selected from halogen, hydroxy, C₁₋₆-alkoxy, amino or carboxy; and

- 5 Ar is arylene, heteroarylene or a divalent heterocyclic group each of which can be optionally substituted with one or more substituents selected from C₁₋₆-alkyl, aryl or C₁₋₆-alkoxy each of which can be optionally substituted with halogen, hydroxy, carboxy, cyano or heterocyclyl; and

R₁ is hydrogen, hydroxy or halogen; or R₁ forms a bond together with R₂; and

10

R₂ is hydrogen or C₁₋₆-alkyl; or R₂ forms a bond together with R₁; and

R₃ is hydrogen, C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, C₄₋₆-alkenynyl, aryl, aralkyl, C₁₋₆-alkoxy, C₁₋₆-alkyl, acyl, heterocyclyl, heteroaryl or heteroaralkyl groups optionally substituted
15 with one or more substituents selected from halogen, perhalomethyl, hydroxy or cyano; and

R₄ is hydrogen, C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, C₄₋₆-alkenynyl or aryl; and


n is an integer ranging from 0 to 3; and

20

m is an integer ranging from 0 to 1.


In another preferred embodiment, the present invention is concerned with compounds of formula (I)

25



wherein A is aryl or heteroaryl and wherein A is optionally substituted with one or more substituents selected from C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, hydroxy, aryloxy, arylthio, aralkoxy, C₁₋₆-alkoxy, C₁₋₆-alkylthio, heteroaryloxy, heteroaralkoxy, halogen, perhalomethyl, perhalomethoxy, acyl, aralkyl, heteroaralkyl, cyano, amino, C₁₋₆-alkylamino, C₁₋₆-dialkylamino, arylamino or methylenedioxy; or

provided X₁ and X₂ is hydrogen, A is selected from the ring systems consisting of



wherein A is optionally substituted with one or more substituents selected from C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, hydroxy, aryloxy, arylthio, aralkoxy, C₁₋₆-alkoxy, C₁₋₆-alkylthio, heteroaryloxy, heteroaralkoxy, halogen, perhalomethyl, perhalomethoxy, acyl, aralkyl, heteroaralkyl, cyano, amino, C₁₋₆-alkylamino, C₁₋₆-dialkylamino, arylamino or methylenedioxy; and

R₅ is hydrogen or C₁₋₆-alkyl; and

10

X₁ and X₂ independently are hydrogen, aryl or heteroaryl optionally substituted with one or more substituents selected from C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, hydroxy, aryloxy, arylthio, aralkoxy, heteroaryloxy, aralkoxy, C₁₋₆-alkoxy, C₁₋₆-alkylthio, halogen, perhalomethyl, perhalomethoxy, acyl, aryl, heteroaryl, aralkyl, heteroaralkyl, cyano, amino, C₁₋₆-alkylamino, C₁₋₆-dialkylamino, arylamino or methylenedioxy; and

15

Y is hydrogen, C₁₋₁₂-alkyl, C₂₋₁₂-alkenyl, C₂₋₁₂-alkynyl, C₄₋₁₂-alkenynyl, aralkyl or heteroaralkyl optionally substituted with one or more substituents selected from halogen, C₁₋₆-alkyl, perhalomethyl, hydroxy, aryl, heteroaryl, carboxy or amino; and

20

Z is hydrogen, halogen, hydroxy, C₁₋₆-alkyl or C₁₋₆-alkoxy optionally substituted with one or more substituents selected from halogen, hydroxy, carboxy, amino, cyano or C₁₋₆-alkoxy; and

Q is O, S or NR₆, wherein R₆ is hydrogen, C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, C₄₋₆-alkenynyl, aralkyl, heteroaralkyl and wherein R₆ is optionally substituted with one or more substituents selected from halogen, hydroxy, C₁₋₆-alkoxy, amino or carboxy; and

- 5 Ar is arylene, heteroarylene or a divalent heterocyclic group each of which can be optionally substituted with one or more substituents selected from C₁₋₆-alkyl, aryl or C₁₋₆-alkoxy each of which can be optionally substituted with halogen, hydroxy, carboxy, cyano or heterocyclyl; and

R₁ is hydrogen, hydroxy or halogen; or R₁ forms a bond together with R₂; and

10

R₂ is hydrogen or C₁₋₆-alkyl; or R₂ forms a bond together with R₁; and

- R₃ is hydrogen, C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, C₄₋₆-alkenynyl, aryl, aralkyl, C₁₋₆-alkoxy, C₁₋₆-alkyl, acyl, heterocyclyl, heteroaryl or heteroaralkyl groups optionally substituted
15 with one or more substituents selected from halogen, perhalomethyl, hydroxy or cyano; and

R₄ is hydrogen, C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, C₄₋₆-alkenynyl or aryl; and


n is an integer ranging from 0 to 3; and

20

m is an integer ranging from 0 to 1.

- In another preferred embodiment, the present invention is concerned with compounds of formula I wherein A is aryl or heteroaryl optionally substituted with one or more substituents
25 selected from C₁₋₆-alkyl, C₂₋₆-alkenyl each of which is optionally substituted with one or more substituents selected from C₁₋₆-alkoxycarbonyl or carboxy, or
A is optionally substituted with aryloxy optionally substituted with one or more C₁₋₆-alkoxy, or
A is optionally substituted with aralkoxy optionally substituted with one or more substituents selected from C₁₋₆-alkoxy, nitro, carboxy or C₁₋₆-alkoxycarbonyl, or
30 A is optionally substituted with C₁₋₆-alkoxy optionally substituted with one or more halogens, or
A is optionally substituted with aralkenyl, C₂₋₆-alkenyloxy, aralkynyl, halogen, perhalomethyl, perhalomethoxy, acyl, aralkyl or methylenedioxy.

In another preferred embodiment, the present invention is concerned with compounds of formula I wherein A is aryl, heteroaryl or



optionally substituted with one or more substituents selected from aryloxy, arylthio, aralkoxy, C₁₋₆-alkoxy, C₁₋₆-alkylthio, halogen, perhalomethyl, aralkyl, heteroaralkyl, heteroaryloxy or heteroaralkoxy.

In another preferred embodiment, the present invention is concerned with compounds of formula I wherein A is aryl optionally substituted with one or more substituents selected from aryloxy, arylthio, aralkoxy, C₁₋₆-alkoxy, C₁₋₆-alkylthio, halogen, perhalomethyl, aralkyl, heteroaralkyl, heteroaryloxy or heteroaralkoxy.

In another preferred embodiment, the present invention is concerned with compounds of formula I wherein A is aryl optionally substituted with one or more substituents selected from C₁₋₆-alkyl, C₂₋₆-alkenyl each of which is optionally substituted with one or more substituents selected from C₁₋₆-alkoxycarbonyl or carboxy, or

A is optionally substituted with aryloxy optionally substituted with one or more C₁₋₆-alkoxy, or A is optionally substituted with aralkoxy optionally substituted with one or more substituents selected from C₁₋₆-alkoxy, nitro, carboxy or C₁₋₆-alkoxycarbonyl, or

A is optionally substituted with C₁₋₆-alkoxy optionally substituted with one or more halogens, or

A is optionally substituted with aralkenyl, C₂₋₆-alkenyloxy, aralkynyl, halogen, perhalomethyl, perhalomethoxy, acyl, aralkyl or methylenedioxy.

In another preferred embodiment, the present invention is concerned with compounds of formula I wherein A is aryl optionally substituted with one or more substituents selected from

from C₁₋₆-alkyl, or

A is optionally substituted with aryloxy optionally substituted with one or more C₁₋₆-alkoxy,

A is optionally substituted with aralkoxy optionally substituted with one or more substituents selected from C₁₋₆-alkoxy, or

A is optionally substituted with C₁₋₆-alkoxy optionally substituted with one or more halogens, or


- 5 A is optionally substituted with aralkenyl, aralkynyl, halogen, perhalomethyl, perhalomethoxy or aralkyl.

- 10 In another preferred embodiment, the present invention is concerned with compounds of formula I wherein A is heteroaryl optionally substituted with one or more substituents selected from aryloxy, arylthio, aralkoxy, C₁₋₆-alkoxy, C₁₋₆-alkylthio, halogen, perhalomethyl, aralkyl, heteroaralkyl, heteroaryloxy or heteroaralkoxy.

In another preferred embodiment, the present invention is concerned with compounds of formula I wherein A is heteroaryl.



15

In another preferred embodiment, the present invention is concerned with compounds of formula I wherein A is



- 20 optionally substituted with one or more substituents selected from aryloxy, arylthio, aralkoxy, C₁₋₆-alkoxy, C₁₋₆-alkylthio, halogen, perhalomethyl, aralkyl, heteroaralkyl, heteroaryloxy or heteroaralkoxy.

- 25 In another preferred embodiment, the present invention is concerned with compounds of formula I wherein A is



optionally substituted with one or more substituents selected from C₁₋₆-alkyl, and wherein R₅ is hydrogen or C₁₋₆-alkyl.

In another preferred embodiment, the present invention is concerned with compounds of formula I wherein X_1 and X_2 independently are hydrogen, aryl or heteroaryl optionally substituted with one or more substituents selected from aryloxy, arylthio, aralkoxy, halogen, perhalomethyl, aryl, heteroaryl, aralkyl, heteroaralkyl, heteroaryloxy or heteroaralkoxy.

In another preferred embodiment, the present invention is concerned with compounds of formula I wherein X_1 and X_2 independently are hydrogen, aryl or heteroaryl optionally substituted with one or more substituents selected from halogen, acyl, aryl, or aryl or heteroaryl optionally substituted with one or more substituents selected from C_{1-6} -alkyl, C_{2-6} -alkynyl each of which is optionally substituted with hydroxy.

In another preferred embodiment, the present invention is concerned with compounds of formula I wherein X_1 and X_2 independently are hydrogen, or aryl or heteroaryl optionally substituted with one or more substituents selected from halogen.

In another preferred embodiment, the present invention is concerned with compounds of formula I wherein X_1 and X_2 independently are hydrogen or aryl optionally substituted with one or more substituents selected from aryloxy, arylthio, aralkoxy, halogen, perhalomethyl, aryl, heteroaryl, aralkyl, heteroaralkyl, heteroaryloxy or heteroaralkoxy.

In another preferred embodiment, the present invention is concerned with compounds of formula I wherein X_1 and X_2 independently are hydrogen, aryl optionally substituted with one or more substituents selected from halogen, acyl, aryl, or aryl optionally substituted with one or more substituents selected from C_{1-6} -alkyl, C_{2-6} -alkynyl each of which is optionally substituted with hydroxy.

In another preferred embodiment, the present invention is concerned with compounds of formula I wherein X_1 and X_2 independently are hydrogen, or

phenyl optionally substituted with one or more substituents selected from halogen.

5 In another preferred embodiment, the present invention is concerned with compounds of formula I wherein X_1 and X_2 independently are hydrogen or heteroaryl optionally substituted with one or more substituents selected from aryloxy, arylthio, aralkoxy, halogen, perhalomethyl, aryl, heteroaryl, aralkyl, heteroaralkyl, heteroaryloxy or heteroaralkoxy.

10 In another preferred embodiment, the present invention is concerned with compounds of formula I wherein X_1 and X_2 independently are hydrogen or heteroaryl.

In another preferred embodiment, the present invention is concerned with compounds of formula I wherein X_1 and X_2 is hydrogen.

15 In another preferred embodiment, the present invention is concerned with compounds of formula I wherein Y is hydrogen or C_{1-12} -alkyl.

In another preferred embodiment, the present invention is concerned with compounds of formula I wherein Y is hydrogen or methyl.

20 In another preferred embodiment, the present invention is concerned with compounds of formula I wherein Z is hydrogen or C_{1-6} -alkoxy.

25 In another preferred embodiment, the present invention is concerned with compounds of formula I wherein Z is hydrogen or C_{1-6} -alkyl.

In another preferred embodiment, the present invention is concerned with compounds of formula I wherein Z is hydrogen.

30 In another preferred embodiment, the present invention is concerned with compounds of formula I wherein Q is O.

In another preferred embodiment, the present invention is concerned with compounds of formula I wherein Ar is arylene optionally substituted with one or more substituents selected from C_{1-6} -alkyl or C_{1-6} -alkoxy each of which can be optionally substituted with carboxy.

In another preferred embodiment, the present invention is concerned with compounds of formula I wherein Ar is arylene.

- 5 In another preferred embodiment, the present invention is concerned with compounds of formula I wherein R_1 is hydrogen or R_1 forms a bond together with R_2 .

In another preferred embodiment, the present invention is concerned with compounds of formula I wherein R_1 is hydrogen.

10

In another preferred embodiment, the present invention is concerned with compounds of formula I wherein R_2 is hydrogen or R_2 forms a bond together with R_1 .

- 15 In another preferred embodiment, the present invention is concerned with compounds of formula I wherein R_2 is hydrogen.

In another preferred embodiment, the present invention is concerned with compounds of formula I wherein R_3 is C_{1-6} -alkyl or aralkyl.

- 20 In another preferred embodiment, the present invention is concerned with compounds of formula I wherein R_3 is C_{1-6} -alkyl.

In another preferred embodiment, the present invention is concerned with compounds of formula I wherein R_4 is hydrogen, C_{1-3} -alkyl.

25

In another preferred embodiment, the present invention is concerned with compounds of formula I wherein R_4 is hydrogen.

- 30 In another preferred embodiment, the present invention is concerned with compounds of formula I wherein n is 1.

In another preferred embodiment, the present invention is concerned with compounds of formula I wherein m is 1.

In another preferred embodiment, the present invention is concerned with compounds of formula I wherein alkyl is methyl, ethyl, n-propyl, iso-propyl, butyl, tert-butyl, pentyl, hexyl, cyclopropyl or cyclopentyl.

- 5 In another preferred embodiment, the present invention is concerned with compounds of formula I wherein alkenyl is vinyl or 1-propenyl.

In another preferred embodiment, the present invention is concerned with compounds of formula I wherein alkynyl is ethynyl, 1-propynyl and 2-propynyl.

10

In another preferred embodiment, the present invention is concerned with compounds of formula I wherein alkoxy is methoxy, ethoxy, propoxy, butoxy, pentoxy, hexoxy, isopropoxy or cyclopentyloxy.

- 15 In another preferred embodiment, the present invention is concerned with compounds of formula I wherein alkylthio is methylthio, ethylthio, propylthio or cyclopentylthio.

In another preferred embodiment, the present invention is concerned with compounds of formula I wherein aryl is phenyl optionally substituted with halogen.

20

In another preferred embodiment, the present invention is concerned with compounds of formula I wherein arylene is phenylene optionally substituted with halogen.

- 25 In another preferred embodiment, the present invention is concerned with compounds of formula I wherein halogen is fluorine or chlorine.

In another preferred embodiment, the present invention is concerned with compounds of formula I wherein perhalomethyl is trifluoromethyl.

- 30 In another preferred embodiment, the present invention is concerned with compounds of formula I wherein acyl is acetyl.

In another preferred embodiment, the present invention is concerned with compounds of formula I wherein heteroaryl is furan, thiophene, pyrrole, imidazole, pyrazole, pyridine, quinoline, isoquinoline, quinazoline, quinoxaline, indole, benzimidazole or benzofuran.

5 In another preferred embodiment, the present invention is concerned with compounds of formula I wherein heteroaryl is furan, pyrrole, indole or benzofuran.

In another preferred embodiment, the present invention is concerned with compounds of formula I wherein heteroarylene is furan, thiophene, pyrrole, imidazole, pyrazole, triazole, pyridine, pyrazine, pyrimidine or pyridazine.

10

In another preferred embodiment, the present invention is concerned with compounds of formula I wherein aralkyl is benzyl or phenethyl.

15 In another preferred embodiment, the present invention is concerned with compounds of formula I wherein aryloxy is phenoxy.

In another preferred embodiment, the present invention is concerned with compounds of formula I wherein aralkoxy is benzyloxy.

20 In another preferred embodiment, the present invention is concerned with compounds of formula I wherein n is an integer ranging from 1 to 3 and m is 1.

In another preferred embodiment, the present invention is concerned with compounds of formula I wherein the substituents Z and Y are arranged in a trans-configuration.

25 In another preferred embodiment, the present invention is concerned with compounds of formula I wherein the substituents Z and Y are arranged in a cis-configuration.

Preferred compounds of the invention are:

30 (E)-(S)-Ethyl 3-[4-(3-biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-ethoxy-propionate,
(E)-(S)-3-[4-(3-Biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-ethoxy-propionic acid,
(E)-(S)-3-[4-[3-(4'-Bromo-biphenyl-4-yl)-but-2-enyloxy]-phenyl]-2-ethoxy-propionic acid ethyl ester,
(E)-(S)-3-[4-[3-(4'-Bromo-biphenyl-4-yl)-but-2-enyloxy]-phenyl]-2-ethoxy-propionic acid,

- (E)-(S)-2-Ethoxy-3-{4-[3-(4-phenoxy-phenyl)-but-2-enyloxy]-phenyl}-propionic acid ethyl ester,
- (E)-(S)-2-Ethoxy-3-{4-[3-(4-phenoxy-phenyl)-but-2-enyloxy]-phenyl}-propionic acid,
- (E)-(S)-2-Ethoxy-3-(4-{3-[4-(4-methoxy-phenoxy)-phenyl]-but-2-enyloxy}-phenyl)-propionic acid ethyl ester,
- 5 (E)-(S)-2-Ethoxy-3-(4-{3-[4-(4-methoxy-phenoxy)-phenyl]-but-2-enyloxy}-phenyl)-propionic acid,
- (E)-(S)-2-Ethoxy-3-{4-[3-(9H-fluoren-2-yl)-but-2-enyloxy]-phenyl}-propionic acid ethyl ester,
- (E)-(S)-2-Ethoxy-3-{4-[3-(9H-fluoren-2-yl)-but-2-enyloxy]-phenyl}-propionic acid,
- 10 (E)-(S)-3-{4-[3-(3,4-Dimethoxy-phenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester,
- (E)-(S)-3-{4-[3-(3,4-Dimethoxy-phenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid,
- (E)-(S)-3-{4-[3-(3,5-Bis-trifluoromethyl-phenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester,
- 15 (E)-(S)-3-{4-[3-(3,5-Bis-trifluoromethyl-phenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid,
- (E)-(S)-3-{4-(3-Biphenyl-4-yl-allyloxy)-phenyl}-2-ethoxy-propionic acid ethyl ester,
- (E)-(S)-3-{4-(3-Biphenyl-4-yl-allyloxy)-phenyl}-2-ethoxy-propionic acid,
- (E)-(S)-2-Ethoxy-3-[4-(3-naphthalen-2-yl-but-2-enyloxy)-phenyl]-propionic acid ethyl ester,
- 20 (E)-(S)-2-Ethoxy-3-[4-(3-naphthalen-2-yl-but-2-enyloxy)-phenyl]-propionic acid,
- (E)-(S)-2-Ethoxy-3-[4-(3-pyridin-2-yl-but-2-enyloxy)-phenyl]-propionic acid ethyl ester,
- (E)-(S)-2-Ethoxy-3-[4-(3-pyridin-2-yl-but-2-enyloxy)-phenyl]-propionic acid,
- (E)-(S)-3-{4-[3-(3,5-Bis-benzyloxy-phenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester,
- 25 (E)-(S)-3-{4-[3-(3,5-Bis-benzyloxy-phenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid,
- (E)-(S)-2-Ethoxy-3-[4-(3-naphthalen-2-yl-allyloxy)-phenyl]-propionic acid ethyl ester,
- (E)-(S)-2-Ethoxy-3-[4-(3-naphthalen-2-yl-allyloxy)-phenyl]-propionic acid,
- (E)-(S)-2-Ethoxy-3-{4-[3-(3-phenoxy-phenyl)-allyloxy]-phenyl}-propionic acid ethyl ester,
- (E)-(S)-2-Ethoxy-3-{4-[3-(3-phenoxy-phenyl)-allyloxy]-phenyl}-propionic acid,
- 30 (S)-3-[4-(2-Benzofuran-3-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid ethyl ester,
- (S)-3-[4-(2-Benzofuran-3-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid;

or a salt thereof with a pharmaceutically acceptable acid or base, or any optical isomer or mixture of optical isomers, including a racemic mixture, or any tautomeric forms.

Also preferred compounds of the invention are:

- (E)-(S)-3-{4-[3-(4-Benzyloxy-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester,
5 (E)-(S)-3-{4-[3-(4-Benzyloxy-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid,
(E)-(S)-3-{4-[3-(4-Benzo[1,3]dioxol-5-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid ethyl ester,
(E)-(S)-3-{4-[3-(4-Benzo[1,3]dioxol-5-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid,
(E)-(S)-3-{4-[3-(4-Allyloxy-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester,
(E)-(S)-3-{4-[3-(4-Allyloxy-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid,
10 (E)-(S)-3-{4-[3-(4-Benzofuran-7-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid ethyl ester,
(E)-(S)-3-{4-[3-(4-Benzofuran-7-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid,
(S)-3-{4-[3-(4-Benzo[1,3]dioxol-4-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid ethyl ester,
(S)-3-{4-[3-(4-Benzo[1,3]dioxol-4-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid,
(E)-(S)-2-Ethoxy-3-{4-[3-(9H-fluoren-2-yl)-allyloxy]-phenyl}-propionic acid ethyl ester,
15 (E)-(S)-2-Ethoxy-3-{4-[3-(9H-fluoren-2-yl)-allyloxy]-phenyl}-propionic acid,
(S)-2-Ethoxy-3-{4-[3-(9H-fluoren-2-yl)-allyloxy]-phenyl}-propionic acid ethyl ester,
(S)-2-Ethoxy-3-{4-[3-(9H-fluoren-2-yl)-allyloxy]-phenyl}-propionic acid,
(E)-(S)-3-{4-[3-(3,5-Bis-benzyloxy-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester,
20 (E)-(S)-3-{4-[3-(3,5-Bis-benzyloxy-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid,
(E)-(S)-3-{4-[3-(3,5-Dimethoxy-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester,
(E)-(S)-3-{4-[3-(3,5-Dimethoxy-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid,
(E)-(S)-2-Ethoxy-3-{4-[3-(3-phenanthren-9-yl-allyloxy)-phenyl]-propionic acid ethyl ester,
(E)-(S)-2-Ethoxy-3-{4-[3-(2-methoxy-naphthalen-1-yl)-allyloxy]-phenyl}-propionic acid ethyl
25 ester,
(E)-(S)-2-Ethoxy-3-{4-[3-(2-methoxy-naphthalen-1-yl)-allyloxy]-phenyl}-propionic acid,
(E)-(S)-Ethyl 3-{4-[3-(4-Bromophenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate,
(E)-(S)-3-{4-[3-(4-Bromophenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid,
(E)-(S)-Ethyl 3-{4-[3-(4'-Chloro-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate,
30 (E)-(S)-3-{4-[3-(4'-Chloro-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid,
(E)-(S)-Ethyl 2-Ethoxy-3-{4-[3-(5'-isopropyl-2'-methoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-
propionate,
(E)-(S)-2-Ethoxy-3-{4-[3-(5'-isopropyl-2'-methoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-
propionic acid,

- (E)-(S)-Ethyl 2-Ethoxy-3-{4-[3-(5'-chloro-2'-methoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionate,
 (E)-(S)-3-{4-[3-(5'-Chloro-2'-methoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid,
 5 (E)-(S)-Ethyl 3-{4-[3-(2',3'-Dichloro-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate,
 (E)-(S)-3-{4-[3-(2',3'-Dichloro-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid,
 (E)-(S)-Ethyl 3-{4-[3-(2',6'-Dimethoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate,
 10 (E)-(S)-3-{4-[3-(2',6'-Dimethoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid,
 (E)-(S)-Ethyl 3-{4-[3-(4-Bromophenyl)-2-methyl-but-2-enyloxy]-phenyl}-2-ethoxy-propionate,
 (E)-(S)-3-{4-[3-(4-Bromophenyl)-2-methyl-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid,
 (Z)-(S)-Ethyl 3-{4-[3-(4-Bromophenyl)-2-methyl-but-2-enyloxy]-phenyl}-2-ethoxy-propionate,
 15 (Z)-(S)-3-{4-[3-(4-Bromophenyl)-2-methyl-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid,
 (E)-(S)-Ethyl 2-Ethoxy-3-{4-(3-[1,1';3',1'']terphenyl-4''-yl)-but-2-enyloxy}-phenyl}-propionate,
 (E)-(S)-2-Ethoxy-3-{4-(3-[1,1';3',1'']terphenyl-4''-yl)-but-2-enyloxy}-phenyl}-propionic acid,
 (E)-(S)-Ethyl 2-Ethoxy-3-{4-[3-(3'-methyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionate,
 (E)-(S)-2-Ethoxy-3-{4-[3-(3'-methyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionic acid,
 20 (E)-(S)-3-{4-[3-(3'-Acetyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate,
 (E)-(S)-2-Ethoxy-3-{4-[3-(5'-isopropyl-2'-methoxy-biphenyl-4-yl)-allyloxy]-phenyl}-propionic acid ethyl ester,
 (E)-(S)-2-Ethoxy-3-{4-[3-(5'-isopropyl-2'-methoxy-biphenyl-4-yl)-allyloxy]-phenyl}-propionic acid,
 25 (E)-(S)-3-{4-[3-(3,5-Distyryl-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester,
 (E)-(S)-3-{4-[3-(3,5-Distyryl-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid;

or a salt thereof with a pharmaceutically acceptable acid or base, or any optical isomer or mixture of optical isomers, including a racemic mixture, or any tautomeric forms.

30

Also preferred compounds of the invention are:

- (E)-(S)-3-{4-[3-(3,5-Diisopropoxy-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester,
 35 (E)-(S)-3-{4-[3-(3,5-Diisopropoxy-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid,

- (S)-3-{4-[3-(3-Bromo-5-styryl-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester,
(S)-3-{4-[3-(3-Bromo-5-styryl-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid,
5 (E)-(S)-2-Ethoxy-3-[4-(3-phenyl-allyloxy)-phenyl]-propionic acid ethyl ester,
(E)-(S)-2-Ethoxy-3-[4-(3-phenyl-allyloxy)-phenyl]-propionic acid,
10 (E)-(S)-3-{4-[3-(2',3'-Dichloro-biphenyl-4-yl)-allyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester,
(E)-(S)-3-{4-[3-(2',3'-Dichloro-biphenyl-4-yl)-allyloxy]-phenyl}-2-ethoxy-propionic acid,
15 (E)-(S)-3-{4-[3-(3,5-Bis-phenylethynyl-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester,
(E)-(S)-3-{4-[3-(3,5-Bis-phenylethynyl-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid,
20 (E)-(S)-3-{4-[3-(3,5-Diphenethyl-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester,
(E)-(S)-3-{4-[3-(3,5-Diphenethyl-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid,
3-4-[3-(3,5-Bis-cyclopentyloxy-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester,
25 (E)-(S)-3-{4-[3-(3,5-Bis-cyclopentyloxy-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid,
(E)-(S)-3-{4-[3-(3,5-Bis-(2,2,2-trifluoro-ethoxy)-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester,
30 (E)-(S)-3-{4-[3-(3,5-Bis-(2,2,2-trifluoro-ethoxy)-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid,
(E)-(S)-Ethyl 2-Ethoxy-3-{4-[3-(4-furan-2-yl-phenyl)-but-2-enyloxy]-phenyl}-propionate,
35 (E)-(S)-Ethyl 2-Ethoxy-3-{4-[3-(2'-methyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionate,
(E)-(S)-2-Ethoxy-3-{4-[3-(2'-methyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionic acid,
40 (E)-(S)-Ethyl 3-{4-[3-(2',5'-Dimethoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate,
(E)-(S)-3-{4-[3-(2',5'-Dimethoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid,
45 (E)-(S)-Ethyl 3-{4-[3-(4-Bromophenyl)-2-ethyl-but-2-enyloxy]-phenyl}-2-ethoxy-propionate,
(E)-(S)-3-{4-[3-(4-Bromophenyl)-2-ethyl-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid,
50 (Z)-(S)-Ethyl 3-{4-[3-(4-Bromophenyl)-2-ethyl-but-2-enyloxy]-phenyl}-2-ethoxy-propionate,

- (Z)-(S)-3-{4-[3-(4-Bromophenyl)-2-ethyl-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid,
- (E)-(S)-Ethyl 3-{4-[3-(4'-tert-Butyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate,
- 5 (E)-(S)-Ethyl 3-{4-[3-(3',5'-Bis-trifluoromethyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate,
- (E)-(S)-3-{4-[3-(3',5'-Bis-trifluoromethyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid,
- 10 (E)-(S)-Ethyl 2-Ethoxy-3-{4-[3-(4'-isopropyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionate,
- (E)-(S)-2-Ethoxy-3-{4-[3-(4'-isopropyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionic acid,
- 15 (E)-(S)- 3-{4-[3-(3,5-Dimethoxy-phenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester,
- (E)-(S)-3-{4-[3-(3'-Acetyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid,
- 20 (E)-(S)-Ethyl 3-{4-[3-(4'-Acetyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate,
- (E)-(S)-3-{4-[3-(4'-Acetyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid,
- 25 (E)-(S)-Ethyl 2-Ethoxy-3-[4-(3-[1,1';3',1'']terphenyl-5'-yl-allyloxy)-phenyl]-propionate,
- (E)-(S)-2-Ethoxy-3-[4-(3-[1,1';3',1'']terphenyl-5'-yl-allyloxy)-phenyl]-propionic acid,
- (E,E)-(S)-Ethyl 3-(4'-{3-[4-(2-Ethoxy-2-ethoxycarbonyl-ethyl)-phenoxy]-1-methyl-propenyl}-biphenyl-3-yl)-but-2-enoate,
- 30 (E,E)-(S)- 3-(4'-{3-[4-(2-Carboxy-2-ethoxy-ethyl)-phenoxy]-1-methyl-propenyl}-biphenyl-3-yl)-but-2-enoic acid,
- 35 (E)-(S)-Ethyl 2-Ethoxy-3-{4-[3-(3'-methoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionate,
- (E)-(S)-2-Ethoxy-3-{4-[3-(3'-methoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionic acid,
- (E)-(S,S/R)-Ethyl 2-Ethoxy-3-(4-{3-[3'-(1-hydroxy-ethyl)-biphenyl-4-yl]-but-2-enyloxy}-phenyl)-propionate,
- 40 (E)-(S,S/R)-2-Ethoxy-3-(4-{3-[3'-(1-hydroxy-ethyl)-biphenyl-4-yl]-but-2-enyloxy}-phenyl)-propionic acid,
- 45 (E)-(S)-Ethyl 3-{4-[3-(3,5-Dibromophenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate,
- (E)-(S)-3-{4-[3-(3,5-Dibromophenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid,
- (E)-(S)-Ethyl 3-{4-[3-(3,5-Dibromophenyl)-allyloxy]-phenyl}-2-ethoxy-propionate,
- 50 (E)-(S)-3-{4-[3-(3,5-Dibromophenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid,

- (E)-(S)-Ethyl 3-{4-[3-(4,4''-Di-*tert*-butyl-[1,1';3',1'']terphenyl-5'-yl)-allyloxy]-phenyl}-2-ethoxy-propionate,
- 5 (E)-(S)-3-{4-[3-(4,4''-Di-*tert*-butyl-[1,1';3',1'']terphenyl-5'-yl)-allyloxy]-phenyl}-2-ethoxy-propionic acid,
- (E)-(S)-Ethyl 3-{4-[3-(3',5'-Dibromo-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate,
- 10 (E)-(S)-3-{4-[3-(3',5'-Dibromo-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid,
- (E)-(S)-Ethyl 3-{4-[3-(3',5'-Dichloro-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate,
- 15 (E)-(S)-3-{4-[3-(3',5'-Dichloro-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid,
- (E)-(S)-Ethyl 3-{4-[3-(3',5'-Dichloro-biphenyl-4-yl)-allyloxy]-phenyl}-2-ethoxy-propionate,
- 20 (E)-(S)-3-{4-[3-(3',5'-Dichloro-biphenyl-4-yl)-allyloxy]-phenyl}-2-ethoxy-propionic acid,
- (E)-(S)-Ethyl 3-{4-[3-(3',5'-di-*tert*-butyl-biphenyl-4-yl)-allyloxy]-phenyl}-2-ethoxy-propionate,
- (E)-(S)-3-{4-[3-(3',5'-Di-*tert*-butyl-biphenyl-4-yl)-allyloxy]-phenyl}-2-ethoxy-propionic acid,
- 25 (E)-(S)-Ethyl 3-{4-[3-(3',5'-Di-*tert*-butyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate,
- (E)-(S)-3-{4-[3-(3',5'-Di-*tert*-butyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid,
- 30 (E)-(S)-3-{4-[3-(3',5'-Di-*tert*-butyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid,
- (E)-(S/R)-Ethyl 3-[4-(3-Biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-isopropoxy-propionate,
- (E)-(S/R)-3-[4-(3-Biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-isopropoxy-propionic acid,
- 35 (E)-(S/R)-Ethyl 3-[4-(3-Biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-butoxy-propionate,
- (E)-(S/R)-3-[4-(3-Biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-butoxy-propionic acid,
- 40 (E)-(S/R)-Ethyl 3-[4-(3-Biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-hexyloxy-propionate,
- (E)-(S/R)-3-[4-(3-Biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-hexyloxy-propionic acid,
- (E)-(S/R)-Ethyl 3-[4-(3-Biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-(3-phenyl-propoxy)-propionate,
- 45 (E)-(S/R)-3-[4-(3-Biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-(3-phenyl-propoxy)-propionic acid,
- (E)-(S/R)-Ethyl 3-[4-(3-Biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-(4-phenyl-butoxy)-propionate,
- 50 (E)-(S/R)-3-[4-(3-Biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-(4-phenyl-butoxy)-propionic acid,

(E)-(S/R)-Propyl 3-[4-(3-Biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-propoxy-propionate,

(E)-(S/R)-3-[4-(3-Biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-propoxy-propionic acid,

5 (E)-(S)-3-{4-[3-(3,5-Diethoxyoxy-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester,

(E)-(S)-3-{4-[3-(3,5-Diethoxyoxy-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid,

(E)-(S)-3-{4-[3-(3,5-Bis-trifluoromethyl-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester,

(E)-(S)-3-{4-[3-(3,5-Bis-trifluoromethyl-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid,

10 (E)-(R,S)-3-[4-(3-Biphenyl-4-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid ethyl ester,

(E)-(R,S)-3-[4-(3-Biphenyl-4-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid,

(E)-(S)-4-(3-{3-[4-(2-Ethoxy-2-ethoxycarbonyl-ethyl)-phenoxy]-propenyl}-phenoxy-methyl)-benzoic acid methyl ester,

15 (E)-(S)-4-(3-{3-[4-(2-Carboxy-2-ethoxy-ethyl)-phenoxy]-propenyl}-phenoxy-methyl)-benzoic acid;

or a salt thereof with a pharmaceutically acceptable acid or base, or any optical isomer or mixture of optical isomers, including a racemic mixture, or any tautomeric forms.

20 Also preferred compounds of the invention are:

(E)-(S)-Ethyl 2-Ethoxy-3-{4-[3-(4'-fluoro-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionate,

(E)-(S)-2-Ethoxy-3-{4-[3-(4'-fluoro-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionic acid,

(E)-(S)-Ethyl 2-Ethoxy-3-{4-[3-(4-Iodophenyl)-but-2-enyloxy]-phenyl}-propionate;

25

or a salt thereof with a pharmaceutically acceptable acid or base, or any optical isomer or mixture of optical isomers, including a racemic mixture, or any tautomeric forms.

30 In the above structural formulas and throughout the present specification, the following terms have the indicated meaning:

The term "C₁₋₁₂-alkyl" as used herein, alone or in combination is intended to include those alkyl groups of the designated length in either a linear or branched or cyclic configuration represents e.g. cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl and the like.

35 Typical C₁₋₁₂-alkyl groups include, but are not limited to, methyl, ethyl, n-propyl, iso-propyl,

butyl, iso-butyl, sec-butyl, tert-butyl, pentyl, iso-pentyl, hexyl, iso-hexyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl and the like, especially preferred is methyl, ethyl, n-propyl, iso-propyl, butyl, tert-butyl, pentyl, hexyl, cyclopropyl and cyclopentyl.

- 5 The term "C₂₋₁₂-alkenyl" as used herein, represents an olefinically unsaturated branched or straight group having from 2 to the specified number of carbon atoms and at least one double bond. Examples of such groups include, but are not limited to, vinyl, 1-propenyl, 2-propenyl, allyl, iso-propenyl, 1,3-butadienyl, 1-butenyl, hexenyl, pentenyl and the like, especially preferred is vinyl and 1-propenyl.

10

The term "C₂₋₁₂-alkynyl", as used herein, represent an unsaturated branched or straight group having from 2 to the specified number of carbon atoms and at least one triple bond. Examples of such groups include, but are not limited to ethynyl, 1-propynyl, 2-propynyl, 1-butynyl, 2-butynyl, 1-pentynyl, 2-pentynyl and the like, especially preferred is ethynyl, 1-propynyl and 2-propynyl.

15

The term "C₄₋₁₂-alkenynyl" as used herein, represent an unsaturated branched or straight hydrocarbon group having from 4 to the specified number of carbon atoms and both at least one double bond and at least one triple bond. Examples of such groups include, but are not limited to, 1-penten-4-yne, 3-penten-1-yne, 1,3-hexadiene-5-yne and the like.

20

The term "C₁₋₆-alkoxy" as used herein, alone or in combination is intended to include those C₁₋₆-alkyl groups of the designated length in either a linear or branched or cyclic configuration linked through an ether oxygen having its free valence bond from the ether oxygen. Examples of linear alkoxy groups are methoxy, ethoxy, propoxy, butoxy, pentoxy, hexoxy and the like, especially preferred is methoxy, ethoxy, propoxy, butoxy, pentoxy, hexoxy. Examples of branched alkoxy are isopropoxy, sec-butoxy, tert-butoxy, isopentoxy, isohexoxy and the like, especially preferred is isopropoxy. Examples of cyclic alkoxy are cyclopropyloxy, cyclobutyloxy, cyclopentyloxy, cyclohexyloxy and the like, especially preferred is cyclopentyloxy.

25

30

The term "C₁₋₆-alkylthio" as used herein, alone or in combination, refers to a straight or branched or cyclic monovalent substituent comprising a C₁₋₆-alkyl group linked through a divalent sulfur atom having its free valence bond from the sulfur atom and having 1 to 6 carbon atoms e.g. methylthio, ethylthio, propylthio, butylthio, pentylthio and the like,

especially preferred is methylthio, ethylthio and propylthio. Examples of cyclic alkylthio are cyclopropylthio, cyclobutylthio, cyclopentylthio, cyclohexylthio and the like, especially preferred is cyclopentylthio.

- 5 The term "C₁₋₆-alkylamino" as used herein, alone or in combination, refers to a straight or branched or cyclic monovalent substituent comprising a C₁₋₆-alkyl group linked through amino having a free valence bond from the nitrogen atom e.g. methylamino, ethylamino, propylamino, butylamino, pentylamino and the like. Examples of cyclic alkylamino are cyclopropylamino, cyclobutylamino, cyclopentylamino, cyclohexylamino and the like.

10

The term "arylamino" as used herein, alone or in combination, refers to an aryl as defined herein linked through amino having a free valence bond from the nitrogen atom e.g. phenylamino, naphthylamino and the like.

- 15 The term "C₁₋₆-alkoxyC₁₋₆-alkyl" as used herein, alone or in combination, refers to a C₁₋₆-alkyl as defined herein whereto is attached a C₁₋₆-alkoxy as defined herein, e.g. methoxymethyl, ethoxymethyl, methoxyethyl, ethoxyethyl and the like.

- The term "aryl" is intended to include a bicyclic aromatic ring, such as carbocyclic aromatic rings selected from the group consisting of phenyl and naphthyl, (1-naphthyl or 2-naphthyl), optionally substituted with halogen, amino, hydroxy, C₁₋₆-alkyl, C₁₋₆-alkoxy, carboxy or C₁₋₆-alkylester and the like, especially preferred is halogen.
- 20

- The term "arylene" is intended to include divalent aromatic rings, such as carbocyclic aromatic rings selected from the group consisting of phenylene, naphthylene and the like optionally substituted with halogen, amino, hydroxy, C₁₋₆-alkyl, C₁₋₆-alkoxy, carboxy or C₁₋₆-alkylester and the like.
- 25

- The term "halogen" means fluorine, chlorine, bromine or iodine, especially preferred is fluorine and chlorine.
- 30

The term "perhalomethyl" means trifluoromethyl, trichloromethyl, tribromomethyl or triiodomethyl, especially preferred is trifluoromethyl.

The term "C₁₋₆-dialkylamino" as used herein refers to an amino group wherein the two hydrogen atoms independently are substituted with a straight or branched, saturated hydrocarbon chain having the indicated number of carbon atoms; such as dimethylamino, N-ethyl-N-methylamino, diethylamino, dipropylamino, N-(n-butyl)-N-methylamino, di(n-pentyl)amino and the like.

The term "acyl" as used herein refers to a monovalent substituent comprising a C₁₋₆-alkyl group linked through a carbonyl group; such as e.g. acetyl, propionyl, butyryl, isobutyryl, pivaloyl, valeryl and the like, especially preferred is acetyl.

The term "heteroaryl" as used herein, alone or in combination, refers to a monovalent substituent comprising a 5-6 membered monocyclic aromatic system or a 9-10 membered bicyclic aromatic system containing one or more heteroatoms selected from nitrogen, oxygen and sulfur, e.g. furan, thiophene, pyrrole, imidazole, pyrazole, triazole, pyridine, pyrazine, pyrimidine, pyridazine, isothiazole, isoxazole, oxazole, oxadiazole, thiadiazole, quinoline, isoquinoline, quinazoline, quinoxaline, indole, benzimidazole, benzofuran, pteridine and purine and the like, preferred is furan, thiophene, pyrrole, imidazole, pyrazole, pyridine, quinoline, isoquinoline, quinazoline, quinoxaline, indole, benzimidazole, benzofuran, especially preferred is furan, pyrrole, indole and benzofuran.

The term "heteroarylene" as used herein, alone or in combination, refers to a divalent group comprising a 5-6 membered monocyclic aromatic system or a 9-10 membered bicyclic aromatic system containing one or more heteroatoms selected from nitrogen, oxygen and sulfur, e.g. furan, thiophene, pyrrole, imidazole, pyrazole, triazole, pyridine, pyrazine, pyrimidine, pyridazine, isothiazole, isoxazole, oxazole, oxadiazole, thiadiazole, quinoline, isoquinoline, quinazoline, quinoxaline, indole, benzimidazole, benzofuran, pteridine and purine and the like, especially preferred is furan, thiophene, pyrrole, imidazole, pyrazole, triazole, pyridine, pyrazine, pyrimidine, pyridazine.

The term "heteroaryloxy" as used herein, alone or in combination, refers to a heteroaryl as defined herein linked to an oxygen atom having its free valence bond from the oxygen atom e.g. pyrrole, imidazole, pyrazole, triazole, pyridine, pyrazine, pyrimidine, pyridazine, isothiazole, isoxazole, oxazole, oxadiazole, thiadiazole, quinoline, isoquinoline, quinazoline,

quinoxaline, indole, benzimidazole, benzofuran, pteridine and purine linked to oxygen, and the like.

5 The term "aralkyl" as used herein refers to a straight or branched saturated carbon chain containing from 1 to 6 carbons substituted with an aromatic carbohydride; such as benzyl, phenethyl, 3-phenylpropyl, 1-naphthylmethyl, 2-(1-naphthyl)ethyl and the like, especially preferred is benzyl and phenethyl.

10 The term "aryloxy" as used herein refers to phenoxy, 1-naphthyloxy, 2-naphthyloxy and the like, especially preferred is phenoxy.

The term "aralkoxy" as used herein refers to a C₁₋₆-alkoxy group substituted with an aromatic carbohydride, such as benzyloxy, phenethoxy, 3-phenylpropoxy, 1-naphthylmethoxy, 2-(1-naphthyl)ethoxy and the like, especially preferred is benzyloxy.

15 The term "heteroaralkyl" as used herein refers to a straight or branched saturated carbon chain containing from 1 to 6 carbons substituted with a heteroaryl group; such as (2-furyl)methyl, (3-furyl)methyl, (2-thienyl)methyl, (3-thienyl)methyl, (2-pyridyl)methyl, 1-methyl-1-(2-pyrimidyl)ethyl and the like.

20 The term "heteroaralkoxy" as used herein refers to a heteroaralkyl as defined herein linked to an oxygen atom having its free valence bond from the oxygen atom, e.g. (2-furyl)methyl, (3-furyl)methyl, (2-thienyl)methyl, (3-thienyl)methyl, (2-pyridyl)methyl, 1-methyl-1-(2-pyrimidyl)ethyl linked to oxygen, and the like.

25 The term "arylthio" as used herein, alone or in combination, refers to an aryl group linked through a divalent sulfur atom having its free valence bond from the sulfur atom, the aryl group optionally being mono- or polysubstituted with C₁₋₆-alkyl, halogen, hydroxy or C₁₋₆-alkoxy; e.g. phenylthio, (4-methylphenyl)-thio, (2-chlorophenyl)thio and the like.

30 As used herein, the phrase "heterocyclyl" means a monovalent saturated or unsaturated non aromatic group being monocyclic and containing one or more, such as from one to four carbon atom(s), and from one to four N, O or S atom(s) or a combination thereof. The phrase "heterocyclyl" includes, but is not limited to, 5-membered heterocycles having one hetero

atom (e.g. pyrrolidine, pyrroline and the like); 5-membered heterocycles having two heteroatoms in 1,2 or 1,3 positions (e.g. pyrazoline, pyrazolidine, 1,2-oxathiolane, imidazolidine, imidazoline, 4-oxazolone and the like); 5-membered heterocycles having three heteroatoms (e.g. tetrahydrofuran and the like); 5-membered heterocycles having four heteroatoms; 6-membered heterocycles with one heteroatom (e.g. piperidine and the like); 6-membered heterocycles with two heteroatoms (e.g. piperazine, morpholine and the like); 6-membered heterocycles with three heteroatoms; and 6-membered heterocycles with four heteroatoms, and the like.

10 As used herein, the phrase "a divalent heterocyclic group" means a divalent saturated or unsaturated system being monocyclic and containing one or more, such as from one to four carbon atom(s), and one to four N, O or S atom(s) or a combination thereof. The phrase a divalent heterocyclic group includes, but is not limited to, 5-membered heterocycles having one hetero atom (e.g. pyrrolidine, pyrroline and the like); 5-membered heterocycles having
15 two heteroatoms in 1,2 or 1,3 positions (e.g. pyrazoline, pyrazolidine, 1,2-oxathiolane, imidazolidine, imidazoline, 4-oxazolone and the like); 5-membered heterocycles having three heteroatoms (e.g. tetrahydrofuran and the like); 5-membered heterocycles having four heteroatoms; 6-membered heterocycles with one heteroatom (e.g. piperidine and the like); 6-membered heterocycles with two heteroatoms (e.g. piperazine, morpholine and the like);
20 membered heterocycles with three heteroatoms; and 6-membered heterocycles with four heteroatoms, and the like.

As used herein the term "treatment" includes treatment, prevention and management of such condition.

25

Certain of the above defined terms may occur more than once in the above formula (I), and upon such occurrence each term shall be defined independently of the other.

The present invention also encompasses pharmaceutically acceptable salts of the present
30 compounds. Such salts include pharmaceutically acceptable acid addition salts, pharmaceutically acceptable base addition salts, pharmaceutically acceptable metal salts, ammonium and alkylated ammonium salts. Acid addition salts include salts of inorganic acids as well as organic acids. Representative examples of suitable inorganic acids include hydrochloric, hydrobromic, hydroiodic, phosphoric, sulfuric, nitric acids and the like. Representative exam-

- ples of suitable organic acids include formic, acetic, trichloroacetic, trifluoroacetic, propionic, benzoic, cinnamic, citric, fumaric, glycolic, lactic, maleic, malic, malonic, mandelic, oxalic, picric, pyruvic, salicylic, succinic, methanesulfonic, ethanesulfonic, tartaric, ascorbic, pamoic, bismethylene salicylic, ethanedisulfonic, gluconic, citraconic, aspartic, stearic, palmitic,
- 5 EDTA, glycolic, p-aminobenzoic, glutamic, benzenesulfonic, p-toluenesulfonic acids, sulphates, nitrates, phosphates, perchlorates, borates, acetates, benzoates, hydroxynaphthoates, glycerophosphates, ketoglutarates and the like. Further examples of pharmaceutically acceptable inorganic or organic acid addition salts include the pharmaceutically acceptable salts listed in J. Pharm. Sci. 1977, 66, 2, which is incorporated herein by reference. Exam-
- 10 ples of metal salts include lithium, sodium, potassium, magnesium salts and the like. Examples of ammonium and alkylated ammonium salts include ammonium, methylammonium, dimethylammonium, trimethylammonium, ethylammonium, hydroxyethylammonium, diethylammonium, butylammonium, tetramethylammonium salts and the like. Examples of organic bases include lysine, arginine, guanidine, diethanolamine, choline and the like.
- 15
- The pharmaceutically acceptable salts are prepared by reacting the compound of formula I with 1 to 4 equivalents of a base such as sodium hydroxide, sodium methoxide, sodium hydride, potassium t-butoxide, calcium hydroxide, magnesium hydroxide and the like, in sol-
- 20 vents like ether, THF, methanol, t-butanol, dioxane, isopropanol, ethanol etc. Mixture of solvents may be used. Organic bases like lysine, arginine, diethanolamine, choline, guanidine and their derivatives etc. may also be used. Alternatively, acid addition salts wherever applicable are prepared by treatment with acids such as hydrochloric acid, hydrobromic acid, nitric acid, dimethylacetic acid, phosphoric acid, p-toluenesulphonic acid, methanesulfonic acid, acetic acid, citric acid, maleic acid salicylic acid, hydroxynaphthoic acid, ascorbic acid, pal-
- 25 mitic acid, succinic acid, benzoic acid, benzenesulfonic acid, tartaric acid and the like in solvents like ethyl acetate, ether, alcohols, acetone, THF, dioxane etc. Mixture of solvents may also be used.
- 30
- The stereoisomers of the compounds forming part of this invention may be prepared by using reactants in their single enantiomeric form in the process wherever possible or by conducting the reaction in the presence of reagents or catalysts in their single enantiomer form or by resolving the mixture of stereoisomers by conventional methods. Some of the preferred methods include use of microbial resolution, enzymatic resolution, resolving the diastereomeric salts formed with chiral acids such as mandelic acid, camphorsulfonic acid, tartaric acid, lac-

tic acid, and the like wherever applicable or chiral bases such as brucine, (R)- or (S)-phenylethylamine, cinchona alkaloids and their derivatives and the like. Commonly used methods are compiled by Jaques et al in "Enantiomers, Racemates and Resolution" (Wiley Interscience, 1981). More specifically the compound of formula I may be converted to a 1:1 mixture of diastereomeric amides by treating with chiral amines, aminoacids, aminoalcohols derived from aminoacids; conventional reaction conditions may be employed to convert acid into an amide; the dia-stereomers may be separated either by fractional crystallization or chromatography and the stereoisomers of compound of formula I may be prepared by hydrolysing the pure diastereomeric amide.

10

Various polymorphs of compound of general formula I forming part of this invention may be prepared by crystallization of compound of formula I under different conditions. For example, using different solvents commonly used or their mixtures for recrystallization; crystallizations at different temperatures; various modes of cooling, ranging from very fast to very slow cooling during crystallizations. Polymorphs may also be obtained by heating or melting the compound followed by gradual or fast cooling. The presence of polymorphs may be determined by solid probe nmr spectroscopy, ir spectroscopy, differential scanning calorimetry, powder X-ray diffraction or such other techniques.

15

20 The invention also encompasses prodrugs of the present compounds, which on administration undergo chemical conversion by metabolic processes before becoming active pharmacological substances. In general, such prodrugs will be functional derivatives of the present compounds, which are readily convertible in vivo into the required compound of the formula (I). Conventional procedures for the selection and preparation of suitable prodrug derivatives are described, for example, in "Design of Prodrugs", ed. H. Bundgaard, Elsevier, 1985.

25

The invention also encompasses active metabolites of the present compounds.

30 Furthermore, the present compounds of formula I can be utilised in the treatment and/or prevention of conditions mediated by nuclear receptors, in particular the Peroxisome Proliferator-Activated Receptors (PPAR).

In a further aspect, the present invention relates to a method of treating and/or preventing Type I or Type II diabetes.

5 In a still further aspect, the present invention relates to the use of one or more compounds of the general formula I or pharmaceutically acceptable salts thereof for the preparation of a medicament for the treatment and/or prevention of Type I or Type II diabetes.

10 In a still further aspect, the present compounds are useful for the treatment and/or prevention of IGT.

In a still further aspect, the present compounds are useful for the treatment and/or prevention of Type 2 diabetes.

15 In a still further aspect, the present compounds are useful for the delaying or prevention of the progression from IGT to Type 2 diabetes.

In a still further aspect, the present compounds are useful for the delaying or prevention of the progression from non-insulin requiring Type 2 diabetes to insulin requiring Type 2 diabetes.

20 In another aspect, the present compounds reduce blood glucose and triglyceride levels and are accordingly useful for the treatment and/or prevention of ailments and disorders such as diabetes and/or obesity.

25 In still another aspect, the present compounds are useful for the treatment and/or prophylaxis of insulin resistance (Type 2 diabetes), impaired glucose tolerance, dyslipidemia, disorders related to Syndrome X such as hypertension, obesity, insulin resistance, hyperglycaemia, atherosclerosis, hyperlipidemia, coronary artery disease, myocardial ischemia and other cardiovascular disorders.

30 In still another aspect, the present compounds are effective in decreasing apoptosis in mammalian cells such as beta cells of Islets of Langerhans.

In still another aspect, the present compounds are useful for the treatment of certain renal diseases including glomerulonephritis, glomerulosclerosis, nephrotic syndrome, hypertensive nephrosclerosis.

- 5 In still another aspect, the present compounds may also be useful for improving cognitive functions in dementia, treating diabetic complications, psoriasis, polycystic ovarian syndrome (PCOS) and prevention and treatment of bone loss, e.g. osteoporosis.

10 The invention also relates to pharmaceutical compositions comprising, as an active ingredient, at least one compound of the formula I or any optical or geometric isomer or tautomeric form thereof including mixtures of these or a pharmaceutically acceptable salt thereof together with one or more pharmaceutically acceptable carriers or diluents.

15 Furthermore, the invention relates to the use of compounds of the general formula I or their tautomeric forms, their stereoisomers, their polymorphs, their pharmaceutically acceptable salts or pharmaceutically acceptable solvates thereof for the preparation of a pharmaceutical composition for the treatment and/or prevention of conditions mediated by nuclear receptors, in particular the Peroxisome Proliferator-Activated Receptors (PPAR) such as the conditions mentioned above.


20

The present invention also relates to a process for the preparation of the above said novel compounds, their derivatives, their analogs, their tautomeric forms, their stereoisomers, their polymorphs, their pharmaceutically acceptable salts or pharmaceutically acceptable solvates.

- 25 The method comprises:

a)

Reacting a compound of formula II




30

(II)

wherein A, X₁, X₂ and Y are defined as above, through a Wittig process with e.g. (EtO)₂PO(CHZ)(CH₂)_tCOOR₆ (wherein R₆ is an alkyl group), in the presence of a base such as sodium hydride, EtONa and the like to give a compound of formula III.

5




10

(III)

wherein A, X₁, X₂, Y, Z and R₆ are defined as above, and wherein t is 0-2, and

b)

- 15 reducing a compound of formula III, wherein A, X₁, X₂, Y, Z, R₆ and t are defined as above with a suitable reagent such as diisobutylaluminum hydride, to give a compound of formula IV.




20

(IV)

wherein A, X₁, X₂, Y, Z and t are defined as above, and

5 c)

reacting a compound of IV, wherein A, X₁, X₂, Y, Z and t are defined as above, with a compound of formula V



10


(V)

15 wherein Q, Ar, R₁, R₂, R₃, R₄ and m are defined as above, under Mitsunobu conditions, using a reagent such as triphenylphosphine/diethylazodicarboxylate and the like to obtain a compound of formula I, wherein A, X₁, X₂, Y, Z, Q, Ar, R₁, R₂, R₃, R₄, n and m are defined as above, except that R₄ is not H, n and m are not 0, and

d)

20 converting the -OH functionality in a compound of formula IV wherein A, X₁, X₂, Y, Z and t are defined as above to an appropriate leaving group (L) such as p-toluenesulfonate, methanesulfonate, halogen (in examples by methods according to: Houben-Weyl, Methoden der organischen Chemie, Alkohole III, 6/1b, Thieme Verlag 1984, 4th Ed., pp. 927-939; Comprehensive Organic Transformations. A guide to functional group preparations, VCH Publishers
25 1989, 1st Ed., pp. 353-363), triflate and the like, to give a compound of formula VI

41




(VI)

e)


reacting a compound of formula VI

5



(VI)

wherein L is a leaving group such as p-toluenesulfonate, methanesulfonate, halogen, triflate
 10 and the like and wherein A, X₁, X₂, Y, Z and t are defined as above with a compound of formula V




15

(V)

wherein Q, Ar, R₁, R₂, R₃, R₄ and m are defined as above, to give a compound of formula I
 wherein A, X₁, X₂, Y, Z, Q, Ar, R₁, R₂, R₃, R₄, n and m are defined as above except that R₄ is
 5 not H, n and m are not 0, or

e)


reacting a compound of formula VII



10

(VII)

wherein A, X₁ and X₂ are defined as above, through a Friedel-Crafts acylation, with in exam-
 ple ClOCCHZ(CH₂)_nR₇ (wherein n and Z is defined as above and R₇ are halogen or OH), in
 15 the presence of a Friedel-Crafts catalysts such as aluminium trichloride and the like, to give a
 compound of formula VIII




20

(VIII)

wherein A, X₁, X₂, Z, R₇ and n are defined as above, and

f)

- 5 reacting a compound of formula VIII, wherein A, X₁, X₂, Z and R₇ are defined as above with a Grignard reagents such a MgBrY or a lithium reagent such as LiY or organozinc reagent such as ZnY, wherein Y is defined as above, followed by a acidic workup to give a compound of formula IX




(IX)

- 15 wherein A, X₁, X₂, Z, Y, R₇ and n are defined as above, and

g)

reacting a compound of IX, wherein A, X₁, X₂, Z, Y, R₇ and n are defined as above, with a compound of formula V




(V)

wherein Q, Ar, R₁, R₂, R₃, R₄ and m are defined as above except that m is not 0, under either basic condition e.g. potassium carbonate/acetone (if R₇ is halogen) or Mitsunobu conditions (if R₇ is OH) using a reagent such as triphenylphosphine/diethylazodicarboxylate and the like to obtain a compound of formula I, wherein A, X₁, X₂, Y, Z, Q, Ar, R₁, R₂, R₃, R₄, n and m are defined as above, except that R₄ is not H, n and m are not 0, or

h)

by chemical or enzymatic saponification of a compound of formula I

10



(I)

wherein A, X₁, X₂, Y, Z, Q, Ar, R₁, R₂, R₃, R₄, n and m are defined as above, except that R₄ is not H, to obtain a compound of formula I, wherein A, X₁, X₂, Y, Z, Q, Ar, R₁, R₂, R₃, R₄, n and m are defined as above, except that R₄ is H.

i)

Trans-cis or cis-trans isomerization of compounds I, III, IV, VI, and IX (Arai et al., Chem. Rev., 93, pp 23-39, 1993; J. March, Advanced Organic Chemistry, 4th Ed., J. Wiley & Sons, New York 1992, pp. 218, 245, 745).

PHARMACOLOGICAL METHODS

In vitro PPAR alpha and PPAR gamma activation activity.

Principle

5

The PPAR gene transcription activation assays were based on transient transfection into human HEK293 cells of two plasmids encoding a chimeric test protein and a reporter protein respectively. The chimeric test protein was a fusion of the DNA binding domain (DBD) from the yeast GAL4 transcription factor to the ligand binding domain (LBD) of the human PPAR proteins. The PPAR LBD harbored in addition to the ligand binding pocket also the native activation domain (activating function 2 = AF2) allowing the fusion protein to function as a PPAR ligand dependent transcription factor. The GAL4 DBD will force the fusion protein to bind only to Gal4 enhancers (of which none existed in HEK293 cells). The reporter plasmid contained a Gal4 enhancer driving the expression of the firefly luciferase protein. After transfection, HEK293 cells expressed the GAL4-DBD-PPAR-LBD fusion protein. The fusion protein will in turn bind to the Gal4 enhancer controlling the luciferase expression, and do nothing in the absence of ligand. Upon addition to the cells of a PPAR ligand, luciferase protein will be produced in amounts corresponding to the activation of the PPAR protein. The amount of luciferase protein is measured by light emission after addition of the appropriate substrate.

20

Methods

25 **In vitro transactivation assays.**

Cell culture and transfection: HEK293 cells were grown in DMEM + 10% FCS. Cells were seeded in 96-well plates the day before transfection to give a confluency of 50-80 % at transfection. A total of 0,8 µg DNA containing 0,64 µg pM1α/γLBD, 0,1 µg pCMVβGal, 0,08 µg pGL2Gal4DBD and 0,02 µg pADVANTAGE was transfected per well using FuGene transfection reagent according to the manufacturers instructions (Roche). Cells were allowed to express protein for 48 h followed by addition of compound.

30

Plasmids: Human PPAR α and γ was obtained by PCR amplification using cDNA synthesized by reverse transcription of mRNA from liver and adipose tissue respectively. Amplified cDNAs were cloned into pCR2.1 and sequenced. The ligand binding domain (LBD) of each PPAR isoform was generated by PCR (PPAR α : aa 167 - C-terminus; PPAR γ : aa 165 - C-terminus) and fused to the DNA binding domain (DBD) of the yeast transcription factor GAL4 by subcloning fragments in frame into the vector pM1 generating the plasmids pM1 α LBD and pM1 γ LBD. Ensuing fusions were verified by sequencing. The reporter was constructed by inserting an oligonucleotide encoding five repeats of the GAL4 recognition sequence (5 x CGGAGTACTGTCCTCCG(AG)) into the vector pGL2 promotor (Promega) generating the plasmid pGL2(GAL4)₅. pCMV β Gal was purchased from Clontech and pADVANTAGE was purchased from Promega.

Luciferase assay: Medium including test compound was aspirated and 100 μ l PBS incl. 1mM Mg⁺⁺ and Ca⁺⁺ was added to each well. The luciferase assay was performed using the LucLite kit according to the manufacturers instructions (Packard Instruments). Light emission was quantified by counting SPC mode on a Packard Instruments top-counter. To measure β -galactosidase activity 25 μ l supernatant from each transfection lysate was transferred to a new microplate. β -galactosidase assays were performed in the microwell plates using a kit from Promega and read in a microplate reader. The β -galactosidase data were used to normalize (transfection efficiency, cell growth etc.) the luciferase data.

Compounds: All compounds were dissolved in DMSO and diluted 1:1000 upon addition to the cells. Compounds were tested in quadruple in five concentrations ranging from 0.01 to 30 μ M. Cells were treated with compound for 24 h followed by luciferase assay. Each compound was tested in three separate experiments. EC₅₀ values were calculated via non-linear regression using GraphPad PRISM 3.02 (GraphPad Software, San Diego, CA). The results were expressed as means.

Table 1.

In vitro PPAR alpha and PPAR gamma activation of examples according to the present invention.

5

Example no	In vitro activation			
	PPAR α		PPAR γ	
	EC ₅₀ , μ M	% max ^a	EC ₅₀ , μ M	% max ^b
4	3.1	212	0.72	156
9	0.038	234	0.35	125
27	0.10	185	0.11	99
57	0.38	178	0.70	110
124	0.35	102	0.30	83
134	2.90	122	0.89	155

Compounds were tested in at least three separate experiments in five concentrations ranging from 0.01 to 30 μ M. EC₅₀'s were not calculated for compounds producing transactivation lower than 25% at 30 μ M. ^aFold activation relative to maximum activation obtained with Wy14643 (approx. 20 fold corresponded to 100%) and with ^brosiglitazone (approx. 120 fold corresponded to 100%).

10

PHARMACEUTICAL COMPOSITIONS

- 15 In another aspect, the present invention includes within its scope pharmaceutical compositions comprising, as an active ingredient, at least one of the compounds of the general formula I or a pharmaceutically acceptable salt thereof together with a pharmaceutically acceptable carrier or diluent.
- 20 The present compounds may also be administered in combination with one or more further pharmacologically active substances eg., selected from antiobesity agents, antidiabetics, antihypertensive agents, agents for the treatment and/or prevention of complications resulting from or associated with diabetes and agents for the treatment and/or prevention of complications and disorders resulting from or associated with obesity.

Thus, in a further aspect of the invention the present compounds may be administered in combination with one or more antiobesity agents or appetite regulating agents.

- 5 Such agents may be selected from the group consisting of CART (cocaine amphetamine regulated transcript) agonists, NPY (neuropeptide Y) antagonists, MC4 (melanocortin 4) agonists, orexin antagonists, TNF (tumor necrosis factor) agonists, CRF (corticotropin releasing factor) agonists, CRF BP (corticotropin releasing factor binding protein) antagonists, urocortin agonists, $\beta 3$ agonists, MSH (melanocyte-stimulating hormone) agonists, MCH
- 10 (melanocyte-concentrating hormone) antagonists, CCK (cholecystokinin) agonists, serotonin re-uptake inhibitors, serotonin and noradrenaline re-uptake inhibitors, mixed serotonin and noradrenergic compounds, 5HT (serotonin) agonists, bombesin agonists, galanin antagonists, growth hormone, growth hormone releasing compounds, TRH (thyrotropin releasing hormone) agonists, UCP 2 or 3 (uncoupling protein 2 or 3) modulators, leptin agonists, DA
- 15 agonists (bromocriptin, doprexin), lipase/amylase inhibitors, RXR (retinoid X receptor) modulators or TR β agonists.

In one embodiment of the invention the antiobesity agent is leptin.

- 20 In another embodiment the antiobesity agent is dexamphetamine or amphetamine.

In another embodiment the antiobesity agent is fenfluramine or dexfenfluramine.

In still another embodiment the antiobesity agent is sibutramine.

25

In a further embodiment the antiobesity agent is orlistat.

In another embodiment the antiobesity agent is mazindol or phentermine.

- 30 Suitable antidiabetics comprise insulin, GLP-1 (glucagons like peptide-1) derivatives such as those disclosed in WO 98/08871 to Novo Nordisk A/S, which is incorporated herein by reference as well as orally active hypoglycaemic agents.

The orally active hypoglycaemic agents preferably comprise sulphonylureas, biguanides, meglitinides, glucosidase inhibitors, glucagon antagonists such as those disclosed in WO 99/01423 to Novo Nordisk A/S and Agouron Pharmaceuticals, Inc., GLP-1 agonists, potassium channel openers such as those disclosed in WO 97/26265 and WO 99/03861 to Novo Nordisk A/S which are incorporated herein by reference, DPP-IV (dipeptidyl peptidase-IV) inhibitors, inhibitors of hepatic enzymes involved in stimulation of gluconeogenesis and/or glycogenolysis, glucose uptake modulators, compounds modifying the lipid metabolism such as antihyperlipidemic agents and antilipidemic agents as HMG CoA inhibitors (statins), compounds lowering food intake, RXR agonists and agents acting on the ATP-dependent potassium channel of the β -cells.

In one embodiment of the invention the present compounds are administered in combination with insulin.

In a further embodiment the present compounds are administered in combination with a sulphonylurea eg. tolbutamide, glibenclamide, glipizide or glicazide.

In another embodiment the present compounds are administered in combination with a biguanide eg. metformin.

In yet another embodiment the present compounds are administered in combination with a meglitinide eg. repaglinide or senaglinide.

In a further embodiment the present compounds are administered in combination with an α -glucosidase inhibitor eg. miglitol or acarbose.

In another embodiment the present compounds are administered in combination with an agent acting on the ATP-dependent potassium channel of the β -cells eg. tolbutamide, glibenclamide, glipizide, glicazide or repaglinide.

Furthermore, the present compounds may be administered in combination with nateglinide.

In still another embodiment the present compounds are administered in combination with an antihyperlipidemic agent or antilipidemic agent eg. cholestyramine, colestipol, clofibrate, gemfibrozil, lovastatin, pravastatin, simvastatin, probucol or dextrothyroxine.

- 5 In a further embodiment the present compounds are administered in combination with more than one of the above-mentioned compounds eg. In combination with a sulphonylurea and metformin, a sulphonylurea and acarbose, repaglinide and metformin, insulin and a sulphonylurea, insulin and metformin, insulin, insulin and lovastatin, etc.
- 10 Furthermore, the present compounds may be administered in combination with one or more antihypertensive agents. Examples of antihypertensive agents are β -blockers such as alprenolol, atenolol, timolol, pindolol, propranolol and metoprolol, ACE (angiotensin converting enzyme) inhibitors such as benazepril, captopril, enalapril, fosinopril, lisinopril, quinapril and ramipril, calcium channel blockers such as nifedipine, felodipine, nicardipine, isradipine, ni-
- 15 modipine, diltiazem and verapamil, and α -blockers such as doxazosin, urapidil, prazosin and terazosin. Further reference can be made to Remington: The Science and Practice of Pharmacy, 19th Edition, Gennaro, Ed., Mack Publishing Co., Easton, PA, 1995.

- It should be understood that any suitable combination of the compounds according to the in-
- 20 vention with one or more of the above-mentioned compounds and optionally one or more further pharmacologically active substances are considered to be within the scope of the present invention.

- Pharmaceutical compositions containing a compound of the present invention may be prepared
- 25 by conventional techniques, e.g. as described in Remington: The Science and Practice of Pharmacy, 19th Ed., 1995. The compositions may appear in conventional forms, for example capsules, tablets, aerosols, solutions, suspensions or topical applications.

- Typical compositions include a compound of formula I or a pharmaceutically acceptable acid
- 30 addition salt thereof, associated with a pharmaceutically acceptable excipient which may be a carrier or a diluent or be diluted by a carrier, or enclosed within a carrier which can be in the form of a capsule, sachet, paper or other container. In making the compositions, conventional techniques for the preparation of pharmaceutical compositions may be used. For example, the active compound will usually be mixed with a carrier, or diluted by a carrier,

or enclosed within a carrier which may be in the form of a ampoule, capsule, sachet, paper, or other container. When the carrier serves as a diluent, it may be solid, semi-solid, or liquid material which acts as a vehicle, excipient, or medium for the active compound. The active compound can be adsorbed on a granular solid container for example in a sachet. Some

5 examples of suitable carriers are water, salt solutions, alcohols, polyethylene glycols, polyhydroxyethoxylated castor oil, peanut oil, olive oil, gelatine, lactose, terra alba, sucrose, cyclodextrin, amylose, magnesium stearate, talc, gelatin, agar, pectin, acacia, stearic acid or lower alkyl ethers of cellulose, silicic acid, fatty acids, fatty acid amines, fatty acid monoglycerides and diglycerides, pentaerythritol fatty acid esters, polyoxyethylene,

10 hydroxymethylcellulose and polyvinylpyrrolidone. Similarly, the carrier or diluent may include any sustained release material known in the art, such as glyceryl monostearate or glyceryl distearate, alone or mixed with a wax. The formulations may also include wetting agents, emulsifying and suspending agents, preserving agents, sweetening agents or flavouring agents. The formulations of the invention may be formulated so as to provide quick,

15 sustained, or delayed release of the active ingredient after administration to the patient by employing procedures well known in the art.

The pharmaceutical compositions can be sterilized and mixed, if desired, with auxiliary agents, emulsifiers, salt for influencing osmotic pressure, buffers and/or colouring substances and the like, which do not deleteriously react with the active compounds.

20

The route of administration may be any route, which effectively transports the active compound to the appropriate or desired site of action, such as oral, nasal, pulmonary, transdermal or parenteral e.g. rectal, depot, subcutaneous, intravenous, intraurethral, intramuscular, intranasal, ophthalmic solution or an ointment, the oral route being preferred.

25

If a solid carrier is used for oral administration, the preparation may be tableted, placed in a hard gelatin capsule in powder or pellet form or it can be in the form of a troche or lozenge. If a liquid carrier is used, the preparation may be in the form of a syrup, emulsion, soft gelatin capsule or sterile injectable liquid such as an aqueous or non-aqueous liquid suspension or solution.

30

For nasal administration, the preparation may contain a compound of formula I dissolved or suspended in a liquid carrier, in particular an aqueous carrier, for aerosol application. The carrier

may contain additives such as solubilizing agents, e.g. propylene glycol, surfactants, absorption enhancers such as lecithin (phosphatidylcholine) or cyclodextrin, or preservatives such as parabenes.

- 5 For parenteral application, particularly suitable are injectable solutions or suspensions, preferably aqueous solutions with the active compound dissolved in polyhydroxylated castor oil.

Tablets, dragees, or capsules having talc and/or a carbohydrate carrier or binder or the like are particularly suitable for oral application. Preferable carriers for tablets, dragees, or capsules include lactose, corn starch, and/or potato starch. A syrup or elixir can be used in cases where a sweetened vehicle can be employed.

A typical tablet which may be prepared by conventional tableting techniques may contain:

15 Core:

Active compound (as free compound or salt thereof)	5 mg
Colloidal silicon dioxide (Aerosil)	1.5 mg
Cellulose, microcryst. (Avicel)	70 mg
Modified cellulose gum (Ac-Di-Sol)	7.5 mg
20 Magnesium stearate	Ad.

Coating:

HPMC approx.	9 mg
*Mywacett 9-40 T approx.	0.9 mg

25

*Acylated monoglyceride used as plasticizer for film coating.

The compounds of the invention may be administered to a mammal, especially a human in need of such treatment, prevention, elimination, alleviation or amelioration of diseases related to the regulation of blood sugar.

30 Such mammals include also animals, both domestic animals, e.g. household pets, and non-domestic animals such as wildlife.

The compounds of the invention are effective over a wide dosage range. For example, in the treatment of adult humans, dosages from about 0.05 to about 100 mg, preferably from about 0.1 to about 100 mg, per day may be used. A most preferable dosage is about 0.1 mg to about 70 mg per day. In choosing a regimen for patients it may frequently be necessary to
5 begin with a dosage of from about 2 to about 70 mg per day and when the condition is under control to reduce the dosage as low as from about 0.1 to about 10 mg per day. The exact dosage will depend upon the mode of administration, on the therapy desired, form in which administered, the subject to be treated and the body weight of the subject to be treated, and the preference and experience of the physician or veterinarian in charge.

10 Generally, the compounds of the present invention are dispensed in unit dosage form comprising from about 0.1 to about 100 mg of active ingredient together with a pharmaceutically acceptable carrier per unit dosage.

15 Usually, dosage forms suitable for oral, nasal, pulmonary or transdermal administration comprise from about 0.001 mg to about 100 mg, preferably from about 0.01 mg to about 50 mg of the compounds of formula I admixed with a pharmaceutically acceptable carrier or diluent.

Any novel feature or combination of features described herein is considered essential to this
20 invention.

EXAMPLES


The process for preparing compounds of formula I, and preparations containing them, is
25 further illustrated in the following examples, which however, are not to be construed as limiting.

The structures of the compounds are confirmed by either elemental analysis (MA) nuclear magnetic resonance (NMR) or mass spectrometry (MS). NMR shifts (δ) are given in parts per million (ppm) and only selected peaks are given. Mp is melting point and is given in °C. Col-
30 umn chromatography was carried out using the technique described by W.C. Still et al, J. Org. Chem. 1978, 43, 2923-2925 on Merck silica gel 60 (Art 9385). Compounds used as starting materials are either known compounds or compounds which can readily be prepared by methods known per se.

Abbreviations:

	THF:	tetrahydrofuran
	DIBAL-H:	diisobutylaluminum hydride
	Na ₂ SO ₄ :	sodium sulfate
5	MgSO ₄ :	magnesium sulfate
	DMSO:	dimethylsulfoxide
	CDCl ₃ :	deuterated chloroform
	DMF:	N,N-dimethylformamide
	HCl:	hydrochloric acid
10	DME:	1,2-dimethoxyethane
	min:	minutes
	h:	hours

15

EXAMPLE 1

(*E*)-(*S*)-Ethyl 3-[4-(3-biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-ethoxy-propionate

20

a)

Sodium (1.75 g, 73.4 mmol) was added to ethanol (45 ml) at 20°C and the mixture stirred until the metal had fully reacted. Triethyl phosphonoacetate (14.69 g, 73.4 mmol) was added, the mixture stirred for 5 min., then 4-acetylbiphenyl (12.00 g, 61.1 mmol) was added to the stirred solution. The mixture was stirred at room temperature for 24h, the resulting suspension filtered, and the filter-cake collected and recrystallised from ethanol to give (*E*)-3-biphenyl-4-yl-but-2-enoic acid ethyl ester as white crystals; 5.73 g (36%)

25

¹H NMR (300 MHz, CDCl₃) δ: 1.32 (3H, t), 2.62 (3H, d), 4.21 (2H, q), 6.2 (1H, d), 7.31-7.65 (9H, m). MS: 267 (M⁺), 266(100%), 221, 194, 178.

30 Microanalysis Calculated % C: 81.00, H: 7.0. Found C: 80.86, H: 6.90.

b)

A 1M solution of DIBAL-H in toluene (40 ml, 40 mmol) was added dropwise at -70°C over 20 min. to a stirred solution of 3-biphenyl-4-yl-but-2-enoic acid ethyl ester (2.66 g, 10.0 mmol) in dry THF (100 ml) and the mixture stirred for 30 min. Methanol (2 ml) was added, followed by saturated aqueous Rochelle's salt (100 ml), and the resulting mixture extracted with ethyl acetate (200 ml), separated and the organic phase washed with brine, dried (Na_2SO_4), evaporated and dried *in vacuo* yielding (*E*)-3-biphenyl-4-yl-but-2-en-1-ol as colorless crystals: 1.94 g (86%)

^1H NMR (300 MHz, CDCl_3) δ : 1.40 (1H, br s), 2.12 (3H, d), 4.45 (2H, dd), 6.05 (1H, dt), 7.35-7.7 (9H, m). MS: 225 (M^+), 224(100%), 209, 181, 165.


Microanalysis Calculated % C: 86.00, H: 7.00. Found C: 85.67, H: 7.29.

c)

Diethyl azodicarboxylate (0.346 ml, 2.2 mmol) was added at 0°C to a stirred solution of triphenyl-phosphine (0.656 g, 2.2 mmol) and (*E*)-3-biphenyl-4-yl-but-2-en-1-ol (0.270 g, 1.2 mmol) in dry THF (20 ml) and the mixture stirred for 5 min. A solution of (*S*)-ethyl 2-ethoxy-3-(4-hydroxy-phenyl)-propionate (0.238 g, 1.0 mmol) in dry THF (10 ml) was added, the mixture allowed to warm to room temperature, and stirring continued for 48 h. The resulting mixture was evaporated *in vacuo* and the residue purified by column chromatography on silica gel (20% ethyl acetate in n-heptane) to give (*E*)-(*S*)-ethyl 3-[4-(3-biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-ethoxy-propionate as an oil; 0.288 g (65%).

^1H NMR (300 MHz, CDCl_3) δ : 1.13-1.25 (6H, m), 2.13 (3H, d), 2.94 (2H, d), 3.29-3.37 (1H, m), 3.54-3.61 (1H, m), 3.97 (1H, t), 4.1 (2H, q), 4.70 (2H, d), 6.11 (1H, dt), 6.86 (2H, d), 7.16 (2H, d), 7.25-7.63 (9H, m).

EXAMPLE 2




(*E*)-(*S*)-3-[4-(3-Biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-ethoxy-propionic acid

Sodium hydroxide (1M, 0.45 ml, 0.45 mmol) was added to a solution of (*E*)-(*S*)-ethyl 3-[4-(3-biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-ethoxy-propionate (example 1) (0.100 g, 0.225 mmol) in ethanol (20 ml) and the mixture stirred at 70°C for 2.5 h. After cooling to room temperature the resulting mixture was partitioned between water (50 ml) and ethyl acetate and the aqueous phase collected. The aqueous phase was acidified with 1N hydrochloric acid (5 ml) and extracted with ethyl acetate (100 ml), and the organic phase collected, washed with brine, dried (Na₂SO₄) and evaporated to give (*E*)-(*S*)-3-[4-(3-biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-ethoxy-propionic acid as a white solid; 0.014 g (15%).

¹H NMR (300 MHz, CDCl₃) δ: 1.19 (3H, t), 2.63 (3H, d), 2.93 (1H, dd), 3.1 (1H, dd), 3.4-3.65 (2H, m), 4.1 (2H, q), 4.72 (2H, d), 6.1 (1H, dt), 6.9 (2H, d), 7.2 (2H, d), 7.35-7.60 (9H, m).

EXAMPLE 3



(*E*)-(*S*)-3-{4-[3-(4'-Bromo-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester.

a)

(*E*)-3-(4'-Bromo-biphenyl-4-yl)-but-2-enoic acid ethyl ester was prepared from 4-(4-bromophenyl)acetophenone (12.0 g, 0.044 mol), sodium (1.25 g, 0.052 mol) and triethyl phosphonoacetate (11.73 g, 0.052 mol) by a procedure analogous to that described in example 1a yielding 11.97 g (80%).

¹H NMR (300 MHz, CDCl₃) δ: 1.32 (3H, t), 2.61 (3H, d), 4.23 (2H, q), 6.19 (1H, d), 7.40-7.58 (8H, m).

b)

(*E*)-3-(4'-bromo-biphenyl-4-yl)-but-2-en-1-ol was prepared from (*E*)-3-(4'-bromo-biphenyl-4-yl)-but-2-enoic acid ethyl ester (3.45 g, 10.0 mmol) and DIBAL-H (1M in toluene, 40 ml, 40 mmol) by a procedure analogous to that described in example 1b, yielding 1.68 g (55%).
¹H NMR (300MHz, CDCl₃) δ: 2.14 (3H, d), 4.4 (2H, t), 6.05 (1H, dt), 7.45-7.55 (8H, m),


5

c)

The title compound was prepared from (*E*)-3-(4'-bromo-biphenyl-4-yl)-but-2-en-1-ol (0.364 g, 1.2 mmol), triphenylphosphine (0.328 g, 1.3 mmol), diethyl azodicarboxylate (0.173 ml, 1.1 mmol) and (*S*)-ethyl 2-ethoxy-3-(4-hydroxy-phenyl)-propionate (0.238 g, 1.0 mmol) by a procedure analogous to that described in example 1c, yielding 0.180 g (34%) of (*E*)-(S)-3-{4-[3-(4'-bromo-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester.
¹H NMR (300 MHz, CDCl₃) δ: 1.15-1.25 (6H, m), 2.15 (3H, d), 2.95 (2H, d), 3.29-3.4 (1H, m), 3.5-3.65 (1H, m), 3.96 (1H, t), 4.15 (2H, q), 4.75 (2H, dd), 6.11 (1H, dt), 6.85 (2H, d), 7.14 (2H, d), 7.4-7.55 (8H, m).

15

EXAMPLE 4




20

(*E*)-(S)-3-{4-[3-(4'-Bromo-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid

The title compound was prepared from (*E*)-(S)-3-{4-[3-(4'-bromo-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester (example 3) (0.150 g, 0.29 mmol) and sodium hydroxide (1M, 0.45 ml, 0.45 mmol) by a procedure analogous to that described in example 2 yielding 0.180 g (34%) of (*E*)-(S)-3-{4-[3-(4'-bromo-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester.

¹H NMR (300 MHz, CDCl₃) δ: 1.14 (3H, t), 2.13 (3H, d), 2.86-3.10 (2H, m), 3.37-3.45 (1H, m), 3.55-3.65 (1H, m), 4.05 (2H, q), 4.70 (2H, dd), 6.12 (1H, dt), 6.9 (2H, d), 7.18 (2H, d), 7.4-7.60 (8H, m).

30

EXAMPLE 5

5


(E)-(S)-2-Ethoxy-3-{4-[3-(4-phenoxy-phenyl)-but-2-enyloxy]-phenyl}-propionic acid ethyl ester

The title compound was prepared from 4-phenoxyacetophenone (12.0 g, 0.056 mol) by a sequence analogous to that described in example 3, yielding 0.190 g (41%) of (E)-(S)-2-ethoxy-3-{4-[3-(4-phenoxy-phenyl)-but-2-enyloxy]-phenyl}-propionic acid ethyl ester.

10

¹H NMR (300 MHz, CDCl₃) δ: 1.2 (6H, m), 2.12 (3H, s), 2.97 (2H, d), 3.30-3.42 (1H, m), 3.59-3.70 (1H, m), 3.98 (1H, t), 4.15 (2H, q), 4.73 (2H, dd), 6.05 (1H, dt), 6.85-7.45 (13H, m).

15

EXAMPLE 6


(E)-(S)-2-Ethoxy-3-{4-[3-(4-phenoxy-phenyl)-but-2-enyloxy]-phenyl}-propionic acid

20

The title compound was prepared from (E)-(S)-2-ethoxy-3-{4-[3-(4-phenoxy-phenyl)-but-2-enyloxy]-phenyl}-propionic acid ethyl ester (example 5) (0.170 g, 0.37 mmol) and sodium hydroxide (1M, 0.74 ml, 0.74 mmol) by a procedure analogous to that described in example 2 yielding 0.136 g (85%) of (E)-(S)-2-ethoxy-3-{4-[3-(4-phenoxy-phenyl)-but-2-enyloxy]-phenyl}-propionic acid.

25

¹H NMR (300 MHz, CDCl₃) δ: 1.14 (3H, t), 2.13 (3H, d), 2.86-3.10 (2H, m), 3.38-3.45 (1H, m), 3.55-3.65 (1H, m), 4.05 (2H, q), 4.70 (2H, dd), 6.12 (1H, dt), 6.9 (2H, d), 7.18 (2H, d), 7.4-7.60 (8H, m).


EXAMPLE 7

(E)-(S)-2-Ethoxy-3-(4-{3-[4-(4-methoxy-phenoxy)-phenyl]-but-2-enyloxy}-phenyl)-propionic acid ethyl ester

The title compound was prepared from 4-(4-methoxyphenoxy)acetophenone (2.63 g, 0.011 mol) by a sequence analogous to that described in example 3 yielding 0.200 g (41%) of (E)-(S)-2-ethoxy-3-(4-{3-[4-(4-methoxy-phenoxy)-phenyl]-but-2-enyloxy}-phenyl)-propionic acid ethyl ester.

¹H NMR (300 MHz, CDCl₃) δ: 1.15-1.23 (6H, m), 2.12 (3H, s), 2.97 (2H, d), 3.30-3.40 (1H, m), 3.57-3.65 (1H, m), 3.80 (3H, s), 3.98 (1H, t), 4.18 (2H, q), 4.63 (2H, dd), 5.97-6.05 (1H, m), 6.85-6.96 (8H, m), 7.15 (2H, d), 7.35 (2H, d).

MS 490 (M⁺), 417, 359 (100%), 269.

EXAMPLE 8


(E)-(S)-2-Ethoxy-3-(4-{3-[4-(4-methoxy-phenoxy)-phenyl]-but-2-enyloxy}-phenyl)-propionic acid

The title compound was prepared from (*E*)-(*S*)-2-ethoxy-3-(4-{3-[4-(4-methoxy-phenoxy)-phenyl]-but-2-enyloxy}-phenyl)-propionic acid ethyl ester (example 7) (0.176 g, 0.36 mmol) and sodium hydroxide (1M, 0.74 ml, 0.74 mmol) by a procedure analogous to that described in example 2 yielding 0.140 g (84%) of (*E*)-(*S*)-2-ethoxy-3-(4-{3-[4-(4-methoxy-phenoxy)-phenyl]-but-2-enyloxy}-phenyl)-propionic acid.

¹H NMR (300 MHz, CDCl₃) δ: 1.15 (3H, t), 2.1 (3H, s), 2.9-3.1 (2H, m), 3.36-3.43 (1H, m), 3.55-3.64 (1H, m), 3.78 (3H, s), 4.00 (1H, dd), 4.70 (2H, dd), 6.0 (1H, dt), 6.8-6.9 (8H, m), 7.19 (2H, d), 7.35 (2H, d).

MS 462 (M⁺)(100%), 436, 359, 252.

EXAMPLE 9




(*E*)-(*S*)-2-Ethoxy-3-{4-[3-(9H-fluoren-2-yl)-but-2-enyloxy]-phenyl}-propionic acid ethyl ester

The title compound was prepared from 2-acetylfluorene (12.0 g, 0.058 mmol) by a sequence analogous to that described in example 3 yielding 0.200 g (41%) of (*E*)-(*S*)-2-ethoxy-3-{4-[3-(9H-fluoren-2-yl)-but-2-enyloxy]-phenyl}-propionic acid ethyl ester.

¹H NMR (300 MHz, CDCl₃) δ: 1.16-1.22 (6H, m), 2.2 (3H, s), 2.96 (2H, d), 3.30-3.40 (1H, m), 3.51-3.65 (1H, m), 3.9 (2H, s), 3.98 (1H, t), 4.15 (2H, q), 4.75 (2H, d), 6.04-6.13 (1H, m), 6.88 (2H, d), 7.17 (2H, d), 7.3-7.8 (7H, m).

MS 456 (M⁺), 410, 325 (100%), 238.

Microanalysis Calculated % C: 78.92, H: 7.06. Found C: 78.72, H: 7.30.

EXAMPLE 10


5

(E)-(S)-2-Ethoxy-3-{4-[3-(9H-fluoren-2-yl)-but-2-enyloxy]-phenyl}-propionic acid

The title compound was prepared from (E)-(S)-2-ethoxy-3-{4-[3-(9H-fluoren-2-yl)-but-2-enyloxy]-phenyl}-propionic acid ethyl ester (example 9) (0.230 g, 0.504 mol) and sodium hydroxide (1M, 1.008 ml, 1.008 mmol) by a procedure analogous to that described in example 2 yielding 0.140 g (84%) of (E)-(S)-2-ethoxy-3-{4-[3-(9H-fluoren-2-yl)-but-2-enyloxy]-phenyl}-propionic acid.

¹H NMR (300 MHz, CDCl₃) δ: 1.20 (3H, t), 2.18 (3H, s), 2.9-3.15 (2H, m), 3.4-3.6 (2H, m), 3.87 (2H, s), 4.05 (1H, dd), 4.75 (2H, d), 6.11 (1H, dt), 6.88 (2H, d), 7.17 (2H, d), 7.3-7.8 (7H, m).

20

EXAMPLE 11


(*E*)-(*S*)-3-{4-[3-(3,4-Dimethoxy-phenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester

The title compound was prepared from 3,4-dimethoxyacetophenone (10.00 g, 0.055 mol) by a sequence analogous to that described in example 3 yielding 0.160 g (31%) of (*E*)-(*S*)-3-{4-[3-(3,4-dimethoxy-phenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester.

¹H NMR (300 MHz, CDCl₃) δ: 1.1-1.19 (6H, m), 2.17 (3H, s), 2.98 (2H, d), 3.37-3.45 (1H, m), 3.58-3.65 (1H, m), 3.9 (6H, ds), 4.02 (1H, t), 4.15 (2H, q), 4.7 (2H, d), 6.0 (1H, dt), 6.81-6.86 (3H, m), 7.0 (2H, d), 7.15 (2H, d).

MS 428 (M⁺), 382, 355, 297 (100%), 207.


EXAMPLE 12



(*E*)-(*S*)-3-{4-[3-(3,4-Dimethoxy-phenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid

The title compound was prepared from (*E*)-(*S*)-3-{4-[3-(3,4-dimethoxy-phenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester (example 11) (0.150 g, 0.350 mmol) and sodium hydroxide (1M, 1.05 ml, 1.05 mmol) by a procedure analogous to that described in example 2 yielding 0.120 g (86%) of (*E*)-(*S*)-3-{4-[3-(3,4-dimethoxy-phenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid.

¹H NMR (300 MHz, CDCl₃) δ: 1.15 (3H, t), 2.15 (3H, s), 2.9-3.15 (2H, m), 3.40-3.48 (1H, m), 3.56-3.63 (1H, m), 3.9 (6H, ds), 4.08 (1H, dd), 4.75 (2H, d), 6.01 (1H, dt), 6.80-6.91 (3H, m), 7.0 (2H, d), 7.15 (2H, d).


EXAMPLE 13

5 *(E)*-(*S*)-3-{4-[3-(3,5-Bis-trifluoromethyl-phenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester

- 10 The title compound was prepared from 3,5-bis(trifluoromethyl)acetophenone (5.12 g, 0.02 mol) by a sequence analogous to that described in example 3 yielding 0.370 g (73%) of *(E)*-(*S*)-3-{4-[3-(3,5-bis-trifluoromethyl-phenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester.

- ¹H NMR (300 MHz, CDCl₃) δ: 1.1-1.25 (6H, m), 2.20 (3H, s), 2.97 (2H, d), 3.3-3.4 (1H, m),
15 3.62-3.7 (1H, m), 4.0 (1H, t), 4.15 (2H, q), 4.75 (2H, d), 6.2 (1H, dt), 6.85 (2H, d), 7.2 (2H, d),
7.78 (1H, br s), 7.87 (2H, br s).

MS 504 (M⁺), 458, 431(100%), 373, 267, 192


EXAMPLE 14

- 25 *(E)*-(*S*)-3-{4-[3-(3,5-Bis-trifluoromethyl-phenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid
-

The title compound was prepared from (*E*)-(*S*)-3-{4-[3-(3,5-bis-trifluoromethyl-phenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester (example 13) (0.200 g, 0.396 mmol) and sodium hydroxide (1M, 0.792 ml, 0.792 mmol) by a procedure analogous to that described in example 2 yielding 0.150 g (79%) of (*E*)-(*S*)-3-{4-[3-(3,5-bis-trifluoromethyl-phenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid.

¹H NMR (300 MHz, CDCl₃) δ: 1.12 (3H, t), 2.18 (3H, s), 2.9 (1H, dd), 3.1 (1H, dd), 3.34-3.42 (1H, m), 3.5-3.65 (1H, m), 4.0 (1H, dd), 4.7 (2H, d), 6.11 (1H, dt), 6.83 (2H, d), 7.19 (2H, d), 7.72 (1H, br s), 7.83 (2H, br s).


EXAMPLE 15



(*E*)-3-[4-(3-Biphenyl-4-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid ethyl ester

The title compound was prepared from 3-biphenyl-4-yl-acrylic acid ethyl ester (2.5 g, 0.01 mol) by a sequence analogous to that described in example 3b-c, yielding 0.370 g (73%) of (*E*)-3-[4-(3-biphenyl-4-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid ethyl ester.


¹H NMR (200 MHz, CDCl₃) δ: 1.1-1.25 (6H, m), 2.97 (2H, d), 3.3-3.4 (1H, m), 3.52-3.7 (1H, m), 4.0 (1H, t), 4.15 (2H, q), 4.75 (2H, dd), 6.35-6.5 (1H, dt), 6.75 (1H, d), 6.87 (2H, d), 7.15 (2H, d), 7.4-7.65 (9H, m).

EXAMPLE 165 (E)-(S)-3-[4-(3-Biphenyl-4-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid

The title compound was prepared from (E)-(S)-3-[4-(3-biphenyl-4-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid ethyl ester (example 15) (0.200 g, 0.464 mmol) and sodium hydroxide (1M, 0.928 ml, 0.928 mmol) by a procedure analogous to that described in example 2 yielding 0.043 g (23%) of (E)-(S)-3-[4-(3-biphenyl-4-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid.

¹H NMR (300 MHz, CDCl₃) δ: 1.15 (3H, t), 2.9 (1H, dd), 3.12 (1H, dd) 3.45-3.55 (2H, m), 3.84-3.96 (2H, m), 4.1 (1H, dd), 4.7 (2H, d), 6.35-6.5 (1H, dt), 6.78 (1H, d), 6.88 (2H, d), 7.15 (2H, d) 7.4-7.6 (9H, m).


15

EXAMPLE 1720 (E)-(S)-2-Ethoxy-3-[4-(3-naphthalen-2-yl-but-2-enyloxy)-phenyl]-propionic acid ethyl ester

The title compound was prepared from 2-acetonaphthone (10.0 g, 0.06 mol) by a sequence analogous to that described in example 3 yielding 0.190 g (38%) of (E)-(S)-2-ethoxy-3-[4-(3-naphthalen-2-yl-but-2-enyloxy)-phenyl]-propionic acid ethyl ester.

¹H NMR (200 MHz, CDCl₃) δ: 1.1-1.2 (6H, m), 2.20 (3H, s), 2.95 (2H, d), 3.3-3.4 (1H, m), 3.52-3.65 (1H, m), 3.95 (1H, t), 4.15 (2H, q), 4.76 (2H, d), 6.2 (1H, t), 6.85 (2H, d), 7.15 (2H, d), 7.35-7.42 (2H, m), 7.6 (1H, dd), 7.75-7.85 (4H, m).


5

EXAMPLE 18**(E)-(S)-2-Ethoxy-3-[4-(3-naphthalen-2-yl-but-2-enyloxy)-phenyl]-propionic acid**

The title compound was prepared from (*E*)-(*S*)-2-ethoxy-3-[4-(3-naphthalen-2-yl-but-2-enyloxy)-phenyl]-propionic acid ethyl ester (example 17) (0.165 g, 0.394 mmol) and sodium hydroxide (1M, 0.789 ml, 0.789 mmol) by a procedure analogous to that described in example 2 yielding 0.030 g (19%) of (*E*)-(*S*)-2-ethoxy-3-[4-(3-naphthalen-2-yl-but-2-enyloxy)-phenyl]-propionic acid.

¹H NMR (400 MHz, CDCl₃) δ: 1.13 (3H, t), 2.18 (3H, s), 2.95 (1H, dd), 3.05 (1H, dd), 3.3-3.45 (1H, m), 3.65-3.63 (1H, m), 3.95 (1H, dd), 4.72 (2H, d), 6.15 (1H, t), 6.84 (2H, d), 7.14 (2H, d), 7.35-7.45 (2H, m), 7.6 (1H, d), 7.7-7.8 (4H, m).

20

EXAMPLE 19

25


(E)-(S)-2-Ethoxy-3-[4-(3-pyridin-2-yl-but-2-enyloxy)-phenyl]-propionic acid ethyl ester

The title compound was prepared from 2-acetylpyridine (9.6 g, 0.08 mol) by a sequence analogous to that described in example 3 yielding 0.230 g (23%) of (*E*)-(*S*)-2-ethoxy-3-[4-(3-pyridin-2-yl-but-2-enyloxy)-phenyl]-propionic acid ethyl ester.

- 5 ¹H NMR (400 MHz, CDCl₃) δ: 1.1-2.5 (6H, m), 2.21 (3H, s), 2.97 (2H, d), 3.3-3.4 (1H, m), 3.58-3.64 (1H, m), 3.97 (1H, t), 4.15 (2H, q), 4.78 (2H, d), 6.65 (1H, t), 6.85 (2H, d), 7.05-7.15 (3H, m), 7.42 (2H, d), 7.6 (1H, dd), 8.52 (1H, d).

10

EXAMPLE 20




(*E*)-(*S*)-2-Ethoxy-3-[4-(3-pyridin-2-yl-but-2-enyloxy)-phenyl]-propionic acid

15

The title compound was prepared from (*E*)-(*S*)-2-ethoxy-3-[4-(3-pyridin-2-yl-but-2-enyloxy)-phenyl]-propionic acid ethyl ester (example 19) (0.220 g, 0.595 mmol) and sodium hydroxide (1M, 1.19 ml, 1.19 mmol) by a procedure analogous to that described in example 2 yielding 0.200 g (98%) of (*E*)-(*S*)-2-ethoxy-3-[4-(3-pyridin-2-yl-but-2-enyloxy)-phenyl]-propionic acid.

- 20 ¹H NMR (300 MHz, CDCl₃) δ: 1.2 (3H, t), 2.1 (3H, s), 2.7-2.85 (1H, m), 3.0-3.25 (2H, m), 3.5-3.6 (1H, m), 3.8-3.92 (1H, m), 4.6 (2H, d), 6.5 (1H, t), 6.75 (2H, d), 7.1-7.2 (3H, m), 7.35 (1H, d), 7.6 (1H, t), 8.5 (1H, d).


25

EXAMPLE 21

- 5 *(E)*-(*S*)-3-{4-[3-(3,5-Bis-benzyloxy-phenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester
-

The title compound was prepared from 3,5-dibenzyloxyacetophenone (6.64 g, 0.02 mol) by a sequence analogous to that described in example 3 yielding 0.460 g (53%) of *(E)*-(*S*)-3-{4-[3-(3,5-bis-benzyloxy-phenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester.

¹H NMR (300 MHz, CDCl₃) δ: 1.1-1.21 (6H, m), 2.14 (3H, s), 2.95 (2H, d), 3.28-3.41 (1H, m), 3.51-3.65 (1H, m), 3.94 (1H, t), 4.12 (2H, q), 4.7 (2H, d), 5.05 (4H, s), 6.05 (1H, t), 6.53-6.57 (1H, m), 6.67 (2H, d), 6.85 (2H, d), 7.12 (2H, d), 7.3-7.45 (10H, m).


EXAMPLE 22

(E)-(S)-3-{4-[3-(3,5-Bis-benzyloxy-phenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid

The title compound was prepared from (E)-(S)-3-{4-[3-(3,5-bis-benzyloxy-phenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester (example 21) (0.430 g, 0.741 mmol) and sodium hydroxide (1M, 1.5 ml, 1.5 mmol) by a procedure analogous to that described in example 2 yielding 0.300 g (73%) of (E)-(S)-3-{4-[3-(3,5-bis-benzyloxy-phenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid.

¹H NMR (300 MHz, CDCl₃) δ: 1.15 (3H, t), 2.1 (3H, s), 2.95 (1H, dd), 3.05 (1H, dd), 3.36-3.44 (1H, m), 3.57-3.65 (1H, m), 4.05 (1H, dd), 4.68 (2H, d), 5.05 (4H, s), 6.05 (1H, t), 6.52 (1H, m), 6.65 (2H, d), 6.85 (2H, d), 7.15 (2H, d), 7.3-7.45 (10H, m).

EXAMPLE 23



(E)-(S)-2-Ethoxy-3-[4-(3-naphthalen-2-yl-allyloxy)-phenyl]-propionic acid ethyl ester

a)

Triethyl phosphonoacetate (8.9 g, 40.0 mmol) was added at 0°C over a period of 10 min. to a stirred suspension of sodium hydride (60% in oil, 1.44 g, 36.0 mmol) in dry THF (145 mL). After stirring at 0°C for 15 min. a solution of 2-naphthaldehyde (3.12 g, 20.0 mmol) in dry THF (15 mL) was added, the mixture slowly warmed to room temperature, and stirring continued for 16h. The reaction mixture was quenched with water (100 mL) and acidified to pH 6 with 1N hydrochloric acid. Additional water (200 mL) was added, the organic phase separated, and the aqueous phase further extracted with ethyl acetate (300 mL). The combined organic phases were washed with water (200 mL x 3), dried (MgSO₄), filtered and concentrated *in vacuo* to give 6.5 g of crude (E)-3-naphthalen-2-yl-acrylic acid ethyl ester.

b)


Crude (*E*)-3-naphthalen-2-yl-acrylic acid ethyl ester (4.5 g, 20.0 mmol) was reduced by a procedure analogous to that described in example 1b. The product was purified by flash column chromatography to give 3.1 g (86%) of (*E*)-3-naphthalen-2-yl-prop-2-en-1-ol.

c)

Under an atmosphere of nitrogen, (*E*)-3-naphthalen-2-yl-prop-2-en-1-ol (190 mg, 0.8 mmol), tributylphosphine (323 mg, 1.6 mmol) and (*S*)-2-ethoxy-3-(4-hydroxy-phenyl)-propionic acid ethyl ester (184 mg, 1.0 mmol) were successively dissolved in dry benzene (20 mL). Solid 1,1'-(azodicarbonyl) dipiperidine (403 mg, 1.6 mmol) was added at 0°C with stirring. After 10 min. the reaction was warmed to room temperature and the stirring continued for 1 h. The reaction mixture was concentrated *in vacuo* and the product purified by flash column chromatography, eluting with heptane/ethyl acetate (3:2), to give 180 mg (55%) of the title compound.

¹H NMR (CDCl₃, 300 MHz) δ: 1.15 (t, 3H), 1.22 (t, 3H), 2.95 (d, 2H), 3.28-3.40 (m, 1H), 3.55-3.65 (m, 1H), 3.96 (t, 1H), 4.15 (q, 2H), 4.72 (dd, 2H), 6.53 (dt, 1H), 6.83-6.93 (m, 3H), 7.18 (d, 2H), 7.40-7.50 (m, 2H), 7.13 (dd, 1H), 7.72-7.85 (m, 4H).

EXAMPLE 24



(*E*)-(S)-2-Ethoxy-3-[4-(3-naphthalen-2-yl-allyloxy)-phenyl]-propionic acid


(*E*)-(S)-2-Ethoxy-3-[4-(3-naphthalen-2-yl-allyloxy)-phenyl]-propionic acid ethyl ester (example 23) (170 mg, 0.42 mmol) was dissolved in ethanol (20 mL) at 35°C and sodium hydroxide (1N, 2.1 mL, 2.1 mmol) added. The mixture was stirred at 35°C for 1 h, the ethanol evapo-

rated *in vacuo* and the mixture acidified to pH 1 with 1N hydrochloric acid. The product was isolated by extraction with ethyl acetate (30 mL x 2). The combined organic phases were dried (MgSO₄), filtered and evaporated to give 155 mg (98%) of the title compound as crystals.

- 5 ¹H NMR (CDCl₃, 300 MHz) δ: 1.18 (t, 3H), 2.90-3.12 (m, 2H), 3.35-3.48 (m, 1H), 3.55-3.68 (m, 1H), 4.03 (q, 1H), 4.70 (dd, 2H), 6.52 (dt, 1H), 6.80-6.95 (m, 3H), 7.18 (d, 2H), 7.40-7.48 (m, 2H), 7.60 (dd, 1H), 7.70-7.80 (m, 4H).

10

EXAMPLE 25




(E)-(S)-2-Ethoxy-3-{4-[3-(3-phenoxy-phenyl)-allyloxy]-phenyl}-propionic acid ethyl ester

15

The title compound was prepared from 3-phenoxybenzaldehyde (4.0 g, 20.0 mmol) by a sequence analogous to that described in example 23.

- 16 ¹H NMR (CDCl₃, 300 MHz) δ: 1.15 (t, 3H), 1.22 (t, 3H), 2.95 (d, 2H), 3.30-3.40 (m, 1H), 3.55-3.68 (m, 1H), 3.98 (t, 1H), 4.15 (q, 2H), 4.72 (dd, 2H), 6.38 (dt, 1H), 6.67 (d, 1H), 6.83-6.93 (m, 3H), 6.97-7.20 (m, 7H), 7.22-7.38 (m, 3H).
- 20


EXAMPLE 265 (E)-(S)-2-Ethoxy-3-{4-[3-(3-phenoxy-phenyl)-allyloxy]-phenyl}-propionic acid

(E)-(S)-2-Ethoxy-3-{4-[3-(3-phenoxy-phenyl)-allyloxy]-phenyl}-propionic acid ethyl ester (example 25) (150 mg, 0.34 mmol) was dissolved in ethanol (7 mL) and sodium hydroxide (1N, 4.4 mL, 4.4 mmol) added. The mixture was heated slightly to obtain a clear solution and then
10 stirred at room temperature for 1.5 h. The ethanol was evaporated *in vacuo* and the mixture acidified to pH 1 with 1N hydrochloric acid. The product was isolated by extraction with ethyl acetate (40 mL x 2). The combined organic phases were dried (MgSO₄), filtered and evaporated to give 130 mg (91%) of the title compound as an oil.

¹H NMR (CDCl₃, 300 MHz) δ : 1.18 (t, 3H), 2.95 (dd, 1H), 3.08 (dd, 1H), 3.38-3.50 (m, 1H),
15 3.55-3.65 (m, 1H), 4.05 (q, 1H), 4.65 (dd, 1H), 6.35 (dt, 1H), 6.66 (d, 1H), 6.85-6.92 (m, 3H), 6.98-7.20 (m, 7H), 7.25-7.40 (m, 3H).

EXAMPLE 27


20

(S)-3-[4-(2-Benzofuran-3-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid ethyl ester

The title compound was prepared from benzo[b]furan-2-carboxaldehyde (9.8 g, 0.07 mol) by a sequence analogous to that described in example 23.

¹H NMR (CDCl₃, 300 MHz) δ: 1.15 (t, 3H), 1.22 (t, 3H), 2.95 (d, 2H), 3.30-3.42 (m, 1H), 3.55-3.65 (m, 1H), 3.98 (t, 1H), 4.15 (q, 2H), 4.73 (d, 2H), 6.65-6.70 (m, 3H), 6.88 (d, 2H), 7.15 (d, 2H), 7.20-7.30 (m, 2H), 7.45 (d, 1H), 7.53 (d, 1H).


EXAMPLE 28



(S)-3-[4-(2-Benzofuran-3-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid

The title compound was prepared from (S)-3-[4-(2-benzofuran-3-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid ethyl ester (example 27) (127 mg, 0.3 mmol) by a procedure analogous to that described in example 26.

¹H NMR (CDCl₃, 300 MHz) δ: 1.15 (t, 3H), 3.30 (dd, 1H), 3.08 (dd, 1H), 2.38-3.50 (m, 1H), 3.55-3.65 (m, 1H), 4.05 (q, 1H), 4.72 (d, 2H), 6.55-6.68 (m, 3H), 6.90 (d, 1H), 7.13-7.30 (m, 5H), 7.42 (d, 1H), 7.50 (d, 1H).

EXAMPLE 29

(*E*)-(*S*)-3-{4-[3-(4-Benzyloxy-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester

5


The title compound was prepared from 4-benzyloxybenzaldehyde (21.2 g, 0.1 mol) by a sequence analogous to that described in example 23. The title compound was purified on HPLC, using ethyl acetate/heptane (20:80) as eluent.

¹H NMR (CDCl₃, 300 MHz) δ: 1.15 (t, 3H), 1.22 (t, 3H), 2.95 (d, 2H), 3.35 (m, 1H), 3.6 (m, 1H), 3.98 (t, 1H), 4.15 (q, 2H), 4.65 (dd, 2H), 5.05 (s, 2H), 2.28 (dt, 1H), 6.65 (d, 1H), 6.85 (d, 2H), 6.93 (d, 2H), 7.15 (d, 2H), 7.30-7.48 (m, 7H).

10

EXAMPLE 30

15




(*E*)-(*S*)-3-{4-[3-(4-Benzyloxy-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid

The title compound was prepared from (*E*)-(*S*)-3-[4-[3-(4-Benzyloxy-phenyl)-allyloxy]-phenyl]-2-ethoxy-propionic acid ethyl ester (example 29) (80 mg, 0.17 mmol) by a procedure analogous to that described in example 26.

- 5 ¹H NMR (CDCl₃, 300 MHz): 1.18 (t, 3H), 2.95 (dd, 1H), 3.12 (dd, 1H), 3.45-3.60 (m, 2H), 4.15 (dd, 1H), 4.65 (dd, 2H), 5.06 (s, 2H), 6.25 (dt, 1H), 6.65 (d, 1H), 6.90 (d, 2H), 6.93 (d, 2H), 7.15 (d, 2H), 7.30-7.45 (m, 7H).

10

EXAMPLE 31




(*E*)-(*S*)-3-[4-(3-Benzo[1,3]dioxol-5-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid ethyl ester

15

The title compound was prepared from piperonal (3.0 g, 20 mmol) by a sequence analogous to that described in example 23. The title compound was purified on HPLC, using ethyl acetate/heptane (10:90) as eluent.


- 20 ¹H NMR (CDCl₃, 300 MHz) δ: 1.15 (t, 3H), 1.22 (t, 3H), 2.96 (d, 2H), 3.30-3.42 (m, 1H), 3.55-3.65 (m, 1H), 3.97 (t, 1H), 4.15 (q, 2H), 4.63 (dd, 2H), 5.96 (s, 2H), 6.25 (dt, 1H), 6.63 (d, 1H), 6.75 (d, 1H), 6.80-6.90 (m, 3H), 6.95 (d, 1H), 7.15 (d, 2H).

EXAMPLE 325 (E)-(S)-3-[4-(3-Benzo[1,3]dioxol-5-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid

The title compound was prepared from (E)-(S)-3-[4-(3-benzo[1,3]dioxol-5-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid ethyl ester (example 31) (100 mg, 0.25 mmol) by a procedure analogous to that described in example 26.

10 ¹H NMR (CDCl₃, 300 MHz): 1.18 (t, 3H), 2.95 (dd, 1H), 3.08 (dd, 1H), 3.38-3.50 (m, 1H), 3.55- 3.68 (m, 1H), 4.05 (dd, 1H), 4.65 (dd, 2H), 5.95 (s, 2H), 6.25 (dt, 1H), 6.63 (d, 1H), 6.75 (d, 1H), 6.83 (dd, 1H), 6.88 (d, 2H), 6.95 (d, 1H), 7.17 (d, 2H).

15

EXAMPLE 33


(E)-(S)-3-[4-[3-(4-Allyloxy-phenyl)-allyloxy]-phenyl]-2-ethoxy-propionic acid ethyl ester

20

The title compound was prepared from 4-allyloxybenzaldehyde (3.24 g, 20 mmol) by a sequence analogous to that described in example 23. The title compound was purified on HPLC, using ethyl acetate/heptane (10:90) as eluent.

¹H NMR (CDCl₃, 300 MHz) δ: 1.18 (t, 3H), 1.22 (t, 3H), 2.95 (d, 2H), 3.30-3.42 (m, 1H), 3.55-3.68 (m, 1H), 3.98 (t, 1H), 4.17 (q, 2H), 4.53 (d, 2H), 4.65 (dd, 2H), 5.29 (dd, 1H), 5.40 (dd, 1H), 5.97-6.13 (m, 1H), 6.28 (dt, 1H), 6.65 (d, 1H), 6.88 (d, 4H), 7.15 (d, 2H), 7.35 (d, 2H).


EXAMPLE 34



(E)-(S)-3-{4-[3-(4-Allyloxy-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid

15 The title compound was prepared from (E)-(S)-3-{4-[3-(4-allyloxy-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester (example 33) (40 mg, 0.1 mmol) by a procedure analogous to that described in example 26.

¹H NMR (CDCl₃, 300 MHz): 1.18 (t, 3H), 2.95 (dd, 1H), 3.10 (dd, 1H), 3.39-3.50 (m, 1H), 3.53-3.65 (m, 1H), 4.05 (dd, 1H), 4.53 (d, 2H), 4.65 (d, 2H), 5.29 (dd, 1H), 5.40 (dd, 1H),
20 5.98-6.14 (m, 1H), 6.28 (dt, 1H), 6.65 (d, 1H), 6.85-6.95 (m, 4H), 7.15 (d, 2H), 7.35 (d, 2H).


EXAMPLE 355 (E)-(S)-3-[4-(3-Benzofuran-7-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid ethyl ester

The title compound was prepared from benzofuran-7-carboxaldehyde (1.46 g, 10 mmol) by a sequence analogous to that described in example 23.

¹H NMR (CDCl₃, 300 MHz) δ: 1.15 (t, 3H), 1.22 (t, 3H), 2.95 (d, 2H), 3.30-3.42 (m, 1H), 3.55-
10 3.65 (m, 1H), 3.98 (t, 1H), 4.15 (q, 2H), 4.75 (dd, 2H), 6.79 (d, 1H), 6.87-7.00 (m, 4H), 7.13-
7.30 (m, 4H), 7.50 (dd, 1H), 7.65 (d, 1H).

EXAMPLE 36


15

(E)-(S)-3-[4-(3-Benzofuran-7-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid

20 The title compound was prepared from (E)-(S)-3-[4-(3-benzofuran-7-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid ethyl ester (example 35) (100 mg, 0.25 mmol) by a procedure analogous to that described in example 26.

^1H NMR (CDCl_3 , 300 MHz) δ : 1.15 (t, 3H), 2.95 (dd, 1H), 3.08 (dd, 1H), 3.35-3.48 (m, 1H) 3.55-3.68 (m, 1H), 4.03 (dd, 1H), 4.75 (dd, 2H), 6.78 (d, 1H), 6.90-7.00 (m, 4H), 7.13-7.32 (m, 4H), 7.50 (dd, 1H), 7.65 (d, 1H), 10.1 (bs, 1H).

5

EXAMPLE 37


10 (S)-3-[4-(3-Benzodioxol-4-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid ethyl ester

The title compound was prepared from 2,3-methylenedioxybenzaldehyde (1.5 g, 10 mmol) by a sequence analogous to that described in example 23.

15 ^1H NMR (CDCl_3 , 300 MHz) δ : 1.18 (t, 3H), 1.22 (t, 3H), 2.95 (d, 2H), 3.30-3.42 (m, 1H), 3.55-3.65 (m, 1H), 3.97 (t, 1H), 4.15 (q, 2H), 4.65 (d, 2H), 6.00 (s, 2H), 6.55-6.92 (m, 7H), 7.15 (d, 2H).

EXAMPLE 38

20




(S)-3-[4-(3-Benzodioxol-4-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid

The title compound was prepared from (S)-3-[4-(3-benzo[1,3]dioxol-4-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid ethyl ester (example 37) (100 mg, 0.24 mmol) by a procedure analogous to that described in example 26.

- 5 ¹H NMR (CDCl₃, 300 MHz) δ: 1.17 (t, 3H), 2.95 (dd, 1H), 3.05 (dd, 1H), 3.35-3.48 (m, 1H), 3.55-3.68 (m, 1H), 4.03 (dd, 1H), 4.65 (d, 2H), 6.00 (s, 2H), 6.55-6.95 (m, 7H), 7.19 (d, 2H).


EXAMPLE 39

10



(E)-(S)-2-Ethoxy-3-(4-[3-(9H-fluoren-2-yl)-allyloxy]-phenyl)-propionic acid ethyl ester


- 15 The title compound was prepared from fluorene-2-carboxaldehyde (9.7 g, 50 mmol) by a sequence analogous to that described in example 23.
- ¹H NMR (CDCl₃, 300 MHz) δ: 1.18 (t, 3H), 1.22 (t, 3H), 2.97 (d, 2H), 3.32-3.42 (m, 1H), 3.55-3.67 (m, 1H), 3.90 (s, 2H), 3.98 (t, 1H), 4.16 (q, 2H), 4.70 (dd, 2H), 6.45 (dt, 1H), 6.80 (d, 1H), 6.90 (d, 2H), 7.1 (d, 2H), 7.24-7.46 (m, 3H), 7.55 (d, 1H), 7.62 (s, 1H), 7.72-7.80 (m, 2H).
- 20

EXAMPLE 405 (E)-(S)-2-Ethoxy-3-(4-[3-(9H-fluoren-2-yl)-allyloxy]-phenyl)-propionic acid

The title compound was prepared from (E)-(S)-2-ethoxy-3-(4-[3-(9H-fluoren-2-yl)-allyloxy]-phenyl)-propionic acid ethyl ester (example 39) (275 mg, 0.6 mmol) by a procedure analogous to that described in example 26.

10 ¹H NMR (CDCl₃, 300 MHz) δ: 1.20 (t, 3H), 3.46 (dd, 1H), 3.12 (dd, 1H), 3.43-3.65 (m, 2H), 3.90 (s, 2H), 4.05 (dd, 1H), 4.70 (dd, 2H), 6.46 (dt, 1H), 6.80 (d, 1H), 6.92 (d, 2H), 7.17 (d, 2H), 7.23-7.46 (m, 3H), 7.53 (d, 1H), 7.60 (s, 1H), 7.70-7.80 (m, 2H).

15


EXAMPLE 41(S)-2-Ethoxy-3-[4-(3-quinolin-2-yl-allyloxy)-phenyl]-propionic acid ethyl ester

20

The title compound was prepared from 2-quinoline-carboxaldehyde (5.12 g, 32.5 mmol) by a sequence analogous to that described in example 23.

¹H NMR (CDCl₃, 300 MHz) δ: 1.18 (t, 3H), 1.22 (t, 3H), 2.98 (d, 2H), 3.32-3.42 (m, 1H), 3.55-3.66 (m, 1H), 3.98 (t, 1H), 4.17 (q, 2H), 4.80 (d, 2H), 6.92 (d, 2H), 7.02 (m, 2H), 7.18 (d, 2H), 7.47-7.60 (m, 2H), 7.70 (dt, 1H), 7.78 (d, 1H), 8.05 (d, 1H), 8.13 (d, 1H).

5


EXAMPLE 42

10 (S)-2-Ethoxy-3-[4-(3-quinolin-2-yl-allyloxy)-phenyl]-propionic acid

(S)-2-Ethoxy-3-[4-(3-quinolin-2-yl-allyloxy)-phenyl]-propionic acid ethyl ester (example 41) (150 mg, 0.37 mmol) was dissolved in ethanol (2 mL) and sodium hydroxide (1N, 2.0 mL, 2.0 mmol) added. The mixture was stirred at room temperature for 16 h. The reaction mixture
15 was concentrated in vacuo, added 2-propanol (2 mL) and diethyl ether (2 mL). The title compound was isolated by filtration.

¹H NMR (CDCl₃/MeOD, 300 MHz) δ: 1.12 (t, 3H), 2.83 (dd, 1H), 3.02 (dd, 1H), 3.32 (m, 1H), 3.56 (dd, 1H), 3.84 (dd, 1H), 4.85 (d, 2H), 6.90-7.10 (m, 4H), 7.25 (m, 2H), 7.5-7.6 (m, 1H), 7.68-7.75 (m, 2H), 7.85 (d, 1H), 8.03 (d, 1H), 8.23 (d, 1H).


20

EXAMPLE 43

- 5 *(E)-(S)*-3-{4-[3-(3,5-Bis-benzyloxy-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester
-

The title compound was prepared from 3,5-dibenzyloxybenzaldehyde (3.1 g, 9.7 mmol) by a sequence analogous to that described in example 23.


- 10 ^1H NMR (CDCl_3 , 300 MHz) δ : 1.16 (t, 3H), 1.22 (t, 3H), 2.96 (d, 2H), 3.30-3.42 (m, 1H), 3.54-3.65 (m, 1H), 3.98 (t, 1H), 4.17 (q, 2H), 4.65 (d, 2H), 5.02 (s, 4H), 6.38 (dt, 1H), 6.55 (s, 1H), 6.58-6.70 (m, 3H), 6.88 (d, 2H), 7.15 (d, 2H), 7.30-7.45 (m, 10H).

EXAMPLE 445 (E)-(S)-3-{4-[3-(3,5-Bis-benzyloxy-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid

The title compound was prepared from (E)-(S)-3-{4-[3-(3,5-bis-benzyloxy-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester (example 43) (587 mg, 1.1 mmol) by a procedure analogous to that described in example 26.

- 10 ^1H NMR (CDCl_3 , 300 MHz) δ : 1.15 (t, 3H), 2.95 (dd, 1H), 3.08 (dd, 1H), 3.38-3.48 (m, 1H), 3.54-3.65 (m 1H), 4.03 (dd, 1H), 4.65 (d, 2H), 5.03 (s, 4H), 6.35 (dt, 1H), 6.54 (t, 1H), 6.60-6.70 (m, 3H), 6.88 (d, 2H), 7.16 (d, 2H), 7.30-7.45 (m, 10 H).

15

EXAMPLE 45


(E)-(S)-3-{4-[3-(3,5-Dimethoxy-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester

The title compound was prepared from 3,5 dimethoxybenzaldehyde (5.5 g, 33.1 mmol) by a sequence analogous to that described in example 23.

- 5 ¹H NMR (CDCl₃, 300 MHz) δ: 1.18 (t, 3H), 1.22 (t, 3H), 2.95 (d, 2H), 3.30-3.40 (m, 1H), 3.53-3.65 (m, 1H), 3.78 (s, 6H), 3.97 (t, 1H), 4.15 (q, 2H), 4.65 (dd, 1H), 6.33-6.43 (m, 2H), 6.55 (d, 2H), 6.88 (d, 2H), 7.15 (d, 2H).

10

EXAMPLE 46




(E)-(S)-3-{4-[3-(3,5-Dimethoxy-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid

15

The title compound was prepared from (E)-(S)-3-{4-[3-(3,5-dimethoxy-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester (example 45) (300 mg, 0.7 mmol) by a procedure analogous to that described in example 26.


- 20 ¹H NMR (CDCl₃, 300 MHz) δ: 1.18 (t, 3H), 2.95 (dd, 1H), 3.07 (dd, 1H), 3.37-3.48 (m, 1H), 3.55-3.67 (m, 1H), 3.80 (s, 6H), 4.05 (dd, 1H), 4.67 (d, 2H), 6.33-6.45 (m, 1H), 6.55 (d, 2H), 6.65 (d, 1H), 6.88 (d, 2H), 7.18 (d, 2H).

EXAMPLE 475 (E)-(S)-2-Ethoxy-3-[4-(3-phenanthren-9-yl-allyloxy)-phenyl]-propionic acid ethyl ester

The title compound was prepared from phenanthrene-9-carboxaldehyde (4.1 g, 20.0 mmol) by a sequence analogous to that described in example 23.

¹H NMR (CDCl₃, 300 MHz) δ: 1.16 (t, 3H), 1.22 (t, 3H), 2.95 (d, 2H), 3.30-3.42 (m, 1H), 3.53-
10 3.65 (m, 1H), 3.98 (t, 1H), 4.15 (q, 2H), 4.47 (d, 2H), 6.47 (dt, 1H), 6.74 (d, 2H), 7.08 (d, 2H),
7.38 (d, 1H), 7.53-7.70 (m, 4H), 7.82 (s, 1H), 7.85 (d, 1H), 8.15 (d, 1H), 8.65 (d, 1H), 8.72 (d,
1H).


15

EXAMPLE 4820 (E)-(S)-2-Ethoxy-3-[4-[3-(2-methoxy-naphthalen-1-yl)-allyloxy]-phenyl]-propionic acid ethyl ester

The title compound was prepared from 2-methoxy-1-naphthaldehyde (4.1 g, 22.1 mmol) by a sequence analogous to that described in example 23.

¹H NMR (CDCl₃, 300 MHz) δ: 1.16 (t, 3H), 1.22 (t, 3H), 2.97 (d, 2H), 3.30-3.42 (m, 1H), 3.55-3.65 (m, 1H), 3.93 (s, 3H), 3.97 (t, 1H), 4.15 (q, 2H), 4.85 (d, 2H), 6.48 (dt, 1H), 6.95 (d, 2H),
5 7.10-7.35 (m, 5H), 7.45 (dt, 1H), 7.75-7.78 (m, 2H), 8.12 (d, 1H).

EXAMPLE 49




(E)-(S)-2-Ethoxy-3-{4-[3-(2-methoxy-naphthalen-1-yl)-allyloxy]-phenyl}-propionic acid

The title compound was prepared from (E)-(S)-2-ethoxy-3-{4-[3-(2-methoxy-naphthalen-1-yl)-allyloxy]-phenyl}-propionic acid ethyl ester (example 48) (327 mg, 0.75 mmol) by a procedure
15 analogous to that described in example 26.

¹H NMR (CDCl₃, 300 MHz) δ: 1.16 (t, 3H), 2.95 (dd, 1H), 3.08 (dd, 1H), 3.35-3.48 (m, 1H),
3.53-3.65 (m, 1H), 3.93 (s, 3H), 4.05 (dd, 1H), 4.82 (dd, 2H), 6.49 (dt, 1H), 6.95 (d, 2H), 7.13
20 (d, 1H), 7.20 (d, 2H), 7.23-7.35 (m, 2H), 7.44 (dt, 1H), 7.74 (d, 2H), 8.12 (d, 1H).

EXAMPLE 50



(*E*)-(*S*)-Ethyl 3-{4-[3-(4-Bromophenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate

a)

- 5 Sodium (5.52 g, 0.24 mol) was added to ethanol (250 ml) at 20°C and the mixture stirred until the metal had fully reacted. Triethyl phosphonoacetate (62.72 g, 0.28 mol) was added, the mixture stirred for 20 min, then a solution of 4-bromoacetophenone (39.81 g, 0.20 mol) in ethanol (250 ml) was added and the reaction mixture heated to 80°C under reflux for 17h. The solution was cooled, the ethanol evaporated and the resulting orange residue partitioned
- 10 between 1N HCl (200 ml) and ethyl acetate (200 ml). The aqueous layer was collected and further extracted with ethyl acetate (2 x 200 ml). The organic layers were combined, washed with brine, dried (MgSO₄) and evaporated to an orange gum. This was purified by column chromatography on silica gel (3% diethyl ether in n-heptane eluent) to give the product, (*E*)-ethyl 3-(4-bromophenyl)-but-2-enoate, as a colourless oil; 44.08 g (82%)
- 15 ¹H NMR (300 MHz, CDCl₃) δ: 1.31 (3H, t), 2.54 (3H, s), 4.21 (2H, q), 6.11 (1H, s), 7.34 (2H, dm), 7.48 (2H, dm). MS: 268/270 (M⁺), 240/242, 239/241, 196/198, 116, 115 (100%).
- Microanalysis Calculated % C: 53.55, H: 4.87. Found % C: 53.86, H: 4.90.

b)


- 20 A 1M solution of DIBAL-H in toluene (42 ml, 42 mmol) was added dropwise, at -70°C over 30 min, to a stirred solution of (*E*)-ethyl 3-(4-bromophenyl)-but-2-enoate (4.55 g, 16.92 mmol) in dry THF (100 ml), and the mixture stirred for 1 h. Methanol (5 ml) was carefully added, followed by 1N HCl (300 ml) and the resulting mixture extracted with ethyl acetate (3 x 200 ml). The combined organic extracts were washed with brine, dried (Na₂SO₄), and evaporated to
- 25 give the crude product as an off-white solid, which was purified by recrystallisation from hot 1:4 ether/n-heptane (250 ml) to give the product (*E*)-3-(4-bromophenyl)-but-2-en-1-ol as colourless needles: 3.10 g (81%)
- Mpt. 58-59.5°C. ¹H NMR (300 MHz, CDCl₃) δ: 1.41 (1H, br s), 2.05 (3H, d), 4.36 (2H, d), 5.96 (1H, tq), 7.27 (2H, dm), 7.44 (2H, dm). MS: 226/228 (M⁺), 211/213, 193/195, 183/185, 147
- 30 (100%), 132, 129, 115. Microanalysis Calculated % C: 52.89, H: 4.88, Br: 35.18. Found C: 53.24, H: 4.86, Br: 35.08.

c)

Azodicarboxylic dipiperidide (0.756 g, 3.0 mmol) was added at 0-5°C to a stirred solution of tributylphosphine (0.74 ml, 0.61 g, 3.0 mmol), (*S*)-ethyl 2-ethoxy-3-(4-hydroxyphenyl)-propionate (0.500 g, 2.10 mmol) and (*E*)-3-(4-bromophenyl)-but-2-en-1-ol (0.454 g, 2.0 mmol) in dry benzene (20 ml), the mixture warmed to room temperature, and stirred for 2.5 days. The resulting mixture was diluted with water and ethyl acetate (50 ml each), the aqueous layer collected and further extracted with ethyl acetate (50 ml). The organic layers were combined, washed with brine, dried (MgSO₄) and evaporated. The crude product was then purified by column chromatography on silica gel (20% ethyl acetate in n-heptane eluent) to give (*E*)-(*S*)-ethyl 3-{4-[3-(4-bromophenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate as an oil; 0.780 g (87%).

¹H NMR (300 MHz, CDCl₃) δ: 1.17 (3H, t), 1.22 (3H, t), 2.10 (3H, s), 2.96 (2H, d), 3.30-3.45 (1H, m), 3.55-3.70 (1H, m), 3.98 (1H, t), 4.17 (2H, q), 4.70 (2H, d), 6.04 (1H, t), 6.86 (2H, m), 7.16 (2H, m), 7.29 (2H, m), 7.44 (2H, m).

EXAMPLE 51




(*E*)-(*S*)-3-{4-[3-(4-Bromophenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid

Sodium hydroxide (1M, 1.10 ml, 1.10 mmol) was added to a solution of (*E*)-(*S*)-ethyl 3-{4-[3-(4-bromophenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate (example 50) (0.245 g, 0.548 mmol) in ethanol (10 ml) and the mixture stirred at room temperature for 18 h. The resulting mixture was partitioned between water (50 ml) and ethyl acetate (50 ml) and the aqueous layer acidified to pH1 by addition of 1N HCl. The aqueous layer was separated and further extracted with ethyl acetate (2 x 50 ml). The combined organic layers were washed with

brine, dried (MgSO_4), evaporated and vacuum dried at 40°C for 18 h, to give (*E*)-(*S*)-3-{4-[3-(4-bromophenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid as a colourless gum which contained 0.1 molar equivalents of ethyl acetate; 0.22 g (96%).

^1H NMR (300 MHz, CDCl_3) δ : 1.18 (3H, t), 1.26 (ethyl acetate impurity, 0.3H, t), 2.04 (ethyl acetate impurity, 0.2H, s), 2.11 (3H, s), 2.96 (1H, dd), 3.08 (1H, dd), 3.40-3.55 (1H, m), 3.55-3.68 (1H, m), 4.06 (1H, dd), 4.15 (ethyl acetate impurity, 0.2H, q), 4.70 (2H, d), 6.04 (1H, t), 6.88 (2H, m), 7.17 (2H, m), 7.29 (2H, m), 7.44 (2H, m), carboxylic acid proton not observed. LCMS: 441/443 ($\text{M}+\text{Na}$), 209/211 (100%).

10

EXAMPLE 52

15 **(*E*)-(*S*)-Ethyl 3-{4-[3-(4'-Chloro-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate**

a)

Tetrakis(triphenylphosphine)palladium(0) (0.26 g, 0.22 mmol, 4 mol%) was added, under nitrogen, to a stirred solution of (*E*)-ethyl 3-(4-bromophenyl)-but-2-enoate (1.5 g, 5.57 mmol) {prepared as detailed in example 50 a} in DME (70 ml), and the resulting orange coloured solution stirred at room temperature for 10 min. Aqueous 2M sodium carbonate (16.7 ml, 33.4 mmol) was then added, the mixture stirred for 10 min, then 4-chlorophenyl boronic acid (1.3 g, 8.36 mmol) was added, and the reaction mixture heated to 80°C for 18 h, under reflux. The reaction mixture was diluted with 1N HCl (100 ml) and the products extracted into ethyl acetate (2 x 100 ml). The combined organic extracts were washed with brine, dried (MgSO_4), and evaporated to give the crude product, which was purified by column chromatography on silica gel (20% ethyl acetate in n-heptane eluent) to give the product, (*E*)-ethyl-3-(4'-chloro-biphenyl-4-yl)-but-2-enoate as a colourless solid; 1.17 g (70%).

^1H NMR (300 MHz, CDCl_3) δ : 1.33 (3H, t), 2.60 (3H, s), 4.23 (2H, q), 6.20 (1H, s), 7.41 (2H, m), 7.52 (2H, m). MS: 300/302 (100%, M^+), 271/273, 255/257, 228/230, 165.

30

b)

A 1M solution of DIBAL-H in toluene (10 ml, 10 mmol) was added dropwise, at -70°C over 10 min, to a stirred solution of (*E*)-ethyl-3-(4'-chloro-biphenyl-4-yl)-but-2-enoate (1.0 g, 3.32 mmol) in dry THF (25 ml), and the mixture warmed to room temperature over 4 h. Methanol (1 ml) was carefully added, followed by 1N HCl (50 ml) and the resulting mixture extracted with ethyl acetate (2 x 50 ml). The combined organic extracts were washed with brine, dried (MgSO_4), and evaporated to give the product, (*E*)-3-(4'-chloro-biphenyl-4-yl)-but-2-en-1-ol as a colourless solid: 0.86 g (100%).


Mpt. $137-142^{\circ}\text{C}$. ^1H NMR (300 MHz, CDCl_3) δ : 1.79 (1H, br s), 2.11 (3H, d), 4.40 (2H, d), 6.05 (1H, tq), 7.41 (2H, dm), 7.45-7.60 (6H, m).

c)

Azodicarboxylic dipiperidide (0.731 g, 2.9 mmol) was added at $0-5^{\circ}\text{C}$ to a stirred solution of tributylphosphine (0.71 ml, 0.58 g, 2.9 mmol), (*S*)-ethyl 2-ethoxy-3-(4-hydroxyphenyl)-propionate (0.483 g, 2.03 mmol) and (*E*)-3-(4'-chloro-biphenyl-4-yl)-but-2-en-1-ol (0.500 g, 1.93 mmol) in dry benzene (15 ml), the mixture warmed to room temperature, and stirred for 3 h. The resulting mixture was diluted with water and ethyl acetate (30 ml each), the aqueous layer collected and further extracted with ethyl acetate (30 ml). The organic layers were combined, washed with brine, dried (MgSO_4) and evaporated. The crude product was then purified by column chromatography on silica (20% ethyl acetate in *n*-heptane eluent) to give (*E*)-(*S*)-ethyl 3-{4-[3-(4'-chloro-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate as a gum; 0.69 g (75%).

^1H NMR (300 MHz, CDCl_3) δ : 1.17 (3H, t), 1.23 (3H, t), 2.16 (3H, s), 2.96 (2H, d), 3.30-3.45 (1H, m), 3.55-3.68 (1H, m), 3.98 (1H, t), 4.17 (2H, q), 4.74 (2H, d), 6.12 (1H, t), 6.88 (2H, m), 7.18 (2H, m), 7.40 (2H, m), 7.45-7.60 (6H, m).

EXAMPLE 53




(E)-(S)-3-{4-[3-(4'-Chloro-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid

Sodium hydroxide (1M, 2.3 ml, 2.3 mmol) was added to a solution of (E)-(S)-ethyl 3-{4-[3-(4'-chloro-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate (example 52) (0.600 g, 1.25 mmol) in ethanol (10 ml) and the mixture stirred at room temperature for 18h, then heated to 80°C for 2 h. The resulting mixture was partitioned between water (50 ml) and ethyl acetate (50 ml) and the aqueous layer acidified to pH1 by addition of 1N HCl. The aqueous layer was separated and further extracted with ethyl acetate (2 x 50 ml). The combined organic layers were washed with brine, dried (MgSO₄), evaporated, and the product vacuum dried at 40°C for 72 h, to give (E)-(S)-3-{4-[3-(4'-chloro-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid as a colourless solid; 0.53 g (94%).

¹H NMR (300 MHz, CDCl₃) δ: 1.18 (3H, t), 2.16 (3H, s), 2.97 (1H, dd), 3.08 (1H, dd), 3.40-3.53 (1H, m), 3.55-3.68 (1H, m), 4.07 (1H, dd), 4.74 (2H, d), 6.11 (1H, t), 6.90 (2H, m), 7.17 (2H, m), 7.39 (2H, m), 7.45-7.60 (6H, m), carboxylic acid proton not observed.

EXAMPLE 54



(E)-(S)-Ethyl 2-Ethoxy-3-{4-[3-(5'-isopropyl-2'-methoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionate

25 a)

Tetrakis(triphenylphosphine)palladium(0) (0.20 g, 0.18 mmol, 4 mol%) was added, under nitrogen, to a stirred solution of (E)-3-(4-bromophenyl)-but-2-en-1-ol (1.0 g, 4.40 mmol) {prepared as detailed in example 50 b) in DME (55 ml), and the resulting orange coloured solution stirred at room temperature for 10 min. Aqueous 2M sodium carbonate (13.2 ml, 26.4


mmol) was then added, the mixture stirred for 10 min, then 5-isopropyl-2-methoxyphenylboronic acid (1.28 g, 6.60 mmol) was added, and the reaction mixture heated to 80°C for 18 h, under reflux. The reaction mixture was diluted with 1N HCl (100 ml) and the products extracted into ethyl acetate (2 x 100 ml). The combined organic extracts were
5 washed with brine, dried (MgSO₄), and evaporated to give the crude product, which was purified by column chromatography on silica gel (1% methanol in dichloromethane eluent) to give the product, 3-(5'-isopropyl-2'-methoxy-biphenyl-4-yl)-but-2-en-1-ol as a colourless oil; 1.15 g (88%).

¹H NMR (300 MHz, CDCl₃) δ: 1.26 (6H, d), 1.33 (1H, br t), 2.12 (3H, s), 2.91 (1H, septet),
10 3.80 (3H, s), 4.39 (2H, br t), 6.04 (1H, t), 6.92 (1H, d), 7.15-7.20 (2H, m), 7.42-7.55 (4H, m).
MS: 296 (100%, M⁺), 281, 263, 253.

b)

Azodicarboxylic dipiperidide (0.756 g, 3.0 mmol) was added at 0-5°C to a stirred solution of
15 tributylphosphine (0.74 ml, 0.61 g, 3.0 mmol), (S)-ethyl 2-ethoxy-3-(4-hydroxyphenyl)-propionate (0.50 g, 2.10 mmol) and 3-(5'-isopropyl-2'-methoxy-biphenyl-4-yl)-but-2-en-1-ol (0.593 g, 2.0 mmol) in dry benzene (15 ml), the mixture warmed to room temperature, and stirred for 4 h. The resulting mixture was diluted with water (100 ml) and ethyl acetate (50 ml), the aqueous layer collected and further extracted with ethyl acetate (50 ml). The organic
20 layers were combined, washed with brine, dried (MgSO₄) and evaporated. The crude product was then purified by column chromatography on silica (10% ethyl acetate in n-heptane eluent) to give (E)-(S)-ethyl 2-ethoxy-3-{4-[3-(5'-isopropyl-2'-methoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionate as a colourless oil; 0.67 g (65%).

¹H NMR (300 MHz, CDCl₃) δ: 1.17 (3H, t), 1.22 (3H, t), 1.26 (6H, d), 2.16 (3H, s), 2.91 (1H, septet), 2.96 (2H, d), 3.30-3.45 (1H, m), 3.54-3.66 (1H, m), 3.79 (3H, s), 3.98 (1H, t), 4.17
25 (2H, q), 4.74 (2H, d), 6.10 (1H, t), 6.84-6.95 (3H, m), 7.12-7.20 (4H, m), 7.42-7.57 (4H, m).


EXAMPLE 55

5 *(E)-(S)*-2-Ethoxy-3-{4-[3-(5'-isopropyl-2'-methoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionic acid

The title compound was prepared from *(E)-(S)*-ethyl 2-ethoxy-3-{4-[3-(5'-isopropyl-2'-methoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionate (example 54) (0.50 g, 0.968 mmol) and sodium hydroxide (1M, 1.93 ml, 1.93 mmol) by a procedure analogous to that described in example 51, yielding *(E)-(S)*-2-ethoxy-3-{4-[3-(5'-isopropyl-2'-methoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionic acid as a colourless gum, which contained 0.44 mol equivalents of ethyl acetate; 0.48 g (94%).

15 ¹H NMR (300 MHz, CDCl₃) δ: 1.18 (3H, t), 1.26 (6H, d), 1.26 (ethyl acetate impurity, 1.32H, t), 2.04 (ethyl acetate impurity, 0.88H, s), 2.16 (3H, s), 2.82-3.02 (2H, m), 3.08 (1H, dd), 3.40-3.52 (1H, m), 3.52-3.68 (1H, m), 3.79 (3H, s), 4.06 (1H, dd), 4.15 (ethyl acetate impurity, 0.88H, q), 4.75 (2H, d), 6.09 (1H, t), 6.88-6.95 (3H, m), 7.12-7.20 (4H, m), 7.42-7.57 (4H, m), carboxylic acid proton not observed.

20

EXAMPLE 56

(*E*)-(*S*)-Ethyl 2-Ethoxy-3-{4-[3-(5'-chloro-2'-methoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionate

a)

- 5 (*E*)-Ethyl 3-(5'-chloro-2'-methoxy-biphenyl-4-yl)-but-2-enoate (1.07 g, 91% yield) was prepared from 5-chloro-2-methoxyphenylboronic acid (1.0 g, 5.36 mmol) and (*E*)-ethyl 3-(4-bromophenyl)-but-2-enoate (0.96 g, 3.57 mmol) by a procedure analogous to that described in example 52 a.

¹H NMR (300 MHz, CDCl₃) δ: 1.33 (3H, t), 2.60 (3H, s), 3.81 (3H, s), 4.23 (2H, q), 6.20 (1H, s), 6.91 (1H, d), 7.25-7.33 (2H, m), 7.47-7.57 (4H, m). MS: 330/332 (100%, M⁺).

10

b)


- (*E*)-Ethyl 3-(5'-chloro-2'-methoxy-biphenyl-4-yl)-but-2-enoate (0.90 g, 2.72 mmol) was reduced with DIBAL-H by a procedure analogous to that described in example 52 b to give (*E*)-3-(5'-chloro-2'-methoxy-biphenyl-4-yl)-but-2-en-1-ol as a colourless oil; 0.785 g (100%).
- 15

¹H NMR (300 MHz, CDCl₃) δ: 1.49 (1H, br s), 2.11 (3H, s), 3.80 (3H, s), 4.39 (2H, d), 6.04 (1H, t), 6.90 (1H, d), 7.22-7.32 (2H, m), 7.47-7.57 (4H, m). MS: 288/290 (100%, M⁺), 270/272, 255/257, 245/247.

20 c)

The title compound (0.54 g, 61% yield) was prepared from (*E*)-3-(5'-chloro-2'-methoxy-biphenyl-4-yl)-but-2-en-1-ol (0.50 g, 1.73 mmol) and (*S*)-ethyl 2-ethoxy-3-(4-hydroxyphenyl)-propionate (0.433 g, 1.82 mmol) by a procedure analogous to that described in example 52 c.

- 25 ¹H NMR (300 MHz, CDCl₃) δ: 1.17 (3H, t), 1.23 (3H, t), 2.16 (3H, s), 2.96 (2H, d), 3.30-3.43 (1H, m), 3.55-3.65 (1H, m), 3.79 (3H, s), 3.98 (1H, t), 4.17 (2H, q), 4.74 (2H, d), 6.10 (1H, t), 6.84-6.92 (3H, m), 7.12-7.20 (2H, m), 7.22-7.32 (2H, m), 7.45-7.50 (4H, m).
LCMS: 331/333 (M+Na).

EXAMPLE 57


5 *(E)-(S)*-3-{4-[3-(5'-Chloro-2'-methoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid

The title compound was prepared from *(E)-(S)*-ethyl 2-ethoxy-3-{4-[3-(5'-chloro-2'-methoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionate (example 56) (0.47 g, 0.92 mmol) and so-
 10 dium hydroxide (1M, 1.8 ml, 1.8 mmol) by a procedure analogous to that described in exam-
 ple 51, yielding *(E)-(S)*-3-{4-[3-(5'-chloro-2'-methoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-
 ethoxy-propionic acid as a colourless gum, which contained 0.2 mol equivalents of ethyl ace-
 tate; 0.43 g (98%).

¹H NMR (300 MHz, CDCl₃) δ: 1.19 (3H, t), 1.26 (ethyl acetate impurity, 0.6H, t), 2.04 (ethyl
 15 acetate impurity, 0.4H, s), 2.16 (3H, s), 2.96 (1H, dd), 3.10 (1H, dd), 3.42-3.52 (1H, m), 3.53-
 3.68 (1H, m), 3.80 (3H, s), 4.07 (1H, dd), 4.12 (ethyl acetate impurity, 0.4H), 4.74 (2H, d),
 6.10 (1H, t), 6.85-6.95 (3H, m), 7.12-7.20 (2H, m), 7.21-7.32 (2H, m), 7.45-7.50 (4H, m), car-
 boxylic acid proton not observed.

LCMS: 503/505 (M+Na).

20

EXAMPLE 58

25 *(E)-(S)*-Ethyl 3-{4-[3-(2',3'-Dichloro-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate

a)

(*E*)-Ethyl 3-(2',3'-Dichloro-biphenyl-4-yl)-but-2-enoate (1.07 g, 73% yield) was prepared from 2,3-dichlorophenylboronic acid (1.26 g, 6.60 mmol) and (*E*)-ethyl 3-(4-bromophenyl)-but-2-enoate (1.0 g, 4.40 mmol) by a procedure analogous to that described in example 52 a.

5 Mpt. 64-66°C. ¹H NMR (300 MHz, CDCl₃) δ: 1.33 (3H, t), 2.62 (3H, d), 4.23 (2H, q), 6.21 (1H, m), 7.20-7.30 (2H, m), 7.40-7.50 (3H, m), 7.50-7.60 (2H, m). MS: 334/336/338 (100%, M⁺), 305/307/309, 289/291/293, 262/264/266, 189/191.

b)

10 (*E*)-Ethyl 3-(2',3'-dichloro-biphenyl-4-yl)-but-2-enoate (1.07 g, 3.19 mmol) was reduced with DIBAL-H by a procedure analogous to that described in example 52 b to give (*E*)-3-(2',3'-dichloro-biphenyl-4-yl)-but-2-en-1-ol as a colourless solid; 0.74 g (79%).

Mpt. 95-100°C. ¹H NMR (300 MHz, CDCl₃) δ: 1.45 (1H, br s), 2.13 (3H, s), 4.40 (2H, d), 6.07 (1H, t), 7.20-7.28 (2H, m), 7.35-7.42 (2H, m), 7.42-7.53 (3H, m).


15

c)

The title compound (0.41 g, 80% yield) was prepared from (*E*)-3-(2',3'-dichloro-biphenyl-4-yl)-but-2-en-1-ol (0.293 g, 1.0 mmol) and (*S*)-ethyl 2-ethoxy-3-(4-hydroxyphenyl)-propionate (0.25 g, 1.05 mmol) by a procedure analogous to that described in example 52 c.

20 ¹H NMR (300 MHz, CDCl₃) δ: 1.17 (3H, t), 1.23 (3H, t), 2.18 (3H, s), 2.96 (2H, d), 3.30-3.43 (1H, m), 3.55-3.65 (1H, m), 3.98 (1H, t), 4.17 (2H, q), 4.75 (2H, d), 6.13 (1H, t), 6.84-6.92 (2H, m), 7.12-7.20 (2H, m), 7.21-7.32 (2H, m), 7.35-7.42 (2H, m), 7.43-7.53 (3H, m).

25


EXAMPLE 59

(*E*)-(S)-3-{4-[3-(2',3'-Dichloro-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid

The title compound was prepared from (*E*)-(*S*)-ethyl 3-{4-[3-(2',3'-dichloro-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate (example 58) (0.325 g, 0.63 mmol) and sodium hydroxide (1M, 1.27 ml, 1.27 mmol) by a procedure analogous to that described in example 51, giving (*E*)-(*S*)-3-{4-[3-(2',3'-dichloro-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid as a gum, which contained 0.28 mol equivalents of ethyl acetate; 0.24 g (80%).

¹H NMR (300 MHz, CDCl₃) δ: 1.19 (3H, t), 1.26 (ethyl acetate impurity, 0.84H, t), 2.05 (ethyl acetate impurity, 0.56H, s), 2.18 (3H, m), 2.98 (1H, dd), 3.08 (1H, dd), 3.42-3.52 (1H, m), 3.53-3.68 (1H, m), 3.80 (3H, s), 4.07 (1H, dd), 4.12 (ethyl acetate impurity, 0.56H), 4.75 (2H, d), 6.13 (1H, tm), 6.85-6.95 (2H, m), 7.14-7.20 (2H, m), 7.21-7.30 (2H, m), 7.35-7.42 (2H, m), 7.42-7.53 (3H, m), carboxylic acid proton not observed.

EXAMPLE 60



(*E*)-(*S*)-Ethyl 3-{4-[3-(2',6'-Dimethoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate

20 a)

(*E*)-Ethyl 3-(2',6'-dimethoxy-biphenyl-4-yl)-but-2-enoate (1.02 g, 73% yield) was prepared from 2,6-dimethoxyphenylboronic acid (1.20 g, 6.60 mmol) and (*E*)-ethyl 3-(4-bromophenyl)-but-2-enoate (1.0 g, 4.40 mmol) by a procedure analogous to that described in example 52 a. Mpt. 120-123.5°C. ¹H NMR (300 MHz, CDCl₃) δ: 1.32 (3H, t), 2.62 (3H, d), 3.75 (6H, s), 4.23 (2H, q), 6.22 (1H, m), 6.67 (2H, d), 7.29 (1H, t), 7.38 (2H, dm), 7.53 (2H, dm). MS: 326 (100%, M⁺), 297, 281.

b)

(*E*)-Ethyl 3-(2',6'-dimethoxy-biphenyl-4-yl)-but-2-enoate (0.90 g, 2.76 mmol) was reduced with DIBAL-H by a procedure analogous to that described in example 52 b to give (*E*)-3-(2',6'-dimethoxy-biphenyl-4-yl)-but-2-en-1-ol as a colourless solid; 0.82 g (100%).


Mpt. 70-75°C. ¹H NMR (300 MHz, CDCl₃) δ: 1.44 (1H, br s), 2.12 (3H, d), 3.74 (6H, s), 4.38 (2H, d), 6.06 (1H, tm), 6.66 (2H, d), 7.13-7.37 (3H, m), 7.42-7.50 (2H, m). MS: 284 (100%, M⁺), 266, 251, 241.

c)

The title compound (0.41 g, 80% yield) was prepared from (*E*)-3-(2',6'-dimethoxy-biphenyl-4-yl)-but-2-en-1-ol (0.50 g, 1.76 mmol) and (*S*)-ethyl 2-ethoxy-3-(4-hydroxyphenyl)-propionate (0.44 g, 1.85 mmol) by a procedure analogous to that described in example 52 c.

¹H NMR (300 MHz, CDCl₃) δ: 1.17 (3H, t), 1.22 (3H, t), 2.16 (3H, m), 2.96 (2H, d), 3.30-3.43 (1H, m), 3.53-3.65 (1H, m), 3.73 (6H, s), 3.97 (1H, t), 4.17 (2H, q), 4.75 (2H, d), 6.10 (1H, tm), 6.66 (2H, d), 6.84-6.90 (2H, m), 7.13-7.20 (2H, m), 7.27 (1H, t), 7.30-7.38 (2H, m), 7.45-7.52 (2H, m). LCMS: 527 (M+Na), 267 (100%).

EXAMPLE 61




(*E*)-(S)-3-{4-[3-(2',6'-Dimethoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid

The title compound was prepared from (*E*)-(S)-ethyl 3-{4-[3-(2',6'-dimethoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate (example 60) (0.565 g, 1.12 mmol) and sodium hydroxide (1M, 2.20 ml, 2.20 mmol) by a procedure analogous to that described in example 51; giving (*E*)-(S)-3-{4-[3-(2',6'-dimethoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid as a gum; 0.49 g (92%).

¹H NMR (300 MHz, CDCl₃) δ: 1.18 (3H, t), 2.16 (3H, m), 2.98 (1H, dd), 3.08 (1H, dd), 3.42-3.52 (1H, m), 3.53-3.68 (1H, m), 3.73 (6H, s), 4.07 (1H, dd), 4.75 (2H, d), 6.10 (1H, tm), 6.66

(2H, d), 6.86-6.92 (2H, m), 7.13-7.20 (2H, m), 7.27 (1H, t), 7.28-7.35 (2H, m), 7.45-7.50 (2H, m), carboxylic acid proton not observed. LCMS: 499 (M^+), 267 (100%).

5

EXAMPLE 62

(E)-(S)-Ethyl 3-{4-[3-(4-Bromophenyl)-2-methyl-but-2-enyloxy]-phenyl}-2-ethoxy-propionate

10

a)

Sodium (1.37 g, 59.6 mmol) was added to ethanol (50 ml) at 20°C and the mixture stirred until the metal had fully reacted. Triethyl phosphonoacetate (17.78 g, 74.62 mmol) was added, the mixture stirred for 20 min, then a solution of 4-bromoacetophenone (9.90 g, 49.74 mmol) in ethanol (50 ml) was added and the reaction mixture heated to 80°C under reflux for 17h. The solution was cooled, the ethanol evaporated and the residue partitioned between 1N HCl (100 ml) and ethyl acetate (100 ml). The aqueous layer was collected and further extracted with ethyl acetate (2 x 200 ml). The organic layers were combined, washed with brine, dried ($MgSO_4$) and evaporated to an orange gum. This was purified by column chromatography on silica gel (2% diethyl ether in n-heptane eluent) to give the two double-bond isomer products as colourless oils.

20

(E)-Ethyl 3-(4-bromophenyl)-2-methyl-but-2-enoate; 5.38 g (38%).

25

1H NMR (300 MHz, $CDCl_3$) δ : 1.34 (3H, t), 1.75 (3H, m), 2.22 (3H, m), 4.26 (2H, q), 7.04 (2H, dm), 7.49 (2H, dm). MS: 282/284 (M^+), 253/255, 237/239, 208/210, 175, 157, 130, 129 (100%), 115.

And

(Z)-Ethyl 3-(4-bromophenyl)-2-methyl-but-2-enoate; 3.15 g (22%).

30

1H NMR (300 MHz, $CDCl_3$) δ : 0.90 (3H, t), 2.01 (3H, s), 2.06 (3H, s), 3.88 (2H, q), 7.00 (2H, dm), 7.41 (2H, dm). MS: 282/284 (M^+), 253/255, 237/239, 208/210, 157, 130, 129 (100%), 115.

b)

(*E*)-Ethyl 3-(4-bromophenyl)-2-methyl-but-2-enoate (2.83 g, 9.99 mmol) was reduced with DIBAL-H by a procedure analogous to that described in example 52b to give (*E*)-3-(4-bromophenyl)-2-methyl-but-2-en-1-ol as a colourless oil; 1.82 g (75%).


¹H NMR (300 MHz, CDCl₃) δ: 1.60 (1H, br s), 1.66 (3H, m), 2.00 (3H, m), 4.29 (2H, s), 7.01 (2H, dm), 7.44 (2H, dm). MS: 240/242 (M⁺), 225/227, 183/185 (100%), 161, 146, 143, 128, 115.

c)

The title compound (0.83 g, 87% yield) was prepared from (*E*)-3-(4-bromophenyl)-2-methyl-but-2-en-1-ol (0.50 g, 2.07 mmol) and (*S*)-ethyl 2-ethoxy-3-(4-hydroxyphenyl)-propionate (0.519 g, 2.18 mmol) by a procedure analogous to that described in example 52c.

¹H NMR (300 MHz, CDCl₃) δ: 1.18 (3H, t), 1.23 (3H, t), 1.68 (3H, m), 2.04 (3H, m), 2.97 (2H, d), 3.30-3.43 (1H, m), 3.53-3.68 (1H, m), 3.98 (1H, t), 4.18 (2H, q), 4.61 (2H, s), 6.88 (2H, dm), 7.04 (2H, dm), 7.17 (2H, dm), 7.45 (2H, dm).

EXAMPLE 63




(*E*)-(S)-3-{4-[3-(4-Bromophenyl)-2-methyl-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid

The title compound was prepared from (*E*)-(S)-ethyl 3-{4-[3-(4-bromophenyl)-2-methyl-but-2-enyloxy]-phenyl}-2-ethoxy-propionate (example 62) (0.710 g, 1.54 mmol) and sodium hydroxide (1M, 3.10 ml, 3.10 mmol) by a procedure analogous to that described in example 51; giving (*E*)-(S)-3-{4-[3-(4-bromophenyl)-2-methyl-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid as a colourless solid, which contained approximately 13 mol% of ethyl acetate impurity; 0.67 g (98%).

¹H NMR (300 MHz, CDCl₃) δ: 1.19 (3H, t), 1.68 (3H, m), 2.04 (3H, m), 2.98 (1H, dd), 3.08 (1H, dd), 3.42-3.54 (1H, m), 3.54-3.68 (1H, m), 4.07 (1H, dd), 4.61 (2H, s), 6.90 (2H, dm), 7.04 (2H, dm), 7.17 (2H, dm), 7.45 (2H, dm), carboxylic acid proton not observed.

5

EXAMPLE 64

10 (Z)-(S)-Ethyl 3-{4-[3-(4-Bromophenyl)-2-methyl-but-2-enyloxy]-phenyl}-2-ethoxy-propionate

a)

(Z)-Ethyl 3-(4-bromophenyl)-2-methyl-but-2-enoate (1.42 g, 5.01 mmol), which was prepared as described in example 62 a, was reduced with DIBAL-H by a procedure analogous to that described in example 52 b to give (Z)-3-(4-bromophenyl)-2-methyl-but-2-en-1-ol as a colourless oil; 1.19 g (98%).

15


¹H NMR (300 MHz, CDCl₃) δ: 1.38 (1H, br s), 1.89 (3H, s), 1.97 (3H, s), 3.92 (2H, s), 7.01 (2H, dm), 7.42 (2H, dm). MS: 240/242 (M⁺), 225/227, 183/185 (100%), 161, 146, 143, 128, 115.

20

c)

The title compound (0.91 g, 95% yield) was prepared from (Z)-3-(4-bromophenyl)-2-methyl-but-2-en-1-ol (0.50 g, 2.07 mmol) and (S)-ethyl 2-ethoxy-3-(4-hydroxyphenyl)-propionate (0.519 g, 2.18 mmol) by a procedure analogous to that described in example 52 c.


25 ¹H NMR (300 MHz, CDCl₃) δ: 1.16 (3H, t), 1.21 (3H, t), 1.93 (3H, s), 2.02 (3H, s), 2.93 (2H, d), 3.28-3.42 (1H, m), 3.53-3.68 (1H, m), 3.95 (1H, t), 4.16 (2H, q), 4.25 (2H, s), 6.69 (2H, dm), 7.04 (2H, dm), 7.09 (2H, dm), 7.41 (2H, dm).

EXAMPLE 65

5 (Z)-(S)-3-{4-[3-(4-Bromophenyl)-2-methyl-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid

The title compound was prepared from (Z)-(S)-ethyl 3-{4-[3-(4-bromophenyl)-2-methyl-but-2-enyloxy]-phenyl}-2-ethoxy-propionate (example 64) (0.82 g, 1.78 mmol) and sodium hydroxide (1M, 3.60 ml, 3.60 mmol) by a procedure analogous to that described in example 51; giving
 10 (Z)-(S)-3-{4-[3-(4-bromophenyl)-2-methyl-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid as a colourless solid, which contained approximately 15 mol% of ethyl acetate impurity; 0.766 g (100%).

¹H NMR (300 MHz, CDCl₃) δ: 1.17 (3H, t), 1.93 (3H, s), 2.02 (3H, s), 2.93 (1H, dd), 3.04 (1H, dd), 3.40-3.52 (1H, m), 3.52-3.65 (1H, m), 4.03 (1H, dd), 4.26 (2H, s), 6.71 (2H, dm), 7.04.
 15 (2H, dm), 7.09 (2H, dm), 7.41 (2H, dm), carboxylic acid proton not observed.

EXAMPLE 66

20 (E)-(S)-Ethyl 2-Ethoxy-3-[4-(3-[1,1';3',1'']terphenyl-4''-yl)-but-2-enyloxy]-phenyl-propionate

a)

(*E*)-Ethyl 3-[1,1';3',1'']terphenyl-4''-yl-but-2-enoate (1.02 g, 68% yield) was prepared from 3-biphenylboronic acid (1.31 g, 6.60 mmol) and (*E*)-ethyl 3-(4-bromophenyl)-but-2-enoate (1.0 g, 4.40 mmol) by a procedure analogous to that described in example 52 a.

¹H NMR (300 MHz, CDCl₃) δ: 1.33 (3H, t), 2.62 (3H, d), 4.23 (2H, q), 6.21 (1H, s), 7.30-7.70 (12H, m), 7.82 (1H, m). LCMS: 343 (100%, M⁺), 297.

b)

(*E*)-Ethyl 3-[1,1';3',1'']terphenyl-4''-yl-but-2-enoate (0.95 g, 2.77 mmol) was reduced with DI-BAL-H by a procedure analogous to that described in example 52 b to give (*E*)-3-

[1,1';3',1'']terphenyl-4''-yl-but-2-en-1-ol as a colourless solid; 0.81 g (97%).

¹H NMR (300 MHz, CDCl₃) δ: 1.37 (1H, br s), 2.13 (3H, s), 4.40 (2H, d), 6.06 (1H, tm), 7.30-7.70 (12H, m), 7.81 (1H, m). LCMS: 283 (100%, M+H-H₂O).

Microanalysis Calculated % C: 87.96, H: 6.71. Found % C: 87.85, H: 6.74.


c)

The title compound (0.41 g, 80% yield) was prepared from (*E*)-3-[1,1';3',1'']terphenyl-4''-yl-but-2-en-1-ol (0.30 g, 1.0 mmol) and (*S*)-ethyl 2-ethoxy-3-(4-hydroxyphenyl)-propionate (0.25 g, 1.05 mmol) by a procedure analogous to that described in example 52 c.

¹H NMR (300 MHz, CDCl₃) δ: 1.17 (3H, t), 1.22 (3H, t), 2.18 (3H, s), 2.96 (2H, d), 3.30-3.43 (1H, m), 3.55-3.68 (1H, m), 3.98 (1H, t), 4.17 (2H, q), 4.75 (2H, d), 6.13 (1H, t), 6.89 (2H, dm), 7.17 (2H, dm), 7.30-7.70 (12H, m), 7.81 (1H, m).

Microanalysis Calculated % C: 80.74, H: 6.97. Found % C: 80.84, H: 7.28.

EXAMPLE 67




(*E*)-(S)-2-Ethoxy-3-[4-(3-[1,1';3',1'']terphenyl-4''-yl-but-2-enyloxy)-phenyl]-propionic acid

The title compound was prepared from (*E*)-(*S*)-ethyl 2-ethoxy-3-[4-(3-[1,1';3',1'']terphenyl-4"-yl-but-2-enyloxy)-phenyl]-propionate (example 66) (0.185 g, 0.36 mmol) and sodium hydroxide (1M, 0.71 ml, 0.71 mmol) by a procedure analogous to that described in example 51; giving (*E*)-(*S*)-2-ethoxy-3-[4-(3-[1,1';3',1'']terphenyl-4"-yl-but-2-enyloxy)-phenyl]-propionic acid as a gum; 0.145 g (83%).

¹H NMR (300 MHz, CDCl₃) δ: 1.19 (3H, t), 2.18 (3H, m), 2.99 (1H, dd), 3.09 (1H, dd), 3.40-3.53 (1H, m), 3.53-3.68 (1H, m), 4.07 (1H, dd), 4.75 (2H, d), 6.13 (1H, tm), 6.90 (2H, dm), 7.17 (2H, dm), 7.30-7.70 (12H, m), 7.81 (1H, m), carboxylic acid proton not observed.

EXAMPLE 68



15 (*E*)-(*S*)-Ethyl 2-Ethoxy-3-{4-[3-(3'-methyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionate

a)

(*E*)-3-(3'-Methyl-biphenyl-4-yl)-but-2-enoate (0.795 g, 65% yield) was prepared from 3-tolylboronic acid (0.90 g, 6.60 mmol) and (*E*)-ethyl 3-(4-bromophenyl)-but-2-enoate (1.0 g, 4.40 mmol) by a procedure analogous to that described in example 52 a.

¹H NMR (300 MHz, CDCl₃) δ: 1.33 (3H, t), 2.43 (3H, s), 2.61 (3H, s), 4.23 (2H, q), 6.20 (1H, s), 7.18 (1H, dm), 7.34 (1H, tm), 7.41 (2H, dm), 7.52-7.63 (4H, m). LCMS: 281 (M+H), 235 (100%).

25 b)

(*E*)-3-(3'-Methyl-biphenyl-4-yl)-but-2-enoate (0.74 g, 2.64 mmol) was reduced with DIBAL-H by a procedure analogous to that described in example 52 b to give (*E*)-3-(3'-methyl-biphenyl-4-yl)-but-2-en-1-ol as a colourless solid; 0.63 g (85%).

¹H NMR (300 MHz, CDCl₃) δ: 1.36 (1H, br s), 2.12 (3H, s), 2.42 (3H, s), 4.39 (2H, d), 6.05 (1H, tm), 7.16 (1H, dm), 7.33 (1H, tm), 7.40 (2H, dm), 7.48 (2H, dm), 7.56 (2H, dm). LCMS: 221 (100%, M+H-H₂O).

5 c)


The title compound (0.365 g, 78% yield) was prepared from (*E*)-3-(3'-methyl-biphenyl-4-yl)-but-2-en-1-ol (0.238 g, 1.0 mmol) and (*S*)-ethyl 2-ethoxy-3-(4-hydroxyphenyl)-propionate (0.25 g, 1.05 mmol) by a procedure analogous to that described in example 52 c.

¹H NMR (300 MHz, CDCl₃) δ: 1.17 (3H, t), 1.22 (3H, t), 2.17 (3H, s), 2.42 (3H, s), 2.96 (2H, d), 3.30-3.43 (1H, m), 3.55-3.68 (1H, m), 3.98 (1H, t), 4.17 (2H, q), 4.74 (2H, d), 6.11 (1H, t), 6.90 (2H, dm), 7.13-7.23 (3H, m), 7.33 (1H, t), 7.36-7.44 (2H, m), 7.45-7.60 (4H, m).

Microanalysis Calculated % C: 78.57, H: 7.47. Found % C: 78.90, H: 7.70.

15

EXAMPLE 69




(*E*)-(*S*)-2-Ethoxy-3-{4-[3-(3'-methyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionic acid

20

The title compound was prepared from (*E*)-(*S*)-2-ethoxy-3-{4-[3-(3'-methyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionate (example 68) (0.225 g, 0.49 mmol) and sodium hydroxide (1M, 0.98 ml, 0.98 mmol) by a procedure analogous to that described in example 51; giving (*E*)-(*S*)-2-ethoxy-3-{4-[3-(3'-methyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionic acid as a gum; 0.20 g (95%).

25

¹H NMR (300 MHz, CDCl₃) δ: 1.18 (3H, t), 2.17 (3H, m), 2.42 (3H, s), 2.97 (1H, dd), 3.09 (1H, dd), 3.42-3.54 (1H, m), 3.55-3.68 (1H, m), 4.07 (1H, dd), 4.75 (2H, d), 6.11 (1H, tm), 6.90 (2H, dm), 7.10-7.23 (3H, m), 7.35 (1H, t), 7.37-7.44 (2H, m), 7.45-7.60 (4H, m), carboxylic acid proton not observed.

EXAMPLE 70

5

(E)-(S)-Ethyl 3-(4-[3-(3'-Acetyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl)-2-ethoxy-propionate

a)

- 10 3-Acetylphenylboronic acid (7.10 g, 43.3 mmol) was coupled with (E)-3-(4-bromophenyl)-but-2-en-1-ol (5.76 g, 25.0 mmol) by a procedure analogous to that described in example 54 a to give (E)-3-(3'-acetyl-biphenyl-4-yl)-but-2-en-1-ol as an off-white solid; 5.33 g (79%). This solid was recrystallised from aqueous ethanol to give a first crop of very pure (E)-3-(3'-acetyl-biphenyl-4-yl)-but-2-en-1-ol as colourless platelets; 2.78 g (41%) and a second crop of (E)-3-(3'-acetyl-biphenyl-4-yl)-but-2-en-1-ol as an amorphous off-white solid; 2.53 g (37%).
- 15


Mpt. 85-86°C. ¹H NMR (300 MHz, CDCl₃) δ: 1.46 (1H, br t), 2.13 (3H, d), 2.66 (3H, s), 4.41 (2H, br t), 6.07 (1H, tm), 7.50-7.62 (5H, m), 7.80 (1H, dm); 7.92 (1H, dm), 8.19 (1H, m). MS: 266 (M⁺), 251, (M-Me), 248 (M-H₂O), 223 (100%). Microanalysis Calculated % C: 81.17, H: 6.81. Found % C: 81.22, H: 6.83.

20

b)

The title compound (0.16 g, 65% yield) was prepared from (E)-3-(3'-acetyl-biphenyl-4-yl)-but-2-en-1-ol (0.133 g, 0.50 mmol) and (S)-ethyl 2-ethoxy-3-(4-hydroxyphenyl)-propionate (0.125 g, 0.525 mmol) by a procedure analogous to that described in example 52 c.

- 25 ¹H NMR (300 MHz, CDCl₃) δ: 1.17 (3H, t), 1.23 (3H, t), 2.18 (3H, s), 2.66 (3H, s), 2.92 (2H, d), 3.30-3.43 (1H, m), 3.55-3.68 (1H, m), 3.98 (1H, t), 4.17 (2H, q), 4.75 (2H, d), 6.13 (1H, t), 6.89 (2H, dm), 7.17 (2H, dm), 7.50-7.64 (5H, m), 7.80 (1H, dm), 7.92 (1H, dm), 8.19 (1H, m).

EXAMPLE 71

5

(*E*)-(*S*)-2-Ethoxy-3-{4-[3-(5'-isopropyl-2'-methoxy-biphenyl-4-yl)-allyloxy]-phenyl}-propionic acid ethyl ester

10 a)

(*E*)-3-(4-Bromo-phenyl)-acrylic acid ethyl ester was prepared from 4-bromobenzaldehyde (20.0 g, 0.11 mol) by a procedure analogous to that described in example 23 a.

b)

15 (*E*)-3-(4-Bromo-phenyl)-acrylic acid ethyl ester (450 mg, 2.0 mmol) was reacted with 5-isopropyl-2-methoxy-benzene boronic acid (776 mg, 4.0 mmol) by a procedure described in example 52 a, to give (*E*)-3-(5'-Isopropyl-2'-methoxy-biphenyl-4-yl)-acrylic acid ethyl ester.

c)

20 (*E*)-3-(5'-Isopropyl-2'-methoxy-biphenyl-4-yl)-acrylic acid ethyl ester was reduced by DIBAL-H by a procedure analogous to that described in example 52 b to give (*E*)-3-(5'-isopropyl-2'-methoxy-biphenyl-4-yl)-prop-2-en-1-ol.


25

d)

The title compound was prepared from (*E*)-3-(5'-isopropyl-2'-methoxy-biphenyl-4-yl)-prop-2-en-1-ol by a procedure analogous to that described in 52 c.

¹H NMR (300 MHz, CDCl₃) δ: 1.13-1.30 (m, 12H), 2.85-3.0 (m, 3H), 3.30-3.42 (m, 1H), 3.53-3.67 (m, 1H), 2.78 (s, 3H), 3.98 (t, 1H), 4.15 (q, 2H), 4.70 (dd, 2H), 6.43 (dt, 1H), 6.75 (d, 1H), 6.85-6.95 (m, 3H), 7.15 (d, 4H), 7.44 (d, 2H), 7.52 (d, 2H).

EXAMPLE 72




(*E*)-(S)-2-Ethoxy-3-{4-[3-(5'-isopropyl-2'-methoxy-biphenyl-4-yl)-allyloxy]-phenyl}-propionic acid

15

The title compound was prepared from (*E*)-(S)-2-ethoxy-3-{4-[3-(5'-isopropyl-2'-methoxy-biphenyl-4-yl)-allyloxy]-phenyl}-propionic acid ethyl ester (example 71) (370 mg, 0.78 mmol) by a procedure analogous to that described in example 26.

¹H NMR (CDCl₃, 300 MHz) δ: 1.18 (t, 3H), 1.26 (d, 6H), 2.85-3.03 (m, 2H), 3.08 (dd, 1H), 3.35-3.48 (m, 1H), 3.55-3.68 (m, 1H), 3.75 (s, 3H), 4.03 (dd, 1H), 4.67 (d, 2H), 6.43 (dt, 1H), 6.75 (d, 1H), 6.87-6.95 (m, 3H), 7.13-7.23 (m, 4H), 7.43 (d, 2H), 7.53 (d, 2H).

20

EXAMPLE 73

5 (E)-(S)-3-{4-[3-(3,5-Distyryl-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester


a)

Bu₄NBr (2.0 g, 6.3 mmol), K₂CO₃ (7.8 g, 56.7 mmol), Pd(OAc)₂ (250 mg, 1.1 mmol) and styrene (20 mL, 175 mmol) were stirred for 5 min under nitrogen. To the mixture was added 3,5-dibromobenzaldehyde (5.0 g, 18.9 mmol) in dry DMF (5.0 mL), and the mixture was stirred at
10 65°C for 16h. The reaction mixture was diluted with ethyl acetate (20 mL) and the solution filtered. The filtrate was diluted with water and extracted with ethyl acetate (3 x 50 mL). The organic layers were combined, dried over MgSO₄, and concentrated under vacuum. To the residue was added a mixture of toluene/petroleum ether (1:1) (50 mL) and 3,5-distyryl-
15 benzaldehyde (4.95 g, 85%) was isolated by filtration.

b)

The title compound was prepared from 3,5-distyryl-benzaldehyde (3.8 g, 10.0 mmol) by a sequence analogous to that described in example 23 b-c.

20 ¹H NMR (CDCl₃, 300 MHz) δ: 1.15 (t, 3H), 1.22 (t, 3H), 2.98 (d, 2H), 3.32-3.42 (m, 1H), 3.55-3.68 (m, 1H), 3.98 (t, 1H), 4.12 (t, 1H), 4.18 (q, 2H), 4.72 (dd, 2H), 6.50 (dt, 1H), 6.78 (d, 1H), 6.90 (d, 1H), 7.08-7.32 (m, 8H), 7.39 (t, 4H), 7.45 (s, 2H), 7.53 (d, 5H).


EXAMPLE 74

5 *(E)-(S)-3-{4-[3-(3,5-Distyryl-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid*

(E)-(S)-3-{4-[3-(3,5-Distyryl-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester (example 73) (335 mg, 0.6 mmol) was dissolved in warm ethanol (20 mL) and sodium hydroxide (1N, 0.9 mL, 0.9 mmol) added. The mixture was stirred at room temperature for 16h. The title
10 compound as a sodium salt was isolated by filtration and washed with ethanol/water (10:1), yielding 190 mg (57%).

¹H NMR (CDCl₃, 300 MHz) δ : 0.98 (t, 3H), 2.63 (dd, 1H), 2.85 (dd, 1H), 3.05-3.15 (m, 1H), 3.50-3.64 (m, 2H), 4.75 (d, 2H), 6.68 (dt, 1H), 6.80 (d, 1H), 6.90 (d, 2H), 7.15 (d, 2H), 7.25-7.48 (m, 10H), 7.60-7.70 (m, 6H), 7.75 (s, 1H).

15

EXAMPLE 75

20

(E)-(S)-3-{4-[3-(3,5-Diisopropoxy-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester

a)


To a solution of 3,5-dihydroxybenzaldehyde (3.0 g, 22.0 mmol) in DMF (17 mL) was added potassium carbonate (12.1 g, 87.0 mmol) and 2-bromopropane (28.5 g, 232 mmol). The reaction mixture was heated at 100°C for 3 h. The mixture was filtered and washed with ethyl acetate. The filtrate was added water and the organic phase isolated. The aqueous phase was extracted once more with ethyl acetate. The combined organic phases were dried (MgSO₄), filtered and concentrated *in vacuo*. The residue was purified by flash chromatography eluting with toluene to give 3.8 g (79%) of 3,5-diisopropoxy-benzaldehyde as a yellow oil. ¹H NMR (CDCl₃, 300 MHz) δ: 1.35 (d, 12H), 4.60 (heptet, 2H), 6.68 (t, 1H), 6.97 (d, 2H).

b)

The title compound was prepared from 3,5-diisopropoxy-benzaldehyde by a sequence analogous to that described in example 23.

¹H NMR (CDCl₃, 300 MHz) δ: 1.15 (t, 3H), 1.22 (t, 3H), 1.32 (d, 12H), 2.96 (d, 2H), 3.32-3.42 (m, 1H), 3.55-3.65 (m, 1H), 3.98 (t, 1H), 4.16 (q, 2H), 4.53 (heptet, 2H), 4.65 (dd, 2H), 6.30-6.40 (m, 2H), 6.52 (d, 2H), 6.62 (d, 1H), 6.88 (d, 2H), 7.15 (d, 2H).

EXAMPLE 76




(E)-(S)-3-{4-[3-(3,5-Diisopropoxy-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid

The title compound was prepared from (E)-(S)-3-{4-[3-(3,5-diisopropoxy-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester (example 75) (800 mg, 1.7 mmol) by a procedure analogous to that described in example 26.

¹H NMR (CDCl₃, 300 MHz) δ: 1.18 (t, 3H), 1.32 (d, 12H), 2.95 (dd, 1H), 3.10 (dd, 1H), 3.40-3.52 (m, 1H), 3.55-3.65 (m, 1H), 4.05 (dd, 1H), 4.53 (heptet, 2H), 6.30-6.40 (m, 2H), 6.52 (d, 2H), 6.62 (d, 1H), 6.88 (d, 2H), 7.15 (d, 2H).

5

EXAMPLE 77

10 (S)-3-{4-[3-(3-Bromo-5-styryl-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester


a)

A mixture of styrene (2.0 g, 18.9 mmol), potassium carbonate (7.8 g, 56.7 mmol), tetra-N-butylammonium bromide (2.0 g, 6.3 mmol) and palladium(II) acetate (250 mg, 1.11 mmol),
15 under nitrogen, was stirred for 10 min. A solution of 3,5-dibromobenzaldehyde (5.0 g, 18.9 mmol) in dry DMF (10 mL) was added and the mixture heated at 65°C for 16 h. The reaction mixture was concentrated in vacuo, and the product purified by flash chromatography (heptane/ethyl acetate 1:4) to give 1.7 g of 3-bromo-5-styryl-benzaldehyde.

20 b)

The title compound was prepared from 3-bromo-5-styryl-benzaldehyde by a sequence analogous to that described in example 23.

¹H NMR (CDCl₃, 300 MHz) δ: 1.18 (t, 3H), 1.23 (t, 3H), 2.96 (d, 2H), 3.30-3.42 (m, 1H), 3.55-3.65 (m, 1H), 3.98 (t, 1H), 4.15 (q, 2H), 4.68 (d, 2H), 6.43 (dt, 1H), 6.66 (d, 1H), 6.88 (d, 2H),
25 6.93-7.56 (m, 12H).


EXAMPLE 78

5 (S)-3-{4-[3-(3-bromo-5-styryl-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid

The title compound was prepared from (S)-3-{4-[3-(3-bromo-5-styryl-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester (example 77) (800 mg, 1.7 mmol) by a procedure analogous to that described in example 26.

- 10 ¹H NMR (CDCl₃, 300 MHz) δ: 1.18 (t, 3H), 2.98 (dd, 1H), 3.10 (dd, 1H), 3.40-3.53 (m, 1H), 3.54-3.68 (m, 1H), 4.05 (dd, 1H), 4.68 (dd, 2H), 6.43 (dt, 1H), 6.68 (s, 1H), 6.88 (d, 2H), 6.94-7.56 (m, 12H).


15

EXAMPLE 79

20 (E)-3-{4-[3-(3-phenyl-allyloxy)-phenyl]-2-ethoxy-propionic acid ethyl ester

The title compound was prepared from 3-phenyl-prop-2-en-1-ol (270 mg, 2.0 mmol) by a sequence analogous to that described in example 23c.


- 25 ¹H NMR (CDCl₃, 300 MHz) δ: 1.18 (t, 3H), 1.22 (t, 3H), 2.95 (d, 2H), 3.30-3.42 (m, 1H), 3.53-3.65 (m, 1H), 3.98 (t, 1H), 4.15 (q, 2H), 4.68 (dd, 2H), 6.41 (dt, 1H), 6.73 (dt, 1H), 6.88 (d, 2H), 7.15 (d, 2H), 7.21-7.38 (m, 3H), 7.38-7.43 (m, 2H).

EXAMPLE 80

(E)-(S)-2-Ethoxy-3-[4-(3-phenyl-allyloxy)-phenyl]-propionic acid

The title compound was prepared from (E)-(S)-2-ethoxy-3-[4-(3-phenyl-allyloxy)-phenyl]-propionic acid ethyl ester (example 79) (700 mg, 2.0 mmol) by a procedure analogous to that described in example 26.

¹H NMR (CDCl₃, 300 MHz) δ: 1.18 (t, 3H), 2.95 (dd, 1H), 3.10 (dd, 1H), 4.42-3.53 (m, 1H), 3.53-3.64 (m, 1H), 4.05 (dd, 1H), 4.68 (dd, 2H), 6.42 (dt, 1H), 6.72 (d, 1H), 6.89 (d, 2H), 7.15 (d, 1H), 7.22-7.37 (m, 3H), 7.40 (d, 2H).

EXAMPLE 81

(E)-(S)-3-[4-[3-(2',3'-Dichloro-biphenyl-4-yl)-allyloxy]-phenyl]-2-ethoxy-propionic acid ethyl ester

a)

(E)-3-(4-Bromo-phenyl)-acrylic acid ethyl ester was prepared from 3-bromobenzaldehyde (20.0 g, 0.11 mol) by a sequence analogous to that described in example 23a.


b)

The title compound was prepared from (E)-3-(4-bromo-phenyl)-acrylic acid ethyl ester and 2,3-dichlorobenzene boronic acid by a sequence analogous to that described in example 52a-c.

- 5 ¹H NMR (CDCl₃, 300 MHz) δ: 1.18 (t, 3H), 1.23 (t, 3H), 2.95 (d, 2H), 3.30-3.42 (m, 1H), 3.53-3.65 (m, 1H), 3.98 (t, 1H), 4.15 (q, 2H), 4.70 (dd, 2H), 6.47 (dt, 1H), 6.7 (d, 1H), 6.88 (d, 2H), 7.15 (d, 2H), 7.20-7.28 (m, 2H), 7.35 (d, 2H), 7.43-7.52 (m, 3H).

10

EXAMPLE 82




(E)-(S)-3-{4-[3-(2',3'-Dichloro-biphenyl-4-yl)-allyloxy]-phenyl}-2-ethoxy-propionic acid

15

The title compound was prepared from (E)-(S)-3-{4-[3-(2',3'-dichloro-biphenyl-4-yl)-allyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester (example 81) by a procedure analogous to that described in example 26.

- 20 ¹H NMR (MeOD, 300 MHz) δ: 1.12 (t, 3H), 2.88 (dd, 1H), 3.0 (dd, 1H), 3.30-3.42 (m, 1H), 3.3-3.65 (m, 1H), 4.0 (dd, 1H), 4.70 (dd, 2H), 6.52 (dt, 1H), 6.80 (d, 1H), 6.90 (d, 2H), 7.18 (d, 2H), 7.25-7.40 (m, 4H), 7.48-7.55 (m, 3H).

EXAMPLE 83

- 5 (E)-(S)-3-(4-[3-(3,5-Bis-phenylethynyl-phenyl)-allyloxy]-phenyl)-2-ethoxy-propionic acid ethyl ester.
-


a)

A mixture of potassium carbonate (2.1 g, 15.2 mmol), tetra-N-butylammonium bromide (0.75
10 g, 2.4 mmol) and palladium(II) acetate (75 mg, 0.33 mmol) in dry DMF (8 ML), under nitro-
gen, was stirred for 10 min. 3-(3,5-Dibromophenyl)-acrylic acid ethyl ester (1.2 g, 3.6 mmol)
was added and the mixture cooled on ice. Phenylacetylene (4.0 mL, 36.0 mmol) was added
and the mixture stirred at room temperature for 7 days. The reaction mixture was added wa-
ter and the product extracted with ethyl acetate (x 3). The combined organic phases were
15 dried and concentrated in vacuo to give crude 3-(3,5-bis-phenylethynyl-phenyl)-acrylic acid
ethyl ester.

b)

The title compound was prepared from 3-(3,5-bis-phenylethynyl-phenyl)-acrylic acid ethyl
20 ester by a sequence analogous to that described in example 23b-c:

¹H NMR (CDCl₃, 300 MHz) δ: 1.18 (t, 3H), 1.23 (t, 3H), 2.96 (d, 2H), 3.30-3.42 (m, 1H), 3.55-
3.67 (m, 1H), 3.98 (t, 1H), 4.15 (q, 2H), 4.70 (d, 2H), 6.46 (dt, 1H), 6.68 (d, 1H), 6.88 (d, 2H),
7.15 (d, 2H), 7.30-7.38 (m, 6H), 7.48-7.58 (m, 6H), 7.60 (s, 1H).


EXAMPLE 84

5 (E)-(S)-3-{4-[3-(3,5-Bis-phenylethynyl-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid

The title compound was prepared from (E)-(S)-3-{4-[3-(3,5-bis-phenylethynyl-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester (example 83) (130 mg, 0.24 mmol) by a procedure analogous to that described in example 26.

10 ¹H NMR (CDCl₃, 300 MHz) δ: 1.18 (t, 3H), 2.96 (dd, 1H), 3.98 (dd, 1H), 3.37-3.48 (m, 1H), 3.53-3.67 (m, 1H), 4.03 (dd, 1H), 4.68 (d, 2H), 6.47 (dt, 1H), 6.68 (d, 1H), 6.88 (d, 2H), 7.18 (d, 2H), 7.30-7.42 (m, 6H), 7.48-7.58 (m, 6H), 7.60 (s, 1H).

15

EXAMPLE 85

(E)-(S)-3-{4-[3-(3,5-Diphenethyl-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester

a)

A solution of 3,5-distyryl-benzaldehyde (2.0 g, 6.44 mmol) (prepared as described in example 79) in ethyl acetate (150 mL) was hydrogenated at 3 atm for 16 h using 5% Pd-C (2 g) as catalyst. The catalyst was removed by filtration and the solvent evaporated to give (3,5-diphenethyl-phenyl)-methanol (2.0 g) as an oil.


b)

To a solution of (3,5-diphenethyl-phenyl)-methanol (2.0 g, 6.4 mmol) in dry dichloromethane (30 mL) was added pyridinium chlorochromate (1.4 g, 6.4 mmol) and the mixture was stirred at room temperature for 16 h. The product was purified by flash chromatography using dichloromethane as solvent to give 1.3 g 3,5-diphenethyl-benzaldehyde.

c)

The title compound was prepared from 3,5-diphenethyl-benzaldehyde by a sequence analogous to that described in example 23.

EXAMPLE 86




(E)-(S)-3-{4-[3-(3,5-Diphenethyl-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid

The title compound was prepared from (E)-(S)-3-{4-[3-(3,5-Diphenethyl-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester (example 85) (449 mg, 0.80 mmol) by a procedure analogous to that described in example 26.

¹H NMR (CDCl₃, 300 MHz) δ: 1.18 (t, 3H), 2.96 (dd, 1H), 3.08 (dd, 1H), 3.35-3.48 (m, 1H), 3.55-3.67 (m, 1H), 4.03 (dd, 1H), 4.65 (dd, 1H), 6.35 (dt, 1H), 6.68 (d, 1H), 6.82-6.92 (m, 3H), 7.04 (d, 2H), 7.12-7.32 (m, 12H).

5


EXAMPLE 87

10 3-{4-[3-(3,5-Bis-cyclopentyloxy-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester

The title compound was prepared from dihydroxybenzaldehyde (1.0 g, 7.2 mmol) and cyclopentylbromide (4.0 g, 29.0 mmol) by a sequence analogous to that described in example 75.

15 ¹H NMR (CDCl₃, 300 MHz) δ: 1.18 (t, 3H), 1.21 (t, 3H), 1.50-1.68 (m, 4H), 1.68-1.97 (m, 12H), 2.95 (d, 2H), 3.28-3.42 (m, 1H), 3.54-3.65 (m, 1H), 3.97 (t, 1H), 4.15 (q, 2H), 4.67 (dd, 2H), 4.67-4.77 (m, 2H), 6.28-6.40 (m, 2H), 6.50 (d, 2H), 6.60 (d, 1H), 6.87 (d, 2H), 7.15 (d, 2H).

20


EXAMPLE 88

5 (E)-(S)-3-{4-[3-(3,5-Bis-cyclopentyloxy-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid

The title compound was prepared from (E)-(S)-3-{4-[3-(3,5-bis-cyclopentyloxy-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester (example 87) (220 mg, 0.42 mmol) by a procedure analogous to that described in example 26.

10 ¹H NMR (CDCl₃, 300 MHz) δ: 1.18 (t, 3H), 1.52-1.70 (m, 4H), 1.70-1.98 (m, 12H), 2.95 (dd, 1H), 3.07 (dd, 1H), 3.37-3.48 (m, 1H), 3.55-3.65 (m, 1H), 4.03 (dd, 1H), 4.65 (dd, 2H), 4.70-4.78 (m, 2H), 6.29-6.40 (m, 2H), 6.50 (d, 2H), 6.60 (d, 1H), 6.88 (d, 2H), 7.15 (d, 2H).

15

EXAMPLE 89

20 (E)-(S)-3-{4-[3-[3,5-Bis-(2,2,2-trifluoro-ethoxy)-phenyl]-allyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester

a)


To a solution of 3,5-dihydroxybenzaldehyde (2.0 g, 14.5 mmol) in DMF (35 mL) was added potassium carbonate (11.0 g, 80.0 mmol) and 1,1,1-trifluoro-2-iodoethane (33.3 g, 160 mmol). The reaction mixture was heated in a sealed reactor at 50°C for 7 days. The mixture was filtered and washed with ethyl acetate. The filtrate was added water and the organic phase isolated. The aqueous phase was extracted once more with ethyl acetate. The combined organic phases were dried (MgSO₄), filtered and concentrated *in vacuo*. The residue was purified by flash chromatography eluting with toluene to give 906 mg (18%) of 3,5-bis-(2,2,2-trifluoro-ethoxy)-benzaldehyde. ¹H NMR (CDCl₃, 300 MHz) δ: 4.43 (q, 4H), 6.85 (t, 1H), 7.15 (d, 2H), 9.95 (s, 1H).

b)

The title compound was prepared from 3,5-bis-(2,2,2-trifluoro-ethoxy)-benzaldehyde by a sequence analogous to that described in example 23.

¹H NMR (CDCl₃, 300 MHz) δ: 1.15 (t, 3H), 1.22 (t, 3H), 2.95 (d, 2H), 3.30-3.40 (m, 1H), 3.55-3.67 (m, 1H), 3.97 (t, 1H), 4.15 (q, 2H), 4.33 (q, 4H), 4.65 (d, 2H), 6.32-6.48 (m, 2H), 6.55-6.70 (m, 3H), 6.85 (d, 2H), 7.15 (d, 2H).

EXAMPLE 90




(E)-(S)-3-(4-{3-[3,5-Bis-(2,2,2-trifluoro-ethoxy)-phenyl]-allyloxy}-phenyl)-2-ethoxy-propionic acid

The title compound was prepared from (E)-(S)-3-(4-{3-[3,5-bis-(2,2,2-trifluoro-ethoxy)-phenyl]-allyloxy}-phenyl)-2-ethoxy-propionic acid ethyl ester (example 89) (200 mg, 0.36 mmol) by a procedure analogous to that described in example 26.

¹H NMR (CDCl₃, 300 MHz) δ: 1.20 (t, 3H), 2.97 (dd, 1H), 3.10 (dd, 1H), 3.41-3.53 (m, 1H),
5 3.55-3.68 (m, 1H), 4.05 (dd, 1H), 4.35 (q, 4H), 4.67 (d, 2H), 6.35-6.48 (m, 2H), 6.60-6.70 (m, 3H), 6.87 (d, 2H), 7.15 (d, 2H).

EXAMPLE 91



(E)-(S)-Ethyl 2-Ethoxy-3-{4-[3-(4-furan-2-yl-phenyl)-but-2-enyloxy]-phenyl}-propionate

15 a)

Sodium (5.52 g, 0.24 mol) was added to ethanol (250 ml) at 20°C and the mixture stirred until the metal had fully reacted. Triethyl phosphonoacetate (62.72 g, 0.28 mol) was added as an ethanol (50 ml) solution, the mixture stirred for 20 min, then a solution of 4-iodoacetophenone (49.21 g, 0.20 mol) in ethanol (300 ml) was added and the reaction mixture heated to 80°C under reflux for 17h. The solution was cooled, the ethanol evaporated and the resulting orange residue partitioned between 1N HCl (200 ml) and ethyl acetate (200 ml). The aqueous layer was collected and further extracted with ethyl acetate (3 x 200 ml). The organic layers were combined, washed with brine, dried (MgSO₄) and evaporated to an orange/yellow oil, which was purified by column chromatography on silica gel (2% diethyl ether in n-heptane eluent) to give the product, (E)-ethyl 3-(4-iodophenyl)-but-2-enoate, as a pale yellow oil; 54.83 g (87%)

¹H NMR (300 MHz, CDCl₃) δ: 1.31 (3H, t), 2.53 (3H, s), 4.21 (2H, q), 6.11 (1H, s), 7.20 (2H, dm), 7.69 (2H, dm). ¹³CNMR (75MHz, CDCl₃) δ: 13.0 (q), 16.4 (q), 58.6 (t), 93.7 (s), 116.2 (d), 126.7 (d), 136.3 (d), 140.3 (s), 152.8 (s), 165.2 (s). MS: 316 (M⁺), 287, 271, 244, 144,
30 115 (100%). Microanalysis Calculated % C: 45.59, H: 4.14. Found % C: 45.72, H: 4.20.

b)

Tetrakis(triphenylphosphine)palladium(0) (0.69 g, 0.60 mmol, 6 mol%) was added, under nitrogen, to a stirred solution of (*E*)-ethyl 3-(4-iodophenyl)-but-2-enoate (3.16 g, 10.0 mmol) in DME (100 ml), and the resulting orange coloured solution stirred at room temperature for 10 min. Aqueous 2M sodium carbonate (30.0 ml, 60.0 mmol) was then added, the mixture stirred for 10 min, then furan-2-boronic acid (2.25 g, 20.11 mmol) was added, and the reaction mixture heated to 80°C for 20 h, under reflux. The reaction mixture was cooled, diluted with water (100 ml) and the products extracted into ethyl acetate (3 x 100 ml). The combined organic extracts were washed with brine, dried (MgSO₄), and evaporated to give the crude product, which was purified by column chromatography on silica gel (3% diethyl ether in n-heptane eluent) to give the product, (*E*)-ethyl 3-(4-furan-2-yl-phenyl)-but-2-enoate as an off-white solid; 2.46 g (96%).

Mpt. 85.5-86.5°C. ¹H NMR (300 MHz, CDCl₃) δ: 1.32 (3H, t), 2.59 (3H, d), 4.22 (2H, q), 6.18 (1H, m), 6.49 (1H, dd), 6.70 (1H, d), 7.46-7.56 (3H, m), 7.66 (2H, dm). MS: 256 (100%, M⁺), 227, 211, 184, 153, 115. Microanalysis Calculated % C: 74.98, H: 6.29. Found % C: 74.99, H: 6.39.


c)

(*E*)-Ethyl 3-(4-furan-2-yl-phenyl)-but-2-enoate was reduced with DIBAL-H by a procedure analogous to that described in example 50b, to give the colourless solid (*E*)-3-(4-furan-2-yl-phenyl)-but-2-en-1-ol.

d)

The title compound (678 mg, 77%) was prepared from (*E*)-3-(4-furan-2-yl-phenyl)-but-2-en-1-ol (430 mg, 2.0 mmol) and (*S*)-ethyl 2-ethoxy-3-(4-hydroxyphenyl)-propionate (526 mg, 2.21 mmol) by a procedure analogous to that described in example 52c.

¹H NMR (300 MHz, CDCl₃) δ: 1.17 (3H, t), 1.22 (3H, t), 2.14 (3H, d), 2.96 (2H, d), 3.31-3.41 (1H, m), 3.55-3.66 (1H, m), 3.98 (1H, t), 4.16 (2H, q), 4.73 (2H, d), 6.10 (1H, tm), 6.47 (1H, dd), 6.64 (1H, d), 6.88 (2H, dm), 7.17 (2H, dm), 7.43-7.48 (3H, m), 7.62 (2H, dm). LCMS: 457 (M+Na), 452 (M+NH₄), 197 (100%).

EXAMPLE 925 (E)-(S)-Ethyl 2-Ethoxy-3-{4-[3-(2'-methyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionate

a)

The colourless solid, (*E*)-ethyl 3-(2'-methyl-biphenyl-4-yl)-but-2-enoate was prepared from (*E*)-ethyl 3-(4-iodophenyl)-but-2-enoate (example 91a) and *ortho*-tolyl boronic acid by a procedure analogous to that described in example 91b.

10

b)

The colourless oil (*E*)-3-(2'-methyl-biphenyl-4-yl)-but-2-en-1-ol was prepared by DIBAL-H reduction of (*E*)-ethyl 3-(2'-methyl-biphenyl-4-yl)-but-2-enoate by a procedure analogous to that described in example 50b.


15

c)

The title compound (1.80 g, 78%) was prepared as a colourless oil from (*E*)-3-(2'-methyl-biphenyl-4-yl)-but-2-en-1-ol (1.19 g, 4.99 mmol) and (*S*)-ethyl 2-ethoxy-3-(4-hydroxyphenyl)-propionate (1.31 g, 6.48 mmol) by a procedure analogous to that described in example 52c.

20


¹H NMR (300 MHz, CDCl₃) δ: 1.17 (3H, t), 1.22 (3H, t), 2.17 (3H, d), 2.28 (3H, s), 2.96 (2H, d), 3.30-3.41 (1H, m), 3.54-3.66 (1H, m), 3.98 (1H, t), 4.16 (2H, q), 4.74 (2H, d), 6.11 (1H, tm), 6.88 (2H, dm), 7.16 (2H, dm), 7.20-7.32 (6H, m), 7.47 (2H, dm). LCMS: 679 (M+221), 633 (679-EtOH), 481 (M+Na), 476 (M+NH₄), 221 (100%).

EXAMPLE 93

5 (E)-(S)-2-Ethoxy-3-{4-[3-(2'-methyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionic acid

The title compound was prepared by hydrolysis of (E)-(S)-ethyl 2-ethoxy-3-{4-[3-(2'-methyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionate (example 92) (918 mg, 2.0 mmol) with sodium hydroxide by a procedure analogous to that described in example 51, yielding (E)-(S)-2-ethoxy-3-{4-[3-(2'-methyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionic acid as a colourless gum, which contained 0.25 mol equivalents of ethyl acetate; 586 mg (64%).

¹H NMR (300 MHz, CDCl₃) δ: 1.17 (3H, t), 1.27 (0.75H, t, AcOEt), 2.04 (0.75H, s, AcOEt), 2.19 (3H, d), 2.29 (3H, s), 2.96 (1H, dd), 3.09 (1H, dd), 3.41-3.53 (1H, m), 3.53-3.65 (1H, m), 4.06 (1H, dd), 4.12 (0.5H, q, AcOEt), 4.75 (2H, d), 6.12 (1H, tm), 6.89 (2H, dm), 7.16 (2H, dm), 7.20-7.34 (6H, m), 7.48 (2H, dm), carboxylic acid proton not observed. LCMS: 651 (M+221), 453 (M+Na), 221 (100%).

EXAMPLE 94

(E)-(S)-Ethyl 3-{4-[3-(2',5'-Dimethoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate

a)

The colourless oil, (*E*)-ethyl 3-(2',5'-dimethoxy-biphenyl-4-yl)-but-2-enoate was prepared from (*E*)-ethyl 3-(4-bromophenyl)-but-2-enoate (example 50a) and 2,5-dimethoxyphenyl boronic acid by a procedure analogous to that described in example 52a.

5 b)

The colourless gum (*E*)-3-(2',5'-dimethoxy-biphenyl-4-yl)-but-2-en-1-ol was prepared by DIBAL-H reduction of (*E*)-ethyl 3-(2',5'-dimethoxy-biphenyl-4-yl)-but-2-enoate by a procedure analogous to that described in example 52b.


10 c)

The title compound (0.765 g, 61%) was prepared as a colourless gum from (*E*)-3-(2',5'-dimethoxy-biphenyl-4-yl)-but-2-en-1-ol (0.711 g, 2.50 mmol) and (*S*)-ethyl 2-ethoxy-3-(4-hydroxyphenyl)-propionate (0.655 g, 2.75 mmol) by a procedure analogous to that described in example 52c.

15 ¹H NMR (300 MHz, CDCl₃) δ: 1.17 (3H, t), 1.22 (3H, t), 2.16 (3H, d), 2.96 (2H, d), 3.31-3.41 (1H, m), 3.55-3.65 (1H, m), 3.76 (3H, s), 3.81 (3H, s), 3.98 (1H, t), 4.17 (2H, q), 4.74 (2H, d), 6.10 (1H, tm), 6.81-6.95 (5H, m), 7.16 (2H, dm), 7.45-7.53 (4H, m). LCMS: 771 (M+267), 527 (M+Na), 422 (M+NH₄), 267 (100%).

20

EXAMPLE 95




25 (*E*)-(S)-3-{4-[3-(2',5'-Dimethoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid

The title compound was prepared from (*E*)-(S)-ethyl 3-{4-[3-(2',5'-dimethoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate (Example 94) (0.62 g, 1.23 mmol) and sodium hydroxide (1M, 2.0 ml, 2.0 mmol) by a procedure analogous to that described in example 51,

yielding (*E*)-(*S*)-ethyl 3-{4-[3-(2',5'-dimethoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid as a colourless glass; 0.485 g (83%).

¹H NMR (300 MHz, CDCl₃) δ: 1.18 (3H, t), 2.16 (3H, d), 2.96 (1H, dd), 3.10 (1H, dd), 3.42-3.51 (1H, m), 3.51-3.65 (1H, m), 3.75 (3H, s), 3.80 (3H, s), 4.05 (1H, dd), 4.74 (2H, d), 6.10 (1H, tm), 6.81-6.94 (5H, m), 7.17 (2H, dm), 7.43-7.53 (4H, m), carboxylic acid proton not observed. LCMS: 743 (M+267), 499 (M+Na), 494 (M+NH₄), 267 (100%).

EXAMPLE 96



(*E*)-(*S*)-Ethyl 3-{4-[3-(4-Bromophenyl)-2-ethyl-but-2-enyloxy]-phenyl}-2-ethoxy-propionate

15 a)

A solution of triethyl 2-phosphonobutyrate (17.7 g, 70.0 mmol) in dry THF (30 ml) was added dropwise, at 0°C, to a stirred suspension of sodium hydride (50% dispersion in mineral oil, 2.90 g, 60.4 mmol) in dry THF (30 ml) and the mixture stirred for 30 min. A solution of 4-bromoacetophenone (7.96 g, 39.99 mmol) in THF (80 ml) was added over 20 min., the resulting mixture warmed to room temperature and stirring continued overnight. Second portions of triethyl 2-phosphonobutyrate (10.11 g, 40.1 mmol) and sodium hydride (2.90 g, 60.4 mmol) were then added at room temperature, and stirring continued for a further 24 h; TLC at this stage showed that a substantial amount of unreacted 4-bromo acetophenone starting material was still present. The reaction was worked up by adding 1N HCl (200 ml) and ethyl acetate (100ml), the organic layer collected and the aqueous layer extracted with ethyl acetate (3 x 100ml). The combined organic layers were washed with brine, dried (MgSO₄) and concentrated to give an orange gum, which was purified by column chromatography on silica gel (2% diethyl ether in n-heptane eluent) to give the orange oil, (*E/Z*)-ethyl 3-(4-bromophenyl)-2-ethyl-but-2-enoate (3.47 g, 29%) as a mixture of double-bond isomers.

30

b)


A toluene solution of DIBAL-H (1M, 29.0 ml, 29.0 mmol) was added dropwise at -70°C to a stirred THF (100 ml) solution of (*E/Z*)-ethyl 3-(4-bromophenyl)-2-ethyl-but-2-enoate (3.45 g, 11.6 mmol), and the solution stirred for 40 min. Methanol (1 ml) was carefully added, followed by 1N HCl (300 ml) and ethyl acetate (200 ml). The aqueous layer was separated and further extracted with ethyl acetate (2 x 150 ml). The combined organic layers were washed with brine, dried (MgSO_4), and concentrated to give an orange gum, which was separated into its two major constituents by column chromatography on silica gel (15% ethyl acetate in *n*-heptane eluent). The two products, in order of elution, were (*Z*)-3-(4-bromo-phenyl)-2-ethyl-but-2-en-1-ol (0.365 g, 12%) and (*E*)-3-(4-bromophenyl)-2-ethyl-but-2-en-1-ol (0.89 g, 30%).

c)

The title compound (843 mg, 89%) was prepared from (*E*)-3-(4-bromophenyl)-2-ethyl-but-2-en-1-ol (510 mg, 2.0 mmol) and (*S*)-ethyl 2-ethoxy-3-(4-hydroxyphenyl)-propionate (500 mg, 2.10 mmol) by a procedure analogous to that described in example 52c.

^1H NMR (300 MHz, CDCl_3) δ : 0.94 (3H, t), 1.20 (3H, t), 1.23 (3H, t), 2.01 (3H, s), 2.05 (2H, q), 2.97 (2H, d), 3.31-3.43 (1H, m), 3.54-3.68 (1H, m), 3.99 (1H, t), 4.17 (2H, q), 4.61 (2H, s), 6.89 (2H, dm), 7.04 (2H, dm), 7.18 (2H, dm), 7.45 (2H, dm).

EXAMPLE 97




(*E*)-(S)-3-{4-[3-(4-Bromophenyl)-2-ethyl-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid

The title compound was prepared from (*E*)-(S)-ethyl 3-{4-[3-(4-bromophenyl)-2-ethyl-but-2-enyloxy]-phenyl}-2-ethoxy-propionate (Example 96) (0.78 g, 1.64 mmol) and sodium hydroxide (1M, 3.3 ml, 3.3 mmol) by a procedure analogous to that described in example 51, yielding (*E*)-(S)-3-{4-[3-(4-bromophenyl)-2-ethyl-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid

(0.703 g, 96%) as a pale yellow oil, which contained a small amount of dichloromethane; 0.703 g (96%).

¹H NMR (300 MHz, CDCl₃) δ: 1.19 (3H, t), 1.26 (ethyl acetate impurity, 0.6H, t), 2.04 (ethyl acetate impurity, 0.4H, s), 2.16 (3H, s), 2.96 (1H, dd), 3.10 (1H, dd), 3.42-3.52 (1H, m), 3.53-3.68 (1H, m), 3.80 (3H, s), 4.07 (1H, dd), 4.12 (ethyl acetate impurity, 0.4H), 4.74 (2H, d), 5.30 (CH₂Cl₂, trace), 6.10 (1H, t), 6.85-6.95 (3H, m), 7.12-7.20 (2H, m), 7.21-7.32 (2H, m), 7.45-7.50 (4H, m), carboxylic acid proton not observed.

10

EXAMPLE 98


(Z)-(S)-Ethyl 3-{4-[3-(4-Bromophenyl)-2-ethyl-but-2-enyloxy]-phenyl}-2-ethoxy-propionate

15

The title compound (535 mg, 81%) was prepared from (Z)-3-(4-bromophenyl)-2-ethyl-but-2-en-1-ol (prepared as described in example 96b) (355 mg, 1.39 mmol) and (S)-ethyl 2-ethoxy-3-(4-hydroxyphenyl)-propionate (348 mg, 1.46 mmol) by a procedure analogous to that described in example 52c.

¹H NMR (300 MHz, CDCl₃) δ: 1.11 (3H, t), 1.16 (3H, t), 1.21 (3H, t), 2.04 (3H, s), 2.37 (2H, q), 2.93 (2H, d), 3.29-3.40 (1H, m), 3.53-3.65 (1H, m), 3.95 (1H, t), 4.16 (2H, q), 4.25 (2H, s), 6.70 (2H, dm), 7.03-7.12 (4H, m), 7.40 (2H, dm). Microanalysis Calculated % C: 63.16, H: 6.57. Found % C: 63.34, H: 6.66.

25


EXAMPLE 99

(Z)-(S)-3-{4-[3-(4-Bromophenyl)-2-ethyl-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid

The title compound was prepared from (Z)-(S)-ethyl 3-{4-[3-(4-bromophenyl)-2-ethyl-but-2-enyloxy]-phenyl}-2-ethoxy-propionate (Example 98) (475 mg, 1.0 mmol) and sodium hydroxide (1M, 2.0 ml, 2.0 mmol) by a procedure analogous to that described in example 51, yielding (Z)-(S)-3-{4-[3-(4-bromophenyl)-2-ethyl-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid (0.424 g, 95%) as a pale yellow oil.

¹H NMR (300 MHz, CDCl₃) δ: 1.11 (3H, t), 1.18 (3H, t), 2.04 (3H, s), 2.37 (2H, q), 2.94 (1H, dd), 3.04 (1H, dd), 3.40-3.53 (1H, m), 3.53-3.64 (1H, m), 4.03 (1H, dd), 4.25 (2H, s), 6.71 (2H, dm), 7.02-7.14 (4H, m), 7.40 (2H, dm), carboxylic acid proton not observed.

EXAMPLE 100



(E)-(S)-Ethyl 3-{4-[3-(4'-tert-Butyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate

a)

The colourless oil, (E)-ethyl 3-(4'-tert-butyl-biphenyl-4-yl)-but-2-enoate was prepared from (E)-ethyl 3-(4-bromophenyl)-but-2-enoate (example 50a) and 4-tert-butylphenylboronic acid by a procedure analogous to that described in example 52a

b)

The colourless gum (E)-3-(4'-tert-butyl-biphenyl-4-yl)-but-2-en-1-ol was prepared by DIBAL-H reduction of (E)-ethyl 3-(4'-tert-butyl-biphenyl-4-yl)-but-2-enoate by a procedure analogous to that described in example 52b.


c)

The title compound (0.375 g, 75%) was prepared as a colourless gum from (*E*)-3-(4'-*tert*-butyl-biphenyl-4-yl)-but-2-en-1-ol (0.280 g, 1.00 mmol) and (*S*)-ethyl 2-ethoxy-3-(4-hydroxyphenyl)-propionate (0.250 g, 1.05 mmol) by a procedure analogous to that described in example 52c.

- 5 ¹H NMR (300 MHz, CDCl₃) δ: 1.17 (3H, t), 1.22 (3H, t), 1.37 (9H, s), 2.17 (3H, d), 2.97 (2H, d), 3.30-3.43 (1H, m), 3.53-3.67 (1H, m), 3.98 (1H, t), 4.17 (2H, q), 4.73 (2H, d), 6.11 (1H, tm), 6.88 (2H, dm), 7.17 (2H, dm), 7.43-7.60 (8H, m). LCMS: 763 (M+263), 523 (M+Na), 263 (100%).

10

EXAMPLE 101



- 15 (*E*)-(*S*)-Ethyl 3-{4-[3-(3',5'-Bis-trifluoromethyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate
-

a)

- The colourless solid, (*E*)-ethyl 3-(3',5'-bis-trifluoromethyl-biphenyl-4-yl)-but-2-enoate was prepared from (*E*)-ethyl 3-(4-iodophenyl)-but-2-enoate (example 91a) and 3,5-bis-(trifluoromethyl)phenyl boronic acid by a procedure analogous to that described in example 91b.
- 20

b)

- 25 The colourless solid (*E*)-3-(3',5'-bis-trifluoromethyl-biphenyl-4-yl)-but-2-en-1-ol was prepared by DIBAL-H reduction of (*E*)-ethyl 3-(3',5'-bis-trifluoromethyl-biphenyl-4-yl)-but-2-enoate by a procedure analogous to that described in example 50b.


c)

The title compound (656 mg, 81%) was prepared as a colourless oil from (*E*)-3-(3',5'-bis-trifluoromethyl-biphenyl-4-yl)-but-2-en-1-ol (500 mg, 1.39 mmol) and (*S*)-ethyl 2-ethoxy-3-(4-hydroxyphenyl)-propionate (348 mg, 1.46 mmol) by a procedure analogous to that described in example 52c.

- 5 ¹H NMR (300 MHz, CDCl₃) δ: 1.17 (3H, t), 1.23 (3H, t), 2.18 (3H, d), 2.97 (2H, d), 3.30-3.43 (1H, m), 3.56-3.69 (1H, m), 3.98 (1H, t), 4.17 (2H, q), 4.75 (2H, d), 6.15 (1H, tm), 6.89 (2H, dm), 7.18 (2H, dm), 7.52-7.62 (4H, m), 7.85 (1H, s), 8.01 (2H, s). LCMS: 603 (100%, M+Na), 598 (M+NH₄), 343.

10


EXAMPLE 102



- 15 (*E*)-(S)-3-{4-[3-(3',5'-Bis-trifluoromethyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid
-

The title compound was prepared from (*E*)-(S)-ethyl 3-{4-[3-(3',5'-bis-trifluoromethyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate (Example 101) (625 mg, 1.08 mmol) and sodium hydroxide (1M, 4.3 ml, 4.3 mmol) by a procedure analogous to that described in example 51, yielding (*E*)-(S)-3-{4-[3-(3',5'-bis-trifluoromethyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid (535 mg, 90%) as a colourless gum.

- 20 ¹H NMR (300 MHz, CDCl₃) δ: 1.20 (3H, t), 2.18 (3H, d), 2.98 (1H, dd), 3.10 (1H, dd), 3.43-3.53 (1H, m), 3.53-3.66 (1H, m), 4.07 (1H, dd), 4.76 (2H, d), 6.15 (1H, tm), 6.90 (2H, dm),
25 7.19 (2H, dm), 7.50-7.62 (4H, m), 7.85 (1H, s), 8.01 (2H, s), carboxylic acid proton not observed.

EXAMPLE 103

5 (E)-(S)-Ethyl 2-Ethoxy-3-{4-[3-(4'-isopropyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionate

a)

The colourless solid, (*E*)-ethyl 3-(4'-isopropyl-biphenyl-4-yl)-but-2-enoate was prepared from (*E*)-ethyl 3-(4-bromophenyl)-but-2-enoate (example 50a) and 4-isopropylphenyl boronic acid
10 by a procedure analogous to that described in example 52a.

Mpt. 96.5-97.5°C. ¹H NMR (300 MHz, CDCl₃) δ: 1.29 (6H, d), 1.32 (3H, t), 2.61 (3H, d), 2.97 (1H, septet), 4.22 (2H, q), 6.20 (1H, m), 7.32 (2H, dm), 7.50-7.65 (6H, m). MS: 308 (100%, M⁺), 293, 178. Microanalysis Calculated % C: 81.78, H: 7.84. Found % C: 81.96, H: 8.22.

15 b)

The colourless solid (*E*)-3-(4'-isopropyl-biphenyl-4-yl)-but-2-en-1-ol was prepared by DIBAL-H reduction of (*E*)-ethyl 3-(4'-isopropyl-biphenyl-4-yl)-but-2-enoate by a procedure analogous to that described in example 50b.

Mpt. 110.5-112.5°C. ¹H NMR (300 MHz, CDCl₃) δ: 1.29 (6H, d), 2.10 (3H, s), 2.94 (1H, septet), 4.37 (2H, d), 6.03 (1H, t), 7.29 (2H, dm), 7.40-7.60 (6H, m). MS: 266 (M⁺), 251 (M-Me), 223 (100%, M-*i*-Pr). Microanalysis Calculated % C: 85.67, H: 8.32. Found % C: 85.55, H: 8.55.


c)

25 The title compound (410 mg, 84%) was prepared as a colourless solid from (*E*)-3-(4'-isopropyl-biphenyl-4-yl)-but-2-en-1-ol (266 mg, 1.00 mmol) and (*S*)-ethyl 2-ethoxy-3-(4-hydroxyphenyl)-propionate (250 mg, 1.05 mmol) by a procedure analogous to that described in example 52c.

30 Mpt. 70-73°C. ¹H NMR (300 MHz, CDCl₃) δ: 1.17 (3H, t), 1.22 (3H, t), 1.29 (6H, d), 2.17 (3H, d), 2.89-3.01 (3H, m), 3.30-3.41 (1H, m), 3.55-3.66 (1H, m), 3.98 (1H, t), 4.17 (2H, q), 4.74

(2H, d), 6.11 (1H, tm), 6.88 (2H, dm), 7.17 (2H, dm), 7.30 (2H, dm), 7.46-7.57 (6H, m).
LCMS: 735 (M+249), 509 (M+Na), 249 (100%).

5

EXAMPLE 104

(E)-(S)-2-Ethoxy-3-{4-[3-(4'-isopropyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionic acid


10

The title compound was prepared from (*E*)-(*S*)-ethyl 2-ethoxy-3-{4-[3-(4'-isopropyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionate (Example 103) (400 mg, 0.822 mmol) and sodium hydroxide (1M, 3.29 ml, 3.29 mmol) by a procedure analogous to that described in example 51, yielding (*E*)-(*S*)-2-ethoxy-3-{4-[3-(4'-isopropyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionic acid (380 mg, 100%) as a beige coloured solid.

15

¹H NMR (300 MHz, CDCl₃) δ: 1.19 (3H, t), 1.29 (6H, d), 2.17 (3H, d), 2.89-3.01 (2H, m), 3.10 (1H, dd), 3.42-3.64 (2H, m), 4.06 (1H, dd), 4.74 (2H, d), 6.11 (1H, tm), 6.90 (2H, dm), 7.16 (2H, dm), 7.30 (2H, dm), 7.46-7.57 (6H, m), carboxylic acid proton not observed. LCMS: 707 (M+249), 481 (M+Na), 249 (100%).


20

EXAMPLE 105

5 *(E)-(S)-3-{4-[3-(3,5-Dimethoxy-phenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester*

The title compound was prepared from 3',5'-dimethoxyacetophenone (7.0 g, 0.0388 mol) by a sequence analogous to that described in example 3, yielding 0.165 g (35%) of *(E)-(S)-3-{4-[3-(3,5-Dimethoxy-phenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester*.

¹H NMR (200 MHz, CDCl₃) δ: 1.1-1.27 (6H, m), 2.97 (2H, d), 3.3-3.4 (1H, m), 3.52-3.7 (1H, m), 4.0 (1H, t), 4.15 (2H, q), 4.7 (2H, d),), 6.39 (1H, dd), 6.57 (2H, dd), 6.88 (2H, d), 7.17 (2H, d).


EXAMPLE 106

20 *(E)-(S)-3-{4-[3-(3'-Acetyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid*

The title compound was prepared from (*E*)-(*S*)-ethyl 3-{4-[3-(3'-acetyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate (Example 105) (140 mg, 0.288 mmol) and sodium hydroxide (1M, 0.58 ml, 0.58 mmol) by a procedure analogous to that described in example 51, yielding (*E*)-(*S*)-3-{4-[3-(3'-acetyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid (33 mg, 25%) as a yellow coloured solid.

¹H NMR (300 MHz, CDCl₃) δ: 1.19 (3H, t), 2.18 (3H, d), 2.66 (3H, s), 2.97 (1H, dd), 3.10 (1H, dd), 3.41-3.65 (2H, m), 4.07 (1H, dd), 4.75 (2H, d), 6.13 (1H, tm), 6.90 (2H, dm), 7.17 (2H, dm), 7.49-7.63 (5H, m), 7.80 (1H, dm), 7.93 (1H, dm), 8.19 (1H, m), carboxylic acid proton not observed. LCMS: 707 (M+249), 481 (M+Na), 249 (100%).

EXAMPLE 107



(*E*)-(*S*)-Ethyl 3-{4-[3-(4'-Acetyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate

a)

(*E*)-3-(4-iodophenyl)-but-2-en-1-ol was prepared by DIBAL-H reduction of (*E*)-ethyl 3-(4-iodophenyl)-but-2-enoate (example 91a) by a procedure analogous to that described in example 50b.

¹H NMR (300 MHz, CDCl₃) δ: 1.36 (1H, br s), 2.04 (3H, d), 2.66 (3H, s), 4.36 (2H, br d), 5.96 (1H, tm), 7.15 (2H, dm), 7.65 (2H, dm).

b)

The pale yellow solid, (*E*)-1-[4'-(3-hydroxy-1-methyl-propenyl)-biphenyl-4-yl]-ethanone was prepared from 4-acetylphenyl boronic acid and (*E*)-3-(4-iodophenyl)-but-2-en-1-ol by a procedure analogous to that described in example 54a.


¹H NMR (300 MHz, CDCl₃) δ: 1.55 (1H, br s), 2.12 (3H, d), 2.64 (3H, s), 4.41 (2H, d), 6.07 (1H, tm), 7.52 (2H, dm), 7.61 (2H, dm), 7.70 (2H, dm), 8.03 (2H, dm).

c)

The title compound (275 mg, 75%) was prepared from (*E*)-1-[4'-(3-hydroxy-1-methyl-propenyl)-biphenyl-4-yl]-ethanone (200 mg, 0.75 mmol) and (*S*)-ethyl 2-ethoxy-3-(4-hydroxyphenyl)-propionate (188 mg, 0.79 mmol) by a procedure analogous to that described in example 52c.

¹H NMR (300 MHz, CDCl₃) δ: 1.17 (3H, t), 1.23 (3H, t), 2.18 (3H, d), 2.64 (3H, s), 2.97 (2H, d), 3.30-3.42 (1H, m), 3.55-3.67 (1H, m), 3.98 (1H, t), 4.17 (2H, q), 4.75 (2H, d), 6.14 (1H, tm), 6.89 (2H, dm), 7.17 (2H, dm), 7.54 (2H, dm), 7.61 (2H, dm), 7.70 (2H, dm), 8.03 (2H, dm).


EXAMPLE 108



(*E*)-(S)-3-{4-[3-(4'-Acetyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid

The title compound was prepared from (*E*)-(S)-ethyl 3-{4-[3-(4'-acetyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate (Example 107) (200 mg, 0.411 mmol) and sodium hydroxide (1M, 1.64 ml, 1.64 mmol) by a procedure analogous to that described in example 51, yielding (*E*)-(S)-3-{4-[3-(4'-acetyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid (70 mg, 37%) as a yellow coloured solid.

¹H NMR (300 MHz, CDCl₃) δ: 1.19 (3H, t), 2.18 (3H, d), 2.64 (3H, s), 2.97 (1H, dd), 3.11 (1H, dd), 3.45-3.65 (2H, m), 4.08 (1H, dd), 4.75 (2H, d), 6.14 (1H, tm), 6.90 (2H, dm), 7.17 (2H, dm), 7.54 (2H, dm), 7.61 (2H, dm), 7.70 (2H, dm), 8.03 (2H, dm), carboxylic acid proton not observed. LCMS: 707 (M+249), 459 (M+H), 249 (100%).

EXAMPLE 109

5 (E)-(S)-Ethyl 2-Ethoxy-3-[4-(3-[1,1';3',1'']terphenyl-5'-yl-allyloxy)-phenyl]-propionate

a)

Sodium (0.90 g, 39.1 mmol) was added to ethanol (50 ml) at 20°C and the mixture stirred until the metal had fully reacted. Triethyl phosphonoacetate (10.1 g, 45 mmol) was added,
 10 the mixture stirred for 10 min, then a solution of 3,5-dibromobenzaldehyde (7.92 g, 30 mmol) in ethanol (50 ml) was added and the reaction mixture heated to 80°C under reflux for 72h. The solution was cooled, the ethanol evaporated and the resulting yellow residue partitioned between 1N HCl (100 ml) and ethyl acetate (100 ml). The aqueous layer was collected and further extracted with ethyl acetate (3 x 100 ml). The organic layers were combined, washed
 15 with brine, dried (MgSO₄) and evaporated to a yellow solid, which was purified by column chromatography on silica gel (2% diethyl ether in n-heptane eluent) to give the product, (E)-ethyl 3-(3,5-dibromophenyl)-acrylate, as a colourless solid; 4.54 g (45%).
 Mpt. 80-82°C. ¹H NMR (300 MHz, CDCl₃) δ: 1.33 (3H, t), 4.27 (2H, q), 6.42 (1H, d), 7.51 (1H, d), 7.58 (2H, d), 7.66 (1H, t). MS: 336/334/332 (M⁺), 308/306/304, 291/289/287 (100%, M-Oet), 180/182.

b)

The colourless solid, (E)-ethyl 3-[1,1';3',1'']terphenyl-5'-yl-acrylate was prepared from (E)-ethyl 3-(3,5-dibromophenyl)-acrylate and phenylboronic acid by a procedure analogous to
 25 that described in example 52a.

Mpt. 78.5-81.5°C. ¹H NMR (300 MHz, CDCl₃) δ: 1.36 (3H, t), 4.29 (2H, q), 6.58 (1H, d), 7.33-7.54 (6H, m), 7.60-7.68 (4H, m), 7.72 (2H, d), 7.81 (1H, t), 7.82 (1H, d). MS: 328 (100%, M⁺), 283, 256, 252, 241, 239.

c)


The colourless solid (*E*)-3-[1,1';3',1'']terphenyl-5'-yl-prop-2-en-1-ol was prepared by DIBAL-H reduction of (*E*)-ethyl 3-[1,1';3',1'']terphenyl-5'-yl-acrylate by a procedure analogous to that described for example 52b.

- 5 Mpt. 140-141.5°C. ¹H NMR (300 MHz, CDCl₃) δ: 1.54 (1H, br s), 4.38 (2H, d), 6.50 (1H, dt), 6.75 (1H, d), 7.30-7.52 (6H, m), 7.53-7.73 (7H, m). MS: 286 (M⁺), 258 (100%), 243, 230, 165, 91, 77.

d)

- 10 The title compound (426 mg, 80%) was prepared from (*E*)-3-[1,1';3',1'']terphenyl-5'-yl-prop-2-en-1-ol (300 mg, 1.05 mmol) and (*S*)-ethyl 2-ethoxy-3-(4-hydroxyphenyl)-propionate (262 mg, 1.10 mmol) by a procedure analogous to that described in example 52c.
- ¹H NMR (300 MHz, CDCl₃) δ: 1.16 (3H, t), 1.22 (3H, t), 2.96 (2H, d), 3.29-3.41 (1H, m), 3.54-3.66 (1H, m), 3.98 (1H, t), 4.16 (2H, q), 4.73 (2H, d), 6.54 (1H, dt), 6.85 (1H, d), 6.90 (2H, dm), 7.17 (2H, dm), 7.30-7.50 (6H, m), 7.55-7.71 (7H, m). LCMS: 775 (M+269), 729 (100%, M+269-EtOH), 461 (M+H-EtOH), 269.
- 15

EXAMPLE 110




(*E*)-(S)-2-Ethoxy-3-[4-(3-[1,1';3',1'']terphenyl-5'-yl-allyloxy)-phenyl]-propionic acid

- 25 The title compound was prepared from (*E*)-(S)-ethyl 2-ethoxy-3-[4-(3-[1,1';3',1'']terphenyl-5'-yl-allyloxy)-phenyl]-propionate (Example 109) (405 mg, 0.8 mmol) and sodium hydroxide (1M, 1.6 ml, 1.6 mmol) by a procedure analogous to that described in example 51, yielding (*E*)-(S)-2-ethoxy-3-[4-(3-[1,1';3',1'']terphenyl-5'-yl-allyloxy)-phenyl]-propionic acid (352 mg, 92%) as a colourless glass.

^1H NMR (300 MHz, CDCl_3) δ : 1.18 (3H, t), 2.97 (1H, dd), 3.09 (1H, dd), 3.41-3.53 (1H, m), 3.53-3.65 (1H, m), 4.07 (1H, dd), 4.73 (2H, dd), 6.54 (1H, dt), 6.85 (1H, dm), 6.92 (2H, dm), 7.17 (2H, dm), 7.32-7.50 (6H, m), 7.55-7.71 (7H, m), carboxylic acid proton not observed. LCMS: 747 ($\text{M}+269$), 501 ($\text{M}+\text{Na}$), 496 ($\text{M}+\text{NH}_4$), 269 (100%).

5

EXAMPLE 111

10

(*E,E*)-(S)-Ethyl 3-(4'-{3-[4-(2-Ethoxy-2-ethoxycarbonyl-ethyl)-phenoxy]-1-methyl-propenyl}-biphenyl-3-yl)-but-2-enoate

a)

- 15 The yellow oil (*E,E*)-ethyl 3-[4'-(3-hydroxy-1-methyl-propenyl)-biphenyl-3-yl]-but-2-enoate was prepared from (*E*)-1-[4'-(3-hydroxy-1-methyl-propenyl)-biphenyl-3-yl]-ethanone (example 105a) and triethyl phosphonoacetate by a reaction analogous to that described for example 50a.

^1H NMR (300 MHz, CDCl_3) δ : 1.33 (3H, t), 1.37 (1H, br t), 2.13 (3H, d), 2.62 (3H, d), 4.23 (2H, q), 4.41 (2H, br t), 6.06 (1H, tm), 6.19 (1H, m), 7.40-7.62 (7H, m), 7.68 (1H, m). MS: 336 (M^+), 334, 308, 293, 43 (100%).

20

b)


- The title compound (230 mg, 58%) was prepared from (*E,E*)-ethyl 3-[4'-(3-hydroxy-1-methyl-propenyl)-biphenyl-3-yl]-but-2-enoate (235 mg, 0.70 mmol) and (*S*)-ethyl 2-ethoxy-3-(4-hydroxyphenyl)-propionate (175 mg, 0.73 mmol) by a procedure analogous to that described in example 52c.

25

^1H NMR (300 MHz, CDCl_3) δ : 1.17 (3H, t), 1.23 (3H, t), 1.33 (3H, t), 2.17 (3H, d), 2.62 (3H, d), 2.96 (2H, d), 3.30-3.42 (1H, m), 3.55-3.67 (1H, m), 3.98 (1H, t), 4.17 (2H, q), 4.23 (2H, q),

4.75 (2H, d), 6.13 (1H, tm), 6.20 (1H, m), 6.89 (2H, dm), 7.17 (2H, dm), 7.42-7.62 (7H, m), 7.67 (1H, m). MS: 556 (M^+), 319 (100%).

5


EXAMPLE 112

10 *(E,E)*-(*S*)- 3-(4'-(3-[4-(2-Carboxy-2-ethoxy-ethyl)-phenoxy]-1-methyl-propenyl)-biphenyl-3-yl)-but-2-enoic acid

The title compound was prepared from *(E,E)*-(*S*)-ethyl 3-(4'-(3-[4-(2-ethoxy-2-ethoxycarbonyl-ethyl)-phenoxy]-1-methyl-propenyl)-biphenyl-3-yl)-but-2-enoate (example 111) (190 mg, 0.34 mmol) and sodium hydroxide (1M, 1.4 ml, 1.4 mmol) by a procedure analogous to that described in example 51, yielding *(E,E)*-(*S*)- 3-(4'-(3-[4-(2-carboxy-2-ethoxy-ethyl)-phenoxy]-1-methyl-propenyl)-biphenyl-3-yl)-but-2-enoic acid (135 mg, 79%) as a colourless solid.

^1H NMR (300 MHz, CDCl_3) δ : 1.19 (3H, t), 2.18 (3H, m), 2.64 (3H, s), 3.01 (1H, dd), 3.08 (1H, dd), 3.40-3.70 (2H, m), 4.07 (1H, dd), 4.75 (2H, dd), 6.12 (1H, br m), 6.23 (1H, s), 6.89 (2H, dm), 7.18 (2H, dm), 7.40-7.70 (8H, m), carboxylic acid protons not observed.

25

EXAMPLE 113

(E)-(S)-Ethyl 2-Ethoxy-3-{4-[3-(3'-methoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionate

a)

The colourless oil (*E*)-ethyl 3-(3'-methoxy-biphenyl-4-yl)-but-2-enoate was prepared from (*E*)-ethyl 3-(4-bromophenyl)-but-2-enoate (example 50a) and 3-methoxyphenyl boronic acid by a procedure analogous to that described in example 52a.

¹H NMR (300 MHz, CDCl₃) δ: 1.33 (3H, t), 2.61 (3H, d), 3.87 (3H, s), 4.23 (2H, q), 6.20 (1H, m), 6.91 (1H, ddd), 7.13 (1H, dd), 7.19 (1H, ddd), 7.37 (1H, dd), 7.51-7.62 (4H, m). MS: 296 (100%, M⁺), 281, 267, 251, 224. Microanalysis Calculated % C: 77.00, H: 6.80. Found % C: 77.02, H: 6.93.

b)


The colourless solid (*E*)-3-(3'-methoxy-biphenyl-4-yl)-but-2-en-1-ol was prepared by DIBAL-H reduction of (*E*)-ethyl 3-(3'-methoxy-biphenyl-4-yl)-but-2-enoate as described for example 52b.

Mpt. 62-68°C. ¹H NMR (300 MHz, CDCl₃) δ: 1.40 (1H, br s), 2.12 (3H, d), 3.87 (3H, s), 4.39 (2H, d), 6.05 (1H, tm), 6.89 (1H, ddd), 7.13 (1H, dd), 7.19 (1H, ddd), 7.35 (1H, dd), 7.49 (2H, dm), 7.56 (2H, dm). MS: 254 (M⁺), 239, 211 (100%).

c)


The title compound (280 mg, 59%) was prepared as a colourless solid from (*E*)-3-(3'-methoxy-biphenyl-4-yl)-but-2-en-1-ol (254 mg, 1.00 mmol) and (*S*)-ethyl 2-ethoxy-3-(4-hydroxyphenyl)-propionate (250 mg, 1.05 mmol) by a procedure analogous to that described in example 52c.

¹H NMR (300 MHz, CDCl₃) δ: 1.17 (3H, t), 1.22 (3H, t), 2.17 (3H, d), 2.96 (2H, d), 3.30-3.42 (1H, m), 3.55-3.67 (1H, m), 3.87 (3H, s), 3.98 (1H, t), 4.17 (2H, q), 4.74 (2H, d), 6.12 (1H, tm), 6.85-6.92 (3H, m), 7.11-7.22 (4H, m), 7.35 (1H, dd), 7.47-7.59 (4H, m). MS: 474 (M⁺), 237 (100%). Microanalysis Calculated % C: 75.92, H: 7.22. Found % C: 76.04, H: 7.39.

EXAMPLE 1145 (E)-(S)-2-Ethoxy-3-{4-[3-(3'-methoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionic acid

The title compound was prepared from (E)-(S)-ethyl 2-ethoxy-3-{4-[3-(3'-methoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionate (example 113) (230 mg, 0.49 mmol) and sodium hydroxide (1M, 0.97 ml, 0.97 mmol) by a procedure analogous to that described in example 51, yielding (E)-(S)-2-ethoxy-3-{4-[3-(3'-methoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionic acid (183 mg, 85%) as a colourless solid.

¹H NMR (300 MHz, CDCl₃) δ: 1.18 (3H, t), 2.17 (3H, d), 2.97 (1H, dd), 3.10 (1H, dd); 3.41-3.53 (1H, m), 3.53-3.65 (1H, m), 3.87 (3H, s), 4.07 (1H, dd), 4.74 (2H, dd), 6.11 (1H, tm), 6.86-6.93 (3H, m), 7.11-7.22 (4H, m), 7.35 (1H, dd), 7.47-7.59 (4H, m), carboxylic acid proton not observed. LCMS: 683 (M+237), 469 (M+Na), 237 (100%).

EXAMPLE 11520 (E)-(S,S/R)-Ethyl 2-Ethoxy-3-(4-{3-[3'-(1-hydroxy-ethyl)-biphenyl-4-yl]-but-2-enyloxy}-phenyl)-propionate

a)

The colourless solid (*E*)-(*S/R*)-3-[3'-(1-hydroxy-ethyl)-biphenyl-4-yl]-but-2-en-1-ol was prepared by DIBAL-H reduction of (*E*)-1-[4'-(3-hydroxy-1-methyl-propenyl)-biphenyl-3-yl]-ethanone (example 105a) by a procedure analogous to that described in example 52b.

- 5 Mpt. 94-100°C. ¹H NMR (300 MHz, DMSO-d₆) δ: 1.37 (3H, d), 2.02 (3H, d), 4.18 (2H, br dd), 4.75 (1H, br t, OH), 4.78 (1H, dq), 5.21 (1H, d, OH), 5.98 (1H, tm), 7.33 (1H, dm), 7.40 (1H, dd), 7.48-7.54 (3H, m), 7.60-7.66 (3H, m). MS: 268 (100%, M⁺), 253, 235, 225. Microanalysis
Calculated % C: 80.56, H: 7.51. Found % C: 80.21, H: 7.78.


10 b)

The title compound (490 mg, 57%) was prepared as a colourless oil from (*E*)-(*S/R*)-3-[3'-(1-hydroxy-ethyl)-biphenyl-4-yl]-but-2-en-1-ol (500 mg, 1.86 mmol) and (*S*)-ethyl 2-ethoxy-3-(4-hydroxyphenyl)-propionate (422 mg, 1.77 mmol) by a procedure analogous to that described in example 52c.

- 15 ¹H NMR (300 MHz, CDCl₃) δ: 1.17 (3H, t), 1.22 (3H, t), 1.56 (3H, d), 1.89 (1H, d, OH), 2.18 (3H, d), 2.97 (2H, d), 3.30-3.42 (1H, m), 3.54-3.66 (1H, m), 3.98 (1H, t), 4.17 (2H, q), 4.75 (2H, d), 4.99 (1H, dq), 6.12 (1H, tm), 6.85 (2H, dm), 7.18 (2H, dm), 7.32-7.48 (2H, m), 7.48-7.67 (6H, m).

20

EXAMPLE 116




- 25 (*E*)-(*S,S/R*)-2-Ethoxy-3-(4-{3-[3'-(1-hydroxy-ethyl)-biphenyl-4-yl]-but-2-enyloxy}-phenyl)-propionic acid
-

The title compound was prepared from (*E*)-(*S,S/R*)-ethyl 2-ethoxy-3-(4-{3-[3'-(1-hydroxy-ethyl)-biphenyl-4-yl]-but-2-enyloxy}-phenyl)-propionate (example 115) (460 mg, 0.94 mmol)
30 and sodium hydroxide (1M, 1.9 ml, 1.9 mmol) by a procedure analogous to that described in

example 51, yielding (*E*)-(S,S/*R*)-ethyl 2-ethoxy-3-(4-{3-[3'-(1-hydroxy-ethyl)-biphenyl-4-yl]-but-2-enyloxy}-phenyl)-propionic acid (434 mg, 100%) as a colourless gum.

¹H NMR (300 MHz, CDCl₃) δ: 1.18 (3H, t), 1.55 (3H, d), 2.17 (3H, d), 2.96 (1H, dd), 3.09 (1H, dd), 3.41-3.53 (1H, m), 3.54-3.66 (1H, m), 4.06 (1H, dd), 4.75 (2H, d), 4.98 (1H, q), 6.12 (1H, tm), 6.90 (2H, dm), 7.17 (2H, dm), 7.32-7.46 (2H, m), 7.47-7.63 (6H, m), carboxylic acid proton not observed.

EXAMPLE 117



(*E*)-(S)-Ethyl 3-[4-[3-(3,5-Dibromophenyl)-but-2-enyloxy]-phenyl]-2-ethoxy-propionate

15 a)

Sodium (0.49 g, 21.3 mmol) was added to ethanol (50 ml) at room temperature and the mixture stirred until the metal had fully reacted. Triethyl phosphonoacetate (5.49 g, 24.5 mmol) was added, the solution stirred for 15 min, then an ethanol (100 ml) solution of 3,5-dibromoacetophenone (4.60 g, 16.6 mol) was added and the reaction mixture heated to 80°C under reflux for 72 h. The solution was cooled, the ethanol evaporated and the resulting orange residue partitioned between 1N HCl (150 ml) and ethyl acetate (150 ml). The aqueous layer was collected and further extracted with ethyl acetate (2 x 100 ml). The organic layers were combined, washed with brine, dried (MgSO₄) and evaporated to an orange/yellow gum, which was purified by column chromatography on silica gel (3% diethyl ether in n-heptane eluent) to give the product, (*E*)-ethyl 3-(3,5-dibromophenyl)-but-2-enoate, as a colourless wax; 4.06 g (70%).

¹H NMR (300 MHz, CDCl₃) δ: 1.32 (3H, t), 2.51 (3H, d), 4.22 (2H, q), 6.09 (1H, m), 7.52 (2H, d), 7.64 (1H, t). MS: 350/348/346 (M⁺), 304/302/300 (M-EtOH), 115 (100%). Microanalysis Calculated % C: 41.41, H: 3.48, Br: 45.92. Found % C: 41.75, H: 3.52, Br: 45.62.

b)

(*E*)-Ethyl 3-(3,5-dibromophenyl)-but-2-enoate was reduced with DIBAL-H by a procedure analogous to that described in example 50b, to give the colourless oil (*E*)-3-(3,5-dibromophenyl)-but-2-en-1-ol.


- 5 ¹H NMR (300 MHz, CDCl₃) δ: 1.64 (1H, br s), 2.01 (3H, d), 4.36 (2H, d), 5.96 (1H, tm), 7.46 (2H, d), 7.54 (1H, t). MS: 308/306/304 (M⁺), 293/291/289 (M-Me), 266/264/262, 227/225/223, 131, 128, 115 (100%), 102.

c)

- 10 The title compound (851 mg, 81%) was prepared from (*E*)-3-(3,5-dibromo-phenyl)-but-2-en-1-ol (612 mg, 2.0 mmol) and (*S*)-ethyl 2-ethoxy-3-(4-hydroxyphenyl)-propionate (500 mg, 2.10 mmol) by a procedure analogous to that described in example 52c.
- ¹H NMR (300 MHz, CDCl₃) δ: 1.17 (3H, t), 1.23 (3H, t), 2.09 (3H, d), 2.96 (2H, d), 3.30-3.42 (1H, m), 3.54-3.66 (1H, m), 3.98 (1H, t), 4.17 (2H, q), 4.69 (2H, d), 6.04 (1H, tm), 6.85 (2H, dm), 7.16 (2H, dm), 7.48 (2H, d), 7.57 (1H, t). LCMS: 551/549/547 (100%, M+Na), 546/544/542 (M+NH₄), 483/481/479 (M+H-EtOH).
- 15

EXAMPLE 118

20




(*E*)-(S)-3-{4-[3-(3,5-Dibromophenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid

- 25 The title compound was prepared from (*E*)-(S)-ethyl 3-{4-[3-(3,5-dibromophenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate (example 117) (840 mg, 1.60 mmol) and sodium hydroxide (1M, 16 ml, 16 mmol) by a procedure analogous to that described in example 51, yielding (*E*)-(S)-3-{4-[3-(3,5-dibromophenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid (781 mg, 98%) as a colourless gum.

¹H NMR (300 MHz, CDCl₃) δ: 1.19 (3H, t), 2.08 (3H, d), 2.96 (1H, dd), 3.09 (1H, dd), 3.41-3.53 (1H, m), 3.55-3.67 (1H, m), 4.07 (1H, dd), 4.70 (2H, d), 6.04 (1H, tm), 6.87 (2H, dm), 7.17 (2H, dm), 7.48 (2H, d), 7.58 (1H, t), carboxylic acid proton not observed. LCMS: 523/521/519 (100%, M+Na), 518/516/514 (M+NH₄), 455/453/451 (M+H-EtOH), 291/289/287.

5

EXAMPLE 119



10

(E)-(S)-Ethyl 3-{4-[3-(3,5-Dibromophenyl)-allyloxy]-phenyl}-2-ethoxy-propionate

a)

(E)-Ethyl 3-(3,5-dibromophenyl)-acrylate (example 109a) was reduced with DIBAL-H by a procedure analogous to that described in example 50b, to give the colourless solid (E)-3-(3,5-dibromophenyl)-prop-2-en-1-ol.

15

¹H NMR (300 MHz, CDCl₃) δ: 1.52 (1H, t, OH), 4.35 (2H, ddd), 6.36 (1H, dt), 6.50 (1H, dm), 7.44 (2H, d), 7.53 (1H, t). LCMS: 277/275/273 (100%, M+H-H₂O), 196/194 (M+H-H₂O-Br), 100.

20


b)

The title compound (780 mg, 78%) was prepared from (E)-3-(3,5-dibromophenyl)-prop-2-en-1-ol (584 mg, 2.0 mmol) and (S)-ethyl 2-ethoxy-3-(4-hydroxyphenyl)-propionate (500 mg, 2.10 mmol) by a procedure analogous to that described in example 52c.

25

¹H NMR (300 MHz, CDCl₃) δ: 1.17 (3H, t), 1.22 (3H, t), 2.96 (2H, d), 3.30-3.42 (1H, m), 3.55-3.67 (1H, m), 3.97 (1H, t), 4.17 (2H, q), 4.68 (2H, dd), 6.41 (1H, dt), 6.59 (1H, dm), 6.86 (2H, dm), 7.17 (2H, dm), 7.46 (2H, d), 7.53 (1H, t).

30


EXAMPLE 120

5 (E)-(S)-3-{4-[3-(3,5-Dibromophenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid

The title compound was prepared from (E)-(S)-ethyl 3-{4-[3-(3,5-dibromophenyl)-allyloxy]-phenyl}-2-ethoxy-propionate (example 119) (512 mg, 1.0 mmol) and sodium hydroxide (1M, 10 ml, 10 mmol) by a procedure analogous to that described in example 51, yielding (E)-(S)-
 10 3-{4-[3-(3,5-dibromophenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid (96 mg; 20%) as a colourless gum.

¹H NMR (300 MHz, CDCl₃) δ: 1.19 (3H, t), 2.97 (1H, dd), 3.09 (1H, dd), 3.40-3.66 (2H, m), 4.06 (1H, dd), 4.68 (2H, dd), 6.42 (1H, dt), 6.59 (1H, dm), 6.88 (2H, dm), 7.18 (2H, dm), 7.46 (2H, d), 7.58 (1H, t), carboxylic acid proton not observed. LCMS: 509/507/505 (M+Na),

15 504/502/500 (100%, M+NH₄), 441/439/437 (M+H-EtOH), 277/275/273.

EXAMPLE 121

20

(E)-(S)-Ethyl 3-{4-[3-(4,4''-Di-tert-butyl-[1,1';3',1'']terphenyl-5'-yl)-allyloxy]-phenyl}-2-ethoxy-propionate

a)

The colourless glass, (*E*)-ethyl 3-(4,4''-di-*tert*-butyl-[1,1';3',1'']terphenyl-5'-yl)-acrylate was prepared from (*E*)-ethyl 3-(3,5-dibromophenyl)-acrylate (example 109a) and 4-*tert*-butylphenylboronic acid by a procedure analogous to that described in example 52a.

- 5 ¹H NMR (300 MHz, CDCl₃) δ: 1.36 (3H, t), 1.38 (18H, s), 4.29 (2H, q), 6.55 (1H, d), 7.43-7.52 (4H, m), 7.52-7.61 (4H, m), 7.70 (2H, d), 7.81 (1H, t), 7.82 (1H, d). MS: 440 (M⁺), 425 (100%, M-Me), 205.

b)


- 10 The colourless gum (*E*)-3-(4,4''-di-*tert*-butyl-[1,1';3',1'']terphenyl-5'-yl)-prop-2-en-1-ol was prepared by DIBAL-H reduction of (*E*)-ethyl 3-(4,4''-di-*tert*-butyl-[1,1';3',1'']terphenyl-5'-yl)-acrylate by a procedure analogous to that described for example 52b.

- 15 ¹H NMR (300 MHz, CDCl₃) δ: 1.37 (18H, s), 1.48 (1H, br t), 4.37 (2H, m), 6.49 (1H, dt), 6.75 (1H, dm), 7.45-7.52 (4H, m), 7.54-7.61 (6H, m), 7.68 (1H, t). LCMS: 779 (M+381), 761 (779-H₂O), 437, 421 (M+Na), 399 (M+H), 381 (100%, M+H-H₂O).

c)

- 20 The title compound (368 mg, 79%) was prepared from (*E*)-3-(4,4''-di-*tert*-butyl-[1,1';3',1'']terphenyl-5'-yl)-prop-2-en-1-ol (300 mg, 0.75 mmol) and (*S*)-ethyl 2-ethoxy-3-(4-hydroxyphenyl)-propionate (188 mg, 0.79 mmol) by a procedure analogous to that described in example 52c.


- 25 ¹H NMR (300 MHz, CDCl₃) δ: 1.17 (3H, t), 1.22 (3H, t), 1.37 (18H, s), 2.96 (2H, d), 3.30-3.43 (1H, m), 3.54-3.67 (1H, m), 3.98 (1H, t), 4.16 (2H, q), 4.72 (2H, d), 6.52 (1H, dt), 6.85 (1H, d), 6.90 (2H, dm), 7.18 (2H, dm), 7.44-7.52 (4H, m), 7.54-7.63 (6H, m), 7.69 (1H, m). LCMS: 641 (100%, M+Na), 636 (M+NH₄), 381.

EXAMPLE 122

- 5 (E)-(S)-3-{4-[3-(4,4''-Di-*tert*-butyl-[1,1';3',1'']terphenyl-5'-yl)-allyloxy]-phenyl}-2-ethoxy-propionic acid

The title compound was prepared from (E)-(S)-ethyl 3-{4-[3-(4,4''-di-*tert*-butyl-[1,1';3',1'']terphenyl-5'-yl)-allyloxy]-phenyl}-2-ethoxy-propionate (example 121) (345 mg, 0.56 mmol) and sodium hydroxide (1M, 1.1 ml, 1.1 mmol) by a procedure analogous to that described in example 51, yielding (E)-(S)-3-{4-[3-(4,4''-di-*tert*-butyl-[1,1';3',1'']terphenyl-5'-yl)-allyloxy]-phenyl}-2-ethoxy-propionic acid (284 mg, 86%) as a colourless foam.

¹H NMR (300 MHz, CDCl₃) δ: 1.18 (3H, t), 1.38 (18H, s), 2.97 (1H, dd), 3.10 (1H, dd), 3.41-3.65 (2H, m), 4.07 (1H, dd), 4.72 (2H, dm), 6.52 (1H, dt), 6.85 (1H, d), 6.92 (2H, dm), 7.17 (2H, dm), 7.44-7.52 (4H, m), 7.55-7.62 (6H, m), 7.69 (1H, m), carboxylic acid proton not observed.

EXAMPLE 123

(E)-(S)-Ethyl 3-{4-[3-(3',5'-Dibromo-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate

a)

The colourless gum (*E*)-ethyl 3-(3',5'-dibromo-biphenyl-4-yl)-but-2-enoate was prepared from (*E*)-ethyl 3-(4-iodophenyl)-but-2-enoate (example 91a) and 3,5-dibromobenzene boronic acid

5 by a procedure analogous to that described in example 52a.

¹H NMR (300 MHz, CDCl₃) δ: 1.33 (3H, t), 2.61 (3H, d), 4.22 (2H, q), 6.20 (1H, m), 7.44-7.80 (7H, m). LCMS: 427/425/423 (100%, M+H), 381/379/377 (M+H-EtOH).

b)

10 The colourless gum (*E*)-3-(3',5'-dibromo-biphenyl-4-yl)-but-2-en-1-ol was prepared by DIBAL-H reduction of (*E*)-ethyl 3-(3',5'-dibromo-biphenyl-4-yl)-but-2-enoate as described for example 52b, with the purification of the product being carried out by preparative HPLC.

¹H NMR (300 MHz, CDCl₃) δ: 1.45 (1H, br s), 2.12 (3H, d), 4.41 (2H, d), 6.05 (1H, tm), 7.45-7.54 (4H, m), 7.62 (1H, t), 7.66 (2H, d). LCMS: 367/365/363 (100%, M+H-H₂O), 286/284.

15

c)


The title compound (177 mg, 83%) was prepared as a colourless gum from (*E*)-3-(3',5'-dibromo-biphenyl-4-yl)-but-2-en-1-ol (135 mg, 0.35 mmol) and (*S*)-ethyl 2-ethoxy-3-(4-hydroxyphenyl)-propionate (90 mg, 0.38 mmol) by a procedure analogous to that described

20 in example 52c.

¹H NMR (300 MHz, CDCl₃) δ: 1.17 (3H, t), 1.23 (3H, t), 2.16 (3H, d), 2.96 (2H, d), 3.31-3.43 (1H, m), 3.55-3.67 (1H, m), 3.98 (1H, t), 4.17 (2H, q), 4.74 (2H, d), 6.13 (1H, tm), 6.88 (2H, dm), 7.17 (2H, dm), 7.45-7.56 (4H, m), 7.63 (1H, t), 7.66 (2H, d). LCMS: 627/625/623 (100%, M+Na), 365.

25

EXAMPLE 124




30

(E)-(S)-3-{4-[3-(3',5'-Dibromo-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid

The title compound was prepared from (E)-(S)-ethyl 3-{4-[3-(3',5'-dibromo-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate (example 123) (110 mg, 0.18 mmol) and sodium hydroxide (1M, 1.0 ml, 1.0 mmol) by a procedure analogous to that described in example 51, yielding (E)-(S)-3-{4-[3-(3',5'-dibromo-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid (90 mg, 86%) as a colourless gum.

¹H NMR (300 MHz, CDCl₃) δ: 1.20 (3H, t), 2.18 (3H, d), 2.99 (1H, dd), 3.12 (1H, dd), 3.43-3.68 (2H, m), 4.08 (1H, dd), 4.75 (2H, d), 6.13 (1H, tm), 6.91 (2H, dm), 7.19 (2H, dm), 7.45-7.60 (4H, m), 7.60-7.74 (3H, m), carboxylic acid proton not observed. LCMS: 599/597/595 (100%, M+Na), 365.

EXAMPLE 125



(E)-(S)-Ethyl 3-{4-[3-(3',5'-Dichloro-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate

20 a)

The colourless solid (E)-ethyl 3-(3',5'-dichloro-biphenyl-4-yl)-but-2-enoate was prepared from (E)-ethyl 3-(4-iodophenyl)-but-2-enoate (example 91a) and 3,5-dichlorobenzene boronic acid by a procedure analogous to that described in example 52a.

Mpt. 96.3-97.3°C. ¹H NMR (300 MHz, CDCl₃) δ: 1.33 (3H, t), 2.61 (3H, d), 4.23 (2H, q), 6.19 (1H, m), 7.35 (1H, t), 7.47 (2H, d), 7.51-7.61 (4H, m). LCMS: 335/337/339 (100%, M+H), 289/291/293 (M+H-EtOH). Microanalysis Calculated % C: 64.49, H: 4.81, Cl: 21.15; found C: 64.41, H: 4.80, Cl: 20.80.

b)

The colourless oil (*E*)-3-(3',5'-dichloro-biphenyl-4-yl)-but-2-en-1-ol was prepared by DIBAL-H reduction of (*E*)-ethyl 3-(3',5'-dichloro-biphenyl-4-yl)-but-2-enoate as described for example 52b.


¹H NMR (300 MHz, CDCl₃) δ: 1.39 (1H, br s), 2.12 (3H, d), 4.40 (2H, d), 6.05 (1H, tm), 7.33 (1H, t), 7.46 (2H, d), 7.45-7.65 (4H, m). MS: 296/294/292 (100%, M⁺), 281/279/277 (M-Me), 278/276/274 (M-H₂O), 253/251/249.

c)

The title compound (794 mg, 77%) was prepared as a colourless gum from (*E*)-3-(3',5'-dichloro-biphenyl-4-yl)-but-2-en-1-ol (586 mg, 2.0 mmol) and (*S*)-ethyl 2-ethoxy-3-(4-hydroxyphenyl)-propionate (500 mg, 2.10 mmol) by a procedure analogous to that described in example 52c.

¹H NMR (300 MHz, CDCl₃) δ: 1.17 (3H, t), 1.23 (3H, t), 2.17 (3H, d), 2.96 (2H, d), 3.30-3.42 (1H, m), 3.55-3.67 (1H, m), 3.98 (1H, t), 4.17 (2H, q), 4.74 (2H, d), 6.13 (1H, tm), 6.88 (2H, dm), 7.18 (2H, dm), 7.33 (1H, t), 7.46 (2H, d), 7.48-7.54 (4H, m). LCMS: 539/537/535 (100%, M+Na), 534/532/530 (M+NH₄), 279/277/275.

EXAMPLE 126




(*E*)-(S)-3-{4-[3-(3',5'-Dichloro-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid

The title compound was prepared from (*E*)-(S)-ethyl 3-{4-[3-(3',5'-dichloro-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate (example 125) (513 mg, 1.0 mmol) and sodium hydroxide (1M, 5.0 ml, 5.0 mmol) by a procedure analogous to that described in example 51, yielding (*E*)-(S)-3-{4-[3-(3',5'-dichloro-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid (422 mg, 87%) as a colourless glass.

¹H NMR (300 MHz, CDCl₃) δ: 1.19 (3H, t), 2.16 (3H, d), 2.97 (1H, dd), 3.09 (1H, dd), 3.40-3.54 (H, m), 3.54-3.67 (1H, m), 4.07 (1H, dd), 4.74 (2H, d), 6.12 (1H, tm), 6.90 (2H, dm), 7.18 (2H, dm), 7.32 (1H, t), 7.46 (2H, d), 7.48-7.55 (4H, m), carboxylic acid proton not observed. LCMS: 511/509/507 (100%, M+Na), 279/277/275.

5

EXAMPLE 127



10

(*E*)-(*S*)-Ethyl 3-{4-[3-(3',5'-Dichloro-biphenyl-4-yl)-allyloxy]-phenyl}-2-ethoxy-propionate

a)

The colourless solid (*E*)-ethyl 3-(3',5'-dichloro-biphenyl-4-yl)-acrylate was prepared from (*E*)-ethyl 3-(4-bromo-phenyl)-acrylate (example 71a) and 3,5-dichlorobenzene boronic acid by a procedure analogous to that described in example 52a.

Mpt. 70-78°C. ¹H NMR (300 MHz, CDCl₃) δ: 1.35 (3H, t), 4.29 (2H, q), 6.49 (1H, d), 7.36 (1H, t), 7.47 (2H, d), 7.53-7.64 (4H, m), 7.71 (1H, d). LCMS: 325/323/321 (100%, M+H). Micro-analysis Calculated % C: 63.57, H: 4.39; found C: 63.37, H: 4.43.

20

b)

The colourless gum (*E*)-3-(3',5'-dichloro-biphenyl-4-yl)-prop-2-en-1-ol was prepared by DIBAL-H reduction of (*E*)-ethyl 3-(3',5'-dichloro-biphenyl-4-yl)-acrylate as described for example 52b.

¹H NMR (300 MHz, CDCl₃) δ: 1.47 (1H, br t), 4.36 (2H, ddd), 6.43 (1H, dt), 6.65 (1H, dm), 7.33 (1H, t), 7.37-7.55 (6H, m). MS: 282/280/278 (100%, M⁺), 239/237/235, 226/224/222.

c)


The title compound (732 mg, 70%) was prepared as a yellow gum, (containing 0.25 molar equivalents of ethyl acetate) from (*E*)-3-(3',5'-dichloro-biphenyl-4-yl)-prop-2-en-1-ol (559 mg,

30

2.0 mmol) and (S)-ethyl 2-ethoxy-3-(4-hydroxyphenyl)-propionate (500 mg, 2.10 mmol) by a procedure analogous to that described in example 52c.


¹H NMR (300 MHz, CDCl₃) δ: 1.17 (3H, t), 1.22 (3H, t), 1.25 (0.75H, t, AcOEt), 2.04 (0.75H, s, AcOEt), 2.97 (2H, d), 3.30-3.41 (1H, m), 3.53-3.67 (1H, m), 3.98 (1H, t), 4.12 (0.5H, q, AcOEt), 4.17 (2H, q), 4.71 (2H, dd), 6.47 (1H, dt), 6.76 (1H, dm), 6.89 (2H, dm), 7.17 (2H, dm), 7.33 (1H, t), 7.43-7.55 (6H, m). LCMS: 525/523/521 (100%, M+Na), 265/263/261.

EXAMPLE 128



(E)-(S)-3-{4-[3-(3',5'-Dichloro-biphenyl-4-yl)-allyloxy]-phenyl}-2-ethoxy-propionic acid

- 15 The title compound was prepared from (E)-(S)-ethyl 3-{4-[3-(3',5'-dichloro-biphenyl-4-yl)-allyloxy]-phenyl}-2-ethoxy-propionate (example 127) (522 mg, 1.0 mmol) and sodium hydroxide (1M, 10.0 ml, 10.0 mmol) by a procedure analogous to that described in example 51, yielding (E)-(S)-3-{4-[3-(3',5'-dichloro-biphenyl-4-yl)-allyloxy]-phenyl}-2-ethoxy-propionic acid (325 mg, 67%) as a colourless wax, which contained 0.167 molar equivalents of AcOEt.
- 20 ¹H NMR (300 MHz, CDCl₃) δ: 1.19 (3H, t), 1.26 (0.5H, t, AcOEt), 2.04 (0.5H, s, AcOEt), 2.97 (1H, dd), 3.09 (1H, dd), 3.41-3.53 (1H, m), 3.53-3.68 (1H, m), 4.07 (1H, dd), 4.12 (0.33H, q, AcOEt), 4.71 (2H, dd), 6.48 (1H, dt), 6.76 (1H, dm), 6.91 (2H, dm), 7.18 (2H, dm), 7.33 (1H, t), 7.40-7.60 (6H, m), carboxylic acid proton not observed. LCMS: 497/495/493 (100%, M+Na), 492/490/488 (M+NH₄), 265/263/261.

EXAMPLE 129

5 (E)-(S)-Ethyl 3-{4-[3-(3',5'-di-*tert*-butyl-biphenyl-4-yl)-allyloxy]-phenyl}-2-ethoxy-propionate

a)

The colourless solid (*E*)-3-(4-bromo-phenyl)-prop-2-en-1-ol was prepared by DIBAL-H reduction of (*E*)-ethyl 3-(4-bromo-phenyl)-acrylate (example 71a) as described for example 52b.

10 Mpt. 65.5-67.5°C. ¹H NMR (300 MHz, CDCl₃) δ: 1.50 (1H, br s), 4.33 (2H, d), 6.35 (1H, dt), 6.55 (1H, d), 7.23 (2H, dm), 7.43 (2H, dm). MS: 214/212 (M⁺), 171/169, 158/156, 133 (M-Br, 100%), 115, 91, 77.

b)

15 *tert*-Butyl chlorodimethylsilane (1.33 g, 19.5 mmol) was added to a stirred solution of (*E*)-3-(4-bromo-phenyl)-prop-2-en-1-ol (3.20 g, 15.0 mmol), and imidazole (2.72 g, 18.0 mmol) in dry dichloromethane (75 ml) and the resulting mixture stirred at room temperature for 18 h, a colourless precipitate being formed. The mixture was diluted with dichloromethane (100 ml) and 1N hydrochloric acid (100 ml). The aqueous layer was separated, further extracted with
20 dichloromethane (2 x 100ml) and the combined organic layers washed with brine, dried (MgSO₄) and evaporated. The product was purified by column chromatography on silica gel (1% diethyl ether in n-heptane eluent) to give the colourless solid, (*E*)-[3-(4-bromo-phenyl)-allyloxy]-*tert*-butyldimethylsilane (4.39 g, 89%).

Mpt. 46.5-48°C. ¹H NMR (300 MHz, CDCl₃) δ: 0.11 (6H, s), 0.94 (9H, s), 4.34 (2H, dd), 6.27
25 (1H, dt), 6.54 (1H, dt), 7.24 (2H, dm), 7.42 (2H, dm), 7.71 (1H, d). Microanalysis Calculated % C: 55.04, H: 7.08, Br: 24.41; found C: 54.81, H: 7.22, Br: 24.51.

c) *tert*-Butyllithium (1.7M in pentane, 3.5 ml, 6.0 mmol) was added dropwise, at -78°C to a stirred THF (10 ml) solution of (*E*)-[3-(4-bromo-phenyl)-allyloxy]-*tert*-butyldimethylsilane (982
30 mg, 3.0 mmol) and the resulting solution stirred for 45 min. Trimethylborate (0.51 ml, 4.50

mmol) was added, the solution allowed to warm to room temperature over 2h, and the solvents evaporated to give the crude boronate ester as a yellow gum, which was dissolved in DME (10 ml). Tetrakis(triphenylphosphine)palladium(0) (69 mg, 0.06 mmol), was added to a stirred DME (20 ml) solution of 1-bromo-3,5-di-*tert*-butylbenzene (538 mg, 2.0 mmol), the solution stirred for 10 min, aqueous sodium carbonate (2M, 9 ml, 18.0 mmol) added, and stirring continued for a further 10 min. The boronate ester solution was added and the mixture heated to 80°C, under reflux, for 24h. The resulting mixture was diluted with 1N HCl (50 ml), the products extracted into ethyl acetate (3 x 50 ml), and the combined extracts washed with brine, dried (MgSO₄), and evaporated. The resulting yellow gum was dissolved in dry THF (20 ml), tetra-*n*-butyl ammonium fluoride (1.26 g, 4.0 mmol) added and the solution stirred at room temperature for 18h. The resulting mixture was diluted with 1N HCl (50 ml), and the products extracted into ethyl acetate (2 x 50 ml). The combined organic phases were washed with brine, dried (MgSO₄), and evaporated to give the colourless glass, (*E*)-3-(3',5'-di-*tert*-butyl-biphenyl-4-yl)-prop-2-en-1-ol (127 mg, 20%).


¹H NMR (300 MHz, CDCl₃) δ: 1.38 (18H, s), 1.45 (1H, br t), 4.35 (2H, br t), 6.41 (1H, dt), 6.66 (1H, br dm), 7.40-7.49 (5H, m), 7.53-7.58 (2H, m). MS: 322 (M⁺), 307 (100%, M-Me), 57.

d)

The title compound (110 mg, 51%) was prepared as a colourless gum from (*E*)-3-(3',5'-di-*tert*-butyl-biphenyl-4-yl)-prop-2-en-1-ol (127 mg, 0.39 mmol) and (*S*)-ethyl 2-ethoxy-3-(4-hydroxyphenyl)-propionate (99 mg, 0.41 mmol) by a procedure analogous to that described in example 52c.

¹H NMR (300 MHz, CDCl₃) δ: 1.17 (3H, t), 1.23 (3H, t), 1.38 (18H, s), 2.97 (2H, d), 3.29-3.42 (1H, m), 3.53-3.67 (1H, m), 3.98 (1H, t), 4.17 (2H, q), 4.70 (2H, dd), 6.45 (1H, dt), 6.77 (1H, dm), 6.90 (2H, dm), 7.18 (2H, dm), 7.40-7.60 (7H, m).

EXAMPLE 130




(E)-(S)-3-{4-[3-(3',5'-Di-*tert*-butyl-biphenyl-4-yl)-allyloxy]-phenyl}-2-ethoxy-propionic acid

The title compound was prepared from (E)-(S)-ethyl 3-{4-[3-(3',5'-di-*tert*-butyl-biphenyl-4-yl)-allyloxy]-phenyl}-2-ethoxy-propionate (example 129) (110 mg, 0.20 mmol) and sodium hydroxide (1M, 0.8 ml, 0.8 mmol) by a procedure analogous to that described in example 51, yielding ((E)-(S)-3-{4-[3-(3',5'-di-*tert*-butyl-biphenyl-4-yl)-allyloxy]-phenyl}-2-ethoxy-propionic acid (92 mg, 88%) as a colourless solid.

¹H NMR (300 MHz, CDCl₃) δ: 1.20 (3H, t), 1.38 (18H, s), 2.97 (1H, dd), 3.10 (1H, dd), 3.40-3.65 (2H, m), 4.07 (1H, dd), 4.70 (2H, dd), 6.45 (1H, dt), 6.77 (1H, dm), 6.90 (2H, dm), 7.18 (2H, dm), 7.38-7.60 (7H, m), carboxylic acid proton not observed.

EXAMPLE 131



(E)-(S)-Ethyl 3-{4-[3-(3',5'-Di-*tert*-butyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate

20

a)

The colourless oil, (E)-[3-(4-bromophenyl)-but-2-enyloxy]-*tert*-butyl-dimethylsilane was prepared from (E)-3-(4-bromophenyl)-but-2-en-1-ol (example 50b), imidazole and *tert*-butyl chlorodimethylsilane by a procedure analogous to that described in example 129b.

25 ¹H NMR (300 MHz, CDCl₃) δ: 0.10 (6H, s), 0.92 (9H, s), 4.37 (2H, d), 5.88 (1H, tm), 7.25 (2H, dm), 7.42 (2H, dm). MS: 342/340 (M⁺), 327/325 (M-Me), 285/283 (M-Bu), 130, 75 (100%).

b)

The colourless wax, (*E*)-3-(3',5'-di-*tert*-butyl-biphenyl-4-yl)-but-2-en-1-ol was prepared via a metallation, boronation, cross coupling and deprotection sequence analogous to that described for example 129c.


¹H NMR (300 MHz, CDCl₃) δ: 1.26 (1H, br m), 1.38 (18H, s), 2.12 (3H, d), 4.40 (2H, br t),
5 6.05 (1H, dt), 7.40-7.42 (3H, m), 7.43-7.59 (4H, m). LCMS: 331 (M+H), 319 (100%, M+H-H₂O).

c)

The title compound (429 mg, 74%) was prepared as a colourless gum from (*E*)-3-(3',5'-di-*tert*-butyl-biphenyl-4-yl)-but-2-en-1-ol (350 mg, 1.04 mmol) and (*S*)-ethyl 2-ethoxy-3-(4-hydroxyphenyl)-propionate (260 mg, 1.09 mmol) by a procedure analogous to that described in example 52c.

¹H NMR (300 MHz, CDCl₃) δ: 1.17 (3H, t), 1.22 (3H, t), 1.38 (18H, s), 2.17 (3H, d), 2.96 (2H, d), 3.30-3.43 (1H, m), 3.55-3.68 (1H, m), 3.98 (1H, t), 4.17 (2H, q), 4.75 (2H, d), 6.11 (1H, tm), 6.89 (2H, dm), 7.17 (2H, dm), 7.40-7.45 (3H, m), 7.46-7.60 (4H, m). LCMS: 579 (100%, M+Na), 574 (M+NH₄), 511 (M+H-EtOH).

EXAMPLE 132




(*E*)-(S)-3-(4-[3-(3',5'-Di-*tert*-butyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl)-2-ethoxy-propionic acid

The title compound was prepared from (*E*)-(S)-ethyl 3-{4-[3-(3',5'-di-*tert*-butyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate (example 131) (400 mg, 0.72 mmol) and sodium hydroxide (1M, 2.9 ml, 2.9 mmol) by a procedure analogous to that described in example 51,

yielding (*E*)-(*S*)-3-[4-[3-(3',5'-di-*tert*-butyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl]-2-ethoxy-propionic acid (315 mg, 83%) as a colourless gum.

¹H NMR (300 MHz, CDCl₃) δ: 1.19 (3H, t), 1.39 (18H, s), 2.18 (3H, d), 2.97 (1H, dd), 3.10 (1H, dd), 3.40-3.67 (2H, m), 4.07 (1H, dd), 4.70 (2H, d), 6.11 (1H, tm), 6.90 (2H, dm), 7.18 (2H, dm), 7.40-7.45 (3H, m), 7.46-7.60 (4H, m), carboxylic acid proton not observed. LCMS: 847 (M+319), 551 (M+Na), 319 (100%).

EXAMPLE 133



(*E*)-(*S/R*)-Ethyl 3-[4-(3-Biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-isopropoxy-propionate

15 a)

(*S/R*)-Ethyl 2-(diethoxyphosphoryl)-2-isopropoxy-acetate was prepared as a pale green oil by the rhodium(II) acetate dimer catalysed reaction of ethyl diazo-(diethoxyphosphoryl)-acetate with isopropyl alcohol, according to the method described by C.J. Moody et al (*Tetrahedron*, 1992, 48, 3991-4004).

20 ¹H NMR (300 MHz, CDCl₃) δ: 1.21 (3H, d), 1.23 (3H, d), 1.28-1.39 (9H, m), 3.74 (1H, septet), 4.15-4.35 (6H, m), 4.39 (1H, d, J_{HP} = 19.9 Hz). LCMS: 283 (M+H), 241 (100%), 213.

b)

A THF (20 ml) solution of (*S/R*)-ethyl 2-(diethoxyphosphoryl)-2-isopropoxy-acetate (6.40 g, 22.7 mmol) was added dropwise, at 0°C, to a stirred suspension of sodium hydride (60% dispersion in mineral oil, 0.92 g, 23.0 mmol) in THF (20 ml), and the resulting mixture stirred for 30 min. A THF (20 ml) solution of 4-benzyloxybenzaldehyde (3.21 g, 15.1 mmol) was added, the resulting solution allowed to warm to room temperature, and stirring continued for 72 h. The mixture was carefully diluted with 1N HCl (150 ml), the products extracted into ethyl acetate (3 x 100 ml), and the combined organic phases washed with brine, dried (MgSO₄) and

(MgSO₄) and evaporated to give a yellow gum, which was purified by column chromatography on silica gel (10% ethyl acetate in *n*-heptane eluent) to give the intermediate, (*E/Z*)-ethyl 3-(4-benzyloxyphenyl)-2-isopropoxy-acrylate as a colourless gum.

The (*E/Z*)-ethyl 3-(4-benzyloxyphenyl)-2-isopropoxy-acrylate was dissolved in ethanol (100 ml), palladium on activated charcoal (10 wt. %, 0.80 g, 0.75 mmol) added and the mixture hydrogenated at 30 lb/in² H₂ pressure for 18 h. The catalyst was removed by filtration through celite and the solvent evaporated to give (*S/R*)-ethyl 3-(4-hydroxyphenyl)-2-isopropoxy-propionate (3.44 g, 90%) as a pale orange gum.


¹H NMR (300 MHz, CDCl₃) δ: 0.98 (3H, d), 1.15 (3H, d), 1.24 (3H, t), 2.82-2.98 (2H, m), 3.51 (1H, septet), 4.02 (1H, dd), 4.17 (2H, q), 5.49 (1H, br s), 6.75 (2H, dm), 7.09 (2H, dm). LCMS: 275 (M+Na), 253 (M+H), 235, 211, 193, 151, 137 (100%).

c)

The title compound (324 mg, 70%) was prepared as a colourless solid from (*S/R*)-ethyl 3-(4-hydroxyphenyl)-2-isopropoxy-propionate (280 mg, 1.11 mmol) and (*E*)-3-biphenyl-4-yl-but-2-en-1-ol (225 mg, 1.0 mmol) by a procedure analogous to that described in example 52c.

Mpt. 79-81°C. ¹H NMR (300 MHz, CDCl₃) δ: 0.97 (3H, d), 1.16 (3H, d), 1.24 (3H, t), 2.17 (3H, d), 2.85-3.02 (2H, m), 3.51 (1H, septet), 4.01 (1H, dd), 4.08-4.25 (2H, m), 4.75 (2H, d), 6.11 (1H, tm), 6.88 (2H, dm), 7.17 (2H, dm), 7.30-7.38 (1H, m), 7.40-7.65 (8H, m). LCMS: 665 (M+207), 481 (M+Na), 476 (M+NH₄), 207 (100%).

EXAMPLE 134




(*E*)-(S/R)-3-[4-(3-Biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-isopropoxy-propionic acid

The title compound was prepared from (*E*)-(S/R)-ethyl 3-[4-(3-biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-isopropoxy-propionate (example 133) (230 mg, 0.50 mmol) and sodium hydroxide

(1M, 1.5 ml, 1.5 mmol) by a procedure analogous to that described in example 51, yielding (E)-(S/R)-3-[4-(3-biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-isopropoxy-propionic acid (190 mg, 88%) as a colourless solid.

Mpt. 125-127.5°C. ¹H NMR (300 MHz, CDCl₃) δ: 1.03 (3H, d), 1.16 (3H, d), 2.17 (3H, d), 2.90 (1H, dd), 3.08 (1H, dd), 3.55 (1H, septet), 4.10 (1H, dd), 4.75 (2H, d), 6.12 (1H, tm), 6.90 (2H, dm), 7.17 (2H, dm), 7.30-7.38 (1H, m), 7.40-7.65 (8H, m), carboxylic acid proton not observed. LCMS: 637 (M+207), 453 (M+Na), 448 (M+NH₄), 207 (100%).

EXAMPLE 135



(E)-(S/R)-Ethyl 3-[4-(3-Biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-butoxy-propionate

a)

Butyl di-butoxy acetate (23.75 g, 91.2 mmol) was mixed with acetyl chloride (15.5 ml, 218 mmol) and iodine (0.2 g, 0.79 mmol) and the resulting brown solution heated to 60°C, under reflux, for 6 h. The product was then fractionally distilled, under reduced pressure, yielding (S/R)-butyl 2-butoxy-2-chloro-acetate (17.58 g, 79%) as an orange coloured oil.

Bpt. 130-135°C/approx. 15 mmHg. ¹H NMR (300 MHz, CDCl₃) δ: 0.89-0.99 (6H, m), 1.41 (4H, sextet), 1.60-1.75 (4H, m), 3.60 (1H, dt), 3.98 (1H, dt), 4.25 (2H, t), 5.81 (1H, s).

b)

A mixture of triethylphosphite (13.05 ml, 75.0 mmol) and (S/R)-butyl 2-butoxy-2-chloro-acetate (16.70 g, 75.0 mmol) was heated to 140°C, under reflux, for 6h. The resulting oil was fractionally distilled under reduced pressure to give the product, (S/R)-butyl 2-butoxy-2-(diethoxyphosphoryl)-acetate (20.42 g, 84%) as a colourless oil.

Bpt. 170-175°C/1-5 mmHg. ¹H NMR (300 MHz, CDCl₃) δ: 0.87-0.99 (6H, m), 1.30-1.49 (10H, m), 1.57-1.73 (4H, m), 3.52 (1H, dt), 3.66 (1H, dt), 4.15-4.30 (6H, m), 4.30 (1H, d, J_{HP} = 19 Hz). LCMS: 325 (100%, M+H), 269, 167.

5 c)

A THF (40 ml) solution of (S/R)-butyl 2-butoxy-2-(diethoxyphosphoryl)-acetate (14.60 g, 45.0 mmol) was added dropwise, at 0°C, to a stirred suspension of sodium hydride (55% dispersion in mineral oil, 2.61 g, 59.8 mmol) in THF (50 ml), and the resulting mixture stirred for 30 min. A THF (50 ml) solution of 4-benzyloxybenzaldehyde (6.37 g, 30.0 mmol) was added, the
10 resulting solution allowed to warm to room temperature, and stirring continued for 48 h. The mixture was carefully diluted with 1N HCl (200 ml), the products extracted into ethyl acetate (4 x 100 ml), and the combined organic phases washed with brine, dried (MgSO₄) and evaporated to give a yellow gum, which was purified by column chromatography on silica gel (10% ethyl acetate in *n*-heptane eluent) to give the intermediate, (E/Z)-butyl 3-(4-
15 benzyloxyphenyl)-2-butoxy-acrylate as a colourless gum.

The (E/Z)-butyl 3-(4-benzyloxyphenyl)-2-butoxy-acrylate was dissolved in ethanol (200 ml), palladium on activated charcoal (10 wt. %, 1.60 g, 1.5 mmol) added and the mixture hydrogenated at 30 lb/in² H₂ pressure for 18 h. The catalyst was removed by filtration through celite and the solvent evaporated to give an orange gum, which contained both (S/R)-butyl 3-(4-
20 (4-hydroxyphenyl)-2-butoxy-propionate and the trans-esterification product, (S/R)-ethyl 3-(4-hydroxyphenyl)-2-butoxy-propionate. These were separated by column chromatography on silica gel (15% ethyl acetate in *n*-heptane eluent) to give, in respective order of elution, (S/R)-butyl 3-(4-hydroxyphenyl)-2-butoxy-propionate (6.74 g, 76%) and (S/R)-ethyl 3-(4-hydroxyphenyl)-2-butoxy-propionate (0.40 g, 5%) as colourless oils.

25

(S/R)-butyl 3-(4-hydroxyphenyl)-2-butoxy-propionate: ¹H NMR (300 MHz, CDCl₃) δ: 0.85 (3H, t), 0.91 (3H, t), 1.22-1.43 (4H, m), 1.43-1.65 (4H, m), 2.89-2.98 (2H, m), 3.28 (1H, dt), 3.54 (1H, dt), 3.97 (1H, dd), 4.11 (2H, t), 5.56 (1H, br s), 6.74 (2H, dm), 7.08 (2H, dm). LCMS: 317 (M+Na), 295 (M+H), 221 (100%, M+H-BuOH), 193, 179, 165, 137.

30


(S/R)-ethyl 3-(4-hydroxyphenyl)-2-butoxy-propionate: ¹H NMR (300 MHz, CDCl₃) δ: 0.85 (3H, t), 1.23 (3H, t), 1.21-1.39 (2H, m), 1.43-1.60 (2H, m), 2.89-2.99 (2H, m), 3.28 (1H, dt), 3.55 (1H, dt), 3.97 (1H, dd), 4.17 (2H, q), 5.63 (1H, br s), 6.74 (2H, dm), 7.08 (2H, dm). LCMS: 289 (M+Na), 267 (M+H), 193 (100%, M+H-BuOH), 151, 137.

d)

The title compound (420 mg, 71%) was prepared as a colourless solid from (*S/R*)-ethyl 3-(4-hydroxyphenyl)-2-butoxy-propionate (385 mg, 1.45 mmol) and (*E*)-3-biphenyl-4-yl-but-2-en-1-ol (280 mg, 1.25 mmol) by a procedure analogous to that described in example 52c.

Mpt. 62-63.5°C. ¹H NMR (300 MHz, CDCl₃) δ: 0.85 (3H, t), 1.23 (3H, t), 1.20-1.40 (2H, m), 2.17 (3H, d), 2.90-3.00 (2H, m), 3.27 (1H, dt), 3.55 (1H, dt), 3.95 (1H, dd), 4.10-4.23 (2H, m), 4.74 (2H, d), 6.11 (1H, tm), 6.88 (2H, dm), 7.17 (2H, dm), 7.30-7.38 (1H, m), 7.40-7.63 (8H, m). LCMS: 679 (M+207), 495 (M+Na), 490 (M+NH₄), 207 (100%).


EXAMPLE 136



(*E*)-(S/R)-3-[4-(3-biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-butoxy-propionic acid

The title compound was prepared from (*E*)-(S/R)-ethyl 3-[4-(3-biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-butoxy-propionate (example 135) (331 mg, 0.70 mmol) and sodium hydroxide (1M, 2.1 ml, 2.1 mmol) by a procedure analogous to that described in example 51, yielding (*E*)-(S/R)-3-[4-(3-biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-butoxy-propionic acid (42 mg, 13%) as a colourless solid.

¹H NMR (300 MHz, CDCl₃) δ: 0.87 (3H, t), 1.21-1.38 (2H, m), 1.47-1.60 (2H, m), 2.17 (3H, br s), 2.96 (1H, dd), 3.09 (1H, dd), 3.33-3.44 (1H, m), 3.47-3.60 (1H, m), 4.04 (1H, dd), 4.75 (2H, d), 6.12 (1H, br t), 6.90 (2H, dm), 7.17 (2H, dm), 7.30-7.38 (1H, m), 7.38-7.65 (8H, m), carboxylic acid proton not observed. LCMS: 651 (M+207), 467 (100%, M+Na), 207.

EXAMPLE 137

5 (E)-(S/R)-Ethyl 3-[4-(3-Biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-hexyloxy-propionate

a)

(S/R)-Ethyl 2-(diethoxyphosphoryl)-2-hexyloxy-acetate was prepared as a pale green oil by the rhodium(II) acetate dimer catalysed reaction of ethyl diazo-(diethoxyphosphoryl)-acetate
10 with 1-hexanol, by a method analogous to that described for example 133a.

¹H NMR (300 MHz, CDCl₃) δ: 0.88 (3H, t), 1.23-1.44 (15H, m), 1.57-1.69 (2H, m), 3.51 (1H, dt), 3.65 (1H, dt), 4.15-4.38 (7H, m). LCMS: 671 (2M+Na), 649 (2M+H), 325 (100%, M+H), 297, 241.

15 b)

Sodium hydride (60% dispersion in mineral oil, 1.0 g, 25.0 mmol) was added at 0°C, in small portions, to a stirred THF (50 ml) solution of (S/R)-ethyl 2-(diethoxyphosphoryl)-2-hexyloxy-acetate (8.12 g, 25.0 mmol), and the resulting suspension stirred for 30 min. A THF (50 ml) solution of 4-benzyloxybenzaldehyde (4.25 g, 20.0 mmol) was added, the resulting solution
20 allowed to warm to room temperature, and stirring continued for 4 h. The mixture was carefully diluted with 0.5N HCl (150 ml), the products extracted into ethyl acetate (4 x 75 ml), and the combined organic phases washed with brine, dried (MgSO₄) and evaporated to give an orange coloured gum, which was purified by column chromatography on silica gel (15% ethyl acetate in n-heptane eluent) to give the intermediate, (E/Z)-ethyl 3-(4-benzyloxyphenyl)-2-hexyloxy-acrylate as a pale yellow oil.
25

The (E/Z)-ethyl 3-(4-benzyloxyphenyl)-2-hexyloxy-acrylate was dissolved in ethanol (150 ml), palladium on activated charcoal (10 wt. %, 1.40 g, 1.32 mmol) added and the mixture hydrogenated at 30 lb/in² H₂ pressure for 18 h. The catalyst was removed by filtration through celite and the solvent evaporated to give a colourless gum, which was purified by column
30 chromatography on silica gel (10% ethyl acetate in n-heptane eluent) to give (S/R)-ethyl 3-(4-hydroxyphenyl)-2-hexyloxy-propionate (1.83 g, 30%) as a colourless gum.


¹H NMR (300 MHz, CDCl₃) δ: 0.85 (3H, t), 1.14-1.32 (9H, m), 1.45-1.60 (2H, m), 2.94 (2H, d), 3.28 (1H, dt), 3.54 (1H, dt), 3.97 (1H, dd), 4.17 (2H, q), 5.95 (1H, br s), 6.74 (2H, dm), 7.07 (2H, dm). MS: 294 (M⁺), 221 (M-COOEt), 192 (M-hexanol), 137, 107 (100%).

5 c)

The title compound (248 mg, 81%) was prepared as a waxy solid from (*S/R*)-ethyl 3-(4-hydroxyphenyl)-2-hexyloxy-propionate (215 mg, 0.72 mmol) and (*E*)-3-biphenyl-4-yl-but-2-en-1-ol (137 mg, 0.61 mmol) by a procedure analogous to that described in example 52c.

¹H NMR (300 MHz, CDCl₃) δ: 0.86 (3H, t), 1.13-1.35 (9H, m), 1.46-1.60 (2H, m), 2.17 (3H, br s), 2.88-3.00 (2H, m), 3.26 (1H, dt), 3.55 (1H, dt), 3.95 (1H, dd), 4.10-4.23 (2H, m), 4.74 (2H, d), 6.12 (1H, tm), 6.88 (2H, dm), 7.17 (2H, dm), 7.30-7.37 (1H, m), 7.40-7.63 (8H, m).

EXAMPLE 138




(*E*)-(S/R)-3-[4-(3-biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-hexyloxy-propionic acid

20 The title compound was prepared from (*E*)-(S/R)-ethyl 3-[4-(3-biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-hexyloxy-propionate (example 137) (172 mg, 0.34 mmol) and sodium hydroxide (1M, 1.0 ml, 1.0 mmol) by a procedure analogous to that described in example 51, yielding (*E*)-(S/R)-3-[4-(3-biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-hexyloxy-propionic acid (160 mg, 99%) as a colourless solid.

25 Mpt. 117-119°C. ¹H NMR (300 MHz, CDCl₃) δ: 0.87 (3H, t), 1.13-1.38 (6H, m), 1.45-1.60 (2H, m), 2.17 (3H, d), 2.96 (1H, dd), 3.10 (1H, dd), 3.40 (1H, dt), 3.53 (1H, dt), 4.05 (1H, dd), 4.75 (2H, d), 6.12 (1H, tm), 6.90 (2H, dm), 7.16 (2H, dm), 7.30-7.38 (1H, m), 7.40-7.63 (8H, m), carboxylic acid proton not observed.

30

EXAMPLE 139

5 (E)-(S/R)-Ethyl 3-[4-(3-Biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-(3-phenyl-propoxy)-propionate

a)

(S/R)-Ethyl 2-(diethoxyphosphoryl)-2-(3-phenyl-propoxy)-acetate was prepared as a pale green oil by the rhodium(II) acetate \square immer catalysed reaction of 3-phenyl-1-propanol with
 10 ethyl diazo-(diethoxyphosphoryl)-acetate, by a method analogous to that described for example 133a.

^1H NMR (300 MHz, CDCl_3) δ : 1.23-1.40 (9H, m), 1.98 (2H, quintet), 2.72 (2H, t), 3.52 (1H, dt), 3.67 (1H, dt), 4.15-4.35 (7H, m), 7.12-7.22 (3H, m), 7.22-7.32 (2H, m).

15 b)

A THF (50 ml) solution of (S/R)-ethyl 2-(diethoxyphosphoryl)-2-(3-phenyl-propoxy)-acetate (14.2 g, 39.6 mmol) was added dropwise, at 0°C , to a stirred mixture of sodium hydride (60% dispersion in mineral oil, 2.35 g, 58.8 mmol) and 4-benzyloxybenzaldehyde (4.20 g, 19.8 mmol) in THF (50 ml), and the resulting mixture allowed to warm slowly to room temperature
 20 over 18 h. The mixture was carefully diluted with water (150 ml), the products extracted into ethyl acetate (2 x 150 ml), and the combined organic phases washed with brine, dried (MgSO_4) and evaporated to give (E/Z)-ethyl 3-(4-benzyloxyphenyl)-2-(3-phenyl-propoxy)-acrylate as a yellow oil.

The (E/Z)-ethyl 3-(4-benzyloxyphenyl)-2-(3-phenyl-propoxy)-acrylate was dissolved in ethanol (50 ml), palladium on activated charcoal (10 wt. %, 1.0 g, 0.94 mmol) added and the mixture hydrogenated at 30 lb/in 2 H_2 pressure for 18 h. The catalyst was removed by filtration through celite and the solvent evaporated to give a colourless oil.


25 ^1H NMR (300 MHz, CDCl_3) δ : 1.23 (3H, t), 1.72-1.97 (2H, m), 2.52-2.64 (2H, m), 2.87-3.02 (2H, m), 3.17-3.27 (1H, m), 3.53-3.63 (1H, m), 3.94 (1H, dd), 4.17 (2H, q), 4.93 (1H, s), 6.76
 30 (2H, dm), 7.02-7.29 (7H, m).

c)

The title compound (150 mg, 56%) was prepared as a waxy solid from (*S/R*)-ethyl 3-(4-hydroxyphenyl)-2-(3-phenyl-propoxy)-propionate (172 mg, 0.53 mmol) and (*E*)-3-biphenyl-4-yl-but-2-en-1-ol (112 mg, 0.50 mmol) by a procedure analogous to that described in example 52c.

¹H NMR (300 MHz, CDCl₃) δ: 1.23 (3H, t), 1.72-1.97 (2H, m), 2.16 (3H, br s), 2.51-2.65 (2H, m), 2.89-3.05 (2H, m), 3.17-3.27 (1H, m), 3.59 (1H, dt), 3.95 (1H, dd), 4.18 (2H, q), 4.74 (2H, d), 6.11 (1H, tm), 6.90 (2H, dm), 7.01-7.63 (16H, m). LCMS: 741 (M+207), 557 (100%, M+Na), 552 (M+NH₄), 207.


EXAMPLE 140



(*E*)-(S/R)-3-[4-(3-Biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-(3-phenyl-propoxy)-propionic acid

The title compound was prepared from (*E*)-(S/R)-ethyl 3-[4-(3-biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-(3-phenyl-propoxy)-propionate (example 139) (130 mg, 0.24 mmol) and sodium hydroxide (1M, 0.73 ml, 0.73 mmol) by a procedure analogous to that described in example 51, yielding (*E*)-(S/R)-3-[4-(3-biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-(3-phenyl-propoxy)-propionic acid (90 mg, 73%) as a colourless solid.

Mpt. 115-117°C. ¹H NMR (300 MHz, CDCl₃) δ: 1.75-1.98 (2H, m), 2.16 (3H, d), 2.52-2.68 (2H, m), 2.96 (1H, dd), 3.10 (1H, dd), 3.36 (1H, dt), 3.57 (1H, dt), 4.03 (1H, dd), 4.74 (2H, d), 6.11 (1H, tm), 6.91 (2H, dm), 7.06 (2H, dm), 7.12-7.63 (14H, m), carboxylic acid proton not observed. LCMS: 529 (M+Na), 525 (M+NH₄), 207 (100%).

EXAMPLE 141

5 (E)-(S/R)-Ethyl 3-[4-(3-Biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-(4-phenyl-butoxy)-propionate

a)

(S/R)-Ethyl 2-(diethoxyphosphoryl)-2-(4-phenyl-butoxy)-acetate was prepared as a pale green oil by the rhodium(II) acetate \square immer catalysed reaction of 4-phenyl-1-butanol with
10 ethyl diazo-(diethoxyphosphoryl)-acetate, by a method analogous to that described for example 133a.

^1H NMR (300 MHz, CDCl_3) δ : 1.25-1.40 (9H, m), 1.60-1.78 (4H, m), 2.63 (2H, t), 3.47-3.56 (1H, m), 3.61-3.70 (1H, m), 4.12-4.35 (7H, m), 7.12-7.21 (3H, m), 7.21-7.31 (2H, m). LCMS: 745 (2M+H), 373 (100%, M+H), 241.

15

b)

A THF (30 ml) solution of (S/R)-ethyl 2-(diethoxyphosphoryl)-2-(4-phenyl-butoxy)-acetate (15.64 g, 42.0 mmol) was added dropwise, at 0°C , to a stirred suspension of sodium hydride (60% dispersion in mineral oil, 2.52 g, 63.0 mmol) in THF (30 ml), and the resulting mixture
20 stirred for 20 min. A THF (50 ml) solution of 4-benzyloxybenzaldehyde (4.46 g, 21.0 mmol) was added, and the mixture warmed, resulting in a vigorous reaction. The mixture was cooled, carefully diluted with 0.5N HCl (150 ml), the products extracted into ethyl acetate (2 x 150 ml), and the combined organic phases washed with brine, dried (MgSO_4) and evaporated to give (E/Z)-ethyl 3-(4-benzyloxyphenyl)-2-(4-phenyl-butoxy)-acrylate as a yellow oil.
25 The (E/Z)-ethyl 3-(4-benzyloxyphenyl)-2-(4-phenyl-butoxy)-acrylate was dissolved in ethanol (175 ml), palladium on activated charcoal (10 wt. %, 0.50 g, 0.47 mmol) added and the mixture hydrogenated at 30 lb/in 2 H_2 pressure for 18 h. The catalyst was removed by filtration through celite and the solvent evaporated to give a colourless gum, which was purified by column chromatography on silica gel to give (S/R)-ethyl 3-(4-hydroxyphenyl)-2-(4-phenyl-
30 butoxy)-propionate (1.73 g, 24%) as a colourless oil.

¹H NMR (300 MHz, CDCl₃) δ: 1.22 (3H, t), 1.50-1.67 (4H, m), 2.50-2.60 (2H, m), 2.85-3.0 (2H, m), 3.21-3.31 (1H, m), 3.53-3.63 (1H, m), 3.94 (1H, dd), 4.16 (2H, q), 6.72 (2H, dm), 7.06-7.31 (7H, m), phenol proton not observed.


5 c)

The title compound (245 mg, 80%) was prepared as a yellow, waxy solid from (*S/R*)-ethyl 3-(4-hydroxyphenyl)-2-(4-phenyl-butoxy)-propionate (200 mg, 0.58 mmol) and (*E*)-3-biphenyl-4-yl-but-2-en-1-ol (125 mg, 0.56 mmol) by a procedure analogous to that described in example 52c.

10 ¹H NMR (300 MHz, CDCl₃) δ: 1.22 (3H, t), 1.50-1.68 (4H, m), 2.16 (3H, d), 2.50-2.60 (2H, m), 2.88-3.02 (2H, m), 3.21-3.33 (1H, m), 3.52-3.64 (1H, m), 3.95 (1H, dd), 4.17 (2H, q), 4.72 (2H, d), 6.11 (1H, tm), 6.87 (2H, dm), 7.06-7.63 (16H, m).

15

EXAMPLE 142



(*E*)-(*S/R*)-3-[4-(3-Biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-(4-phenyl-butoxy)-propionic acid


20

The title compound was prepared from (*E*)-(*S/R*)-ethyl 3-[4-(3-biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-(4-phenyl-butoxy)-propionate (example 141) (225 mg, 0.41 mmol) and sodium hydroxide (1M, 1.64 ml, 1.64 mmol) by a procedure analogous to that described in example 51, yielding (*E*)-(*S/R*)-3-[4-(3-biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-(4-phenyl-butoxy)-propionic acid (210 mg, 99%) as a pale yellow solid.

25

¹H NMR (300 MHz, CDCl₃) δ: 1.50-1.70 (4H, m), 2.17 (3H, d), 2.53-2.61 (2H, m), 2.95 (1H, dd), 3.09 (1H, dd), 3.34-3.44 (1H, m), 3.50-3.60 (1H, m), 4.04 (1H, dd), 4.72 (2H, d), 6.11 (1H, tm), 6.88 (2H, dm), 7.06-7.63 (16H, m), carboxylic acid proton not observed.

30

EXAMPLE 143

5 (E)-(S/R)-Propyl 3-[4-(3-Biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-propoxy-propionate

a)

Morpholinium dimorpholinoacetate was prepared according to the method described by Bourguignon and Wermuth (Bourguignon, J.J.; Wermuth, C.G. *J. Org. Chem.* **1981**, *46*, 4889-4894): an ethanol (100 ml) solution of morpholine (310 ml, 3.55 mol) was added, at 0°C, to a stirred ethanol (500 ml) solution of glyoxylic acid monohydrate (92.06 g, 1.0 mol) and the resulting mixture refrigerated for 60 h, a colourless precipitate being formed. The solid was collected by filtration, washed with diethyl ether (2 x 300 ml) and vacuum dried at 30°C to give morphilinium dimorpholinoacetate (298 g, 94%) as a colourless solid, which contained a small amount of water.

Mpt. 139-139.5°C. ¹H NMR (300 MHz, CDCl₃) δ: 2.83 (12H, br m), 3.26 (1H, s), 3.78 (12H, br m), 7.78 (2H, br s). Microanalysis Calculated % C: 52.93, H: 8.58, N: 13.24, water: 0.1%; found C: 52.84, H: 8.84, N: 13.15, water: 0.1%.

20 b)

Using a method based on that described by Kerfanto and Jegou, (Kerfanto, M.; Jegou, D. *Compt. Rendus.* **1965**, *261* (11), 2232-2233) morphilinium dimorpholinoacetate (127 g, 0.40 mol) was added to a stirred solution of hydrochloric acid (94.3 g, 6.5 mol) in 1-propanol (600 ml) and the resulting mixture heated to 80°C, under reflux, for 2 h. The resulting colourless suspension was filtered to remove morphiline hydrochloride, and the filtrate fractionally distilled, under reduced pressure, to give excess 1-propanol and the colourless oil, propyl 2,2-dipropoxyacetate (57.14 g, 65%).

¹H NMR (300 MHz, CDCl₃) δ: 0.86-1.05 (9H, m), 1.55-1.78 (6H, m), 3.47-3.65 (4H, m), 4.15 (2H, t), 4.89 (1H, s).

c)

A mixture of propyl 2,2-dipropoxyacetate (43.66 g, 0.20 mol), acetyl chloride (28 ml, 0.394 mol) and iodine (0.25 g, 1.0 mmol) was heated to 55°C, under reflux, for 16 h. Since GC analysis showed that some propyl 2,2-dipropoxyacetate starting material was still present, second portions of acetyl chloride (14 ml, 0.197 mol) and iodine (0.25 g, 1.0 mmol) were added, and heating continued for a further 6 h. The product was then purified by fractional distillation under reduced pressure to give (S/R)-propyl 2-chloro-2-propoxyacetate (32.67 g, 84%) as a pale orange oil (trace of iodine present).

Bpt. 116-119.5°C/approx. 10 mmHg. ¹H NMR (300 MHz, CDCl₃) δ: 0.97 (3H, t), 0.98 (3H, t), 1.63-1.82 (4H, m), 3.58 (1H, dt), 3.93 (1H, dt), 4.13-4.26 (2H, m), 5.83 (1H, s).

d)

Triethylphosphite (27 ml, 0.155 mol) was added to (S/R)-propyl 2-chloro-2-propoxyacetate (29.21 g, 0.15 mol), resulting in an immediate decolourisation of the pale orange acetate, and the resulting mixture heated to 140°C, under reflux, for 6 h, a colourless gas being evolved. The mixture was then fractionally distilled under reduced pressure to give the product, (S/R)-propyl 2-(diethoxyphosphoryl)-2-propoxyacetate (35.78 g, 80%) as a colourless oil.

Bpt. 155-160°C/approx. 3 mmHg. ¹H NMR (300 MHz, CDCl₃) δ: 0.95 (3H, t), 0.98 (3H, t), 1.31-1.39 (6H, m), 1.60-1.76 (4H, m), 3.49 (1H, dt), 3.62 (1H, dt), 4.12-4.30 (6H, m), 4.31 (1H, d, J_{HP} = 19 Hz).

e)

A THF (50 ml) solution of (S/R)-propyl 2-(diethoxyphosphoryl)-2-propoxyacetate (18.52 g, 62.5 mmol) was added dropwise, at 0°C, to a stirred suspension of sodium hydride (60% dispersion in mineral oil, 2.50 g, 62.5 mmol) in THF (50 ml), and the resulting mixture stirred for 30 min. A THF (100 ml) solution of 4-benzyloxybenzaldehyde (10.62 g, 50.0 mmol) was added, the resulting solution allowed to warm to room temperature, and stirring continued for 24 h. TLC showed a considerable amount of unreacted 4-benzyloxybenzaldehyde was still present so a further portion of sodium hydride (60% dispersion in mineral oil, 1.0 g, 25.0 mmol) was added and stirring continued for a further 18 h. The mixture was carefully diluted with 0.5N HCl (400 ml), the products extracted into ethyl acetate (3 x 200 ml), and the combined organic phases washed with brine, dried (MgSO₄) and evaporated to give a yellow gum, which was purified by column chromatography on silica gel (10% ethyl acetate in *n*-

heptane eluent) to give the intermediate, (*E/Z*)-propyl 3-(4-benzyloxyphenyl)-2-propoxy-acrylate as a colourless gum.


The (*E/Z*)-propyl 3-(4-benzyloxyphenyl)-2-propoxy-acrylate was dissolved in ethanol (200 ml), palladium on activated charcoal (10 wt. %, 2.18 g, 2.05 mmol) added and the mixture
5 hydrogenated at 30 lb/in² H₂ pressure for 20 h. The catalyst was removed by filtration through celite and the solvent evaporated to give an orange gum, which contained both the propyl and the ethyl (formed by trans-esterification) esters of (*S/R*)-3-(4-hydroxyphenyl)-2-propoxy-propionic acid. These were separated by column chromatography on silica gel (15% ethyl acetate in *n*-heptane eluent) to give, in respective order of elution, (*S/R*)-propyl 3-(4-
10 hydroxyphenyl)-2-propoxy-propionate (4.52 g, 41%) and a mixture of both (*S/R*)-propyl and (*S/R*)-ethyl 3-(4-hydroxyphenyl)-2-propoxy-propionates (4.98 g, approx. 45%) as colourless oils.

(*S/R*)-propyl 3-(4-hydroxyphenyl)-2-propoxy-propionate: ¹H NMR (300 MHz, CDCl₃) δ: 0.84 (3H, t), 0.90 (3H, t), 1.48-1.69 (4H, m), 2.95 (2H, d), 3.25 (1H, dt), 3.52 (1H, dt), 4.00 (1H, t),
15 4.07 (2H, t), 6.43 (1H, br s), 6.74 (2H, dm), 7.07 (2H, dm). ¹³C NMR (75 MHz, CDCl₃) δ: 10.2 (q), 10.3 (q), 21.8 (t), 22.7 (t), 38.4 (t), 66.6 (t), 72.5 (t), 80.6 (d), 115.2 (d), 128.5 (s), 130.4 (d), 154.7 (s), 173.2 (s). MS: 266 (M⁺), 206 (M-PrOH), 179, 164, 137, 107 (100%).

f)

20 The title compound (350 mg, 74%) was prepared as a colourless gum from (*S/R*)-propyl 3-(4-hydroxyphenyl)-2-propoxy-propionate (280 mg, 1.05 mmol) and (*E*)-3-biphenyl-4-yl-but-2-en-1-ol (224 mg, 1.0 mmol) by a procedure analogous to that described in example 52c.

¹H NMR (300 MHz, CDCl₃) δ: 0.85 (3H, t), 0.90 (3H, t), 1.49-1.69 (4H, m), 2.17 (3H, d), 3.91-3.02 (2H, m), 3.23 (1H, dt), 3.52 (1H, dt), 3.97 (1H, dd), 4.07 (2H, t), 4.74 (2H, d), 6.11 (1H,
25 tm), 6.88 (2H, dm), 7.17 (2H, dm), 7.30-7.38 (1H, m), 7.40-7.63 (8H, m). LCMS: 679 (M+207), 495 (100%, M+Na), 490 (M+NH₄), 207.

EXAMPLE 144


5 (E)-(S/R)-3-[4-(3-Biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-propoxy-propionic acid

The title compound was prepared from (E)-(S/R)-propyl 3-[4-(3-biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-propoxy-propionate (example 143) (330 mg, 0.70 mmol) and sodium hydroxide (1M, 1.4 ml, 1.4 mmol) by a procedure analogous to that described in example 51, yielding

10 (E)-(S/R)-3-[4-(3-biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-propoxy-propionic acid (300 mg, 100%) as a colourless gum.

¹H NMR (300 MHz, CDCl₃) δ: 0.88 (3H, t), 1.58 (2H, sextet), 2.17 (3H, s), 2.96 (1H, dd), 3.10 (1H, dd), 3.37 (1H, dt), 3.50 (1H, dt), 4.06 (1H, dd), 4.75 (2H, d), 6.11 (1H, t), 6.90 (2H, dm), 7.17 (2H, dm), 7.30-7.38 (1H, m), 7.40-7.63 (8H, m), carboxylic acid proton not observed.

15 LCMS: 637 (M+207), 453 (100%, M+Na), 207.

EXAMPLE 145


20 (E)-(S)-3-[4-[3-(3,5-Diethoxyoxy-phenyl)-allyloxy]-phenyl]-2-ethoxy-propionic acid ethyl ester

The title compound was prepared from 3,5-dihydroxybenzaldehyde (3.0 g, 22.0 mmol) and ethyl iodide (17.2 g, 110 mmol) by a sequence analogous to that described in example 75.

25

^1H NMR (300 MHz, CDCl_3) δ : 1.15 (t, 3H), 1.20 (t, 3H), 1.38 (t, 6H), 2.95 (d, 2H), 3.30-3.40 (m, 1H), 3.53-3.65 (m, 1H), 3.98 (q, 4H), 4.15 (q, 2H), 4.63 (d, 2H), 6.28-6.40 (m, 2H), 6.53 (d, 2H), 6.60 (d, 1H), 6.87 (d, 2H), 7.15 (d, 2H).


5

EXAMPLE 14610 (E)-(S)-3-{4-[3-(3,5-Diethoxyoxy-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid

The title compound was prepared from (E)-(S)-3-{4-[3-(3,5-diethoxyoxy-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester (730 mg, 1.6 mmol) by a procedure analogous to that described in example 26.

15 ^1H NMR (300 MHz, CDCl_3) δ : 1.15 (t, 3H), 1.22 (t, 3H), 1.38 (t, 6H), 2.95 (d, 2H), 3.28-3.38 (m, 1H), 3.53-3.65 (m, 1H), 3.98 (q, 4H), 4.15 (q, 2H), 4.63 (d, 2H), 6.28-6.40 (m, 2H), 6.53 (d, 2H), 6.60 (d, 1H), 6.85 (d, 2H), 7.15 (d, 2H).

20


EXAMPLE 147

(*E*)-(*S*)-3-{4-[3-(3,5-Bis-trifluoromethyl-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester

The title compound was prepared from 3,5-bis(trifluoromethyl)benzaldehyde (5.0 g, 20.7 mmol) by a sequence analogous to that described in example 23. The title compound was purified on HPLC, using ethyl acetate/heptane (20:80) as eluent.

¹H NMR (300 MHz, CDCl₃) δ: 1.15 (t, 3H), 1.22 (t, 3H), 2.97 (d, 2H), 3.30-3.42 (m, 1H), 3.55-3.67 (m, 1H), 3.98 (t, 1H), 4.18 (q, 2H), 4.72 (d, 2H), 6.55 (dt, 1H), 6.80 (d, 1H), 6.89 (d, 2H), 7.18 (d, 2H), 7.75 (bs, 1H), 7.82 (bs, 2H).


EXAMPLE 148



(*E*)-(*S*)-3-{4-[3-(3,5-Bis-trifluoromethyl-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid

The title compound was prepared from (*E*)-(*S*)-3-{4-[3-(3,5-bis-trifluoromethyl-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester (0.58 g, 1.2 mmol) by a procedure analogous to that described in example 26.


¹H NMR (300 MHz, CDCl₃) δ: 1.18 (t, 3H), 2.98 (dd, 1H), 3.08 (dd, 1H), 3.36-3.48 (m, 1H), 3.58-3.71 (m, 1H), 4.05 (dd, 1H), 4.72 (d, 2H), 6.55 (dt, 1H), 6.80 (d, 1H), 6.89 (d, 2H), 7.20 (d, 2H), 7.75 (bs, 1H), 7.82 (bs, 2H).

EXAMPLE 1495 (E)-(R,S)-3-[4-(3-Biphenyl-4-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid ethyl ester

The title compound was prepared from 3-biphenyl-4-yl-prop-2-en-1-ol (0.25 g, 0.001 mol) by a procedure analogous to that described in example 3c yielding 0.050 g of (E)-(R,S)-3-[4-(3-biphenyl-4-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid ethyl ester.

- 10 ^1H NMR (200 MHz, CDCl_3) δ : 1.1-1.26 (6H, m), 2.97 (2H, d), 3.3-3.4 (1H, m), 3.52-3.7 (1H, m), 4.0 (1H, t), 4.15 (2H, q), 4.75 (2H, dd), 6.35-6.5 (1H, dt), 6.75 (1H, d), 6.87 (2H, d), 7.15 (2H, d), 7.4-7.7 (9H, m).

15


EXAMPLE 150(E)-(R,S)-3-[4-(3-Biphenyl-4-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid

20

The title compound was prepared from (E)-(S)-3-[4-(3-biphenyl-4-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid ethyl ester (example 149) (0.040 g) by a procedure analogous to that described in example 2 yielding 0.0045 g of (E)-(R,S)-3-[4-(3-biphenyl-4-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid.

^1H NMR (300 MHz, CDCl_3) δ : 1.14 (3H, t), 2.85 (1H, dd), 3.1 (1H, dd) 3.42-3.57 (2H, m), 3.84-3.96 (2H, m), 4.1 (1H, dd), 4.7 (2H, d), 6.3-6.5 (1H, dt), 6.78 (1H, d), 6.88 (2H, d), 7.15 (2H, d) 7.4-7.6 (9H, m).

5

EXAMPLE 151

10

(*E*)-(S)- 4-(3-{3-[4-(2-Ethoxy-2-ethoxycarbonyl-ethyl)-phenoxy]-propenyl}-phenoxy)methyl-benzoic acid methyl ester

a)

15 (*E*)-3-(3-Hydroxy-propenyl)-phenol was prepared from 3-hydroxybenzaldehyde (6.0 g, 0.049 mol) by a procedure analogous to that described in example 1a-b yielding 1.5 g

^1H NMR (300 MHz, CDCl_3) δ : 1.4 (1H, t), 4.27 (2H, m), 4.88 (1H, s), 6.35 (1H, dt), 6.57 (1H, d), 6.68 (1H, dd), 6.87 (1H, s), 6.96 (1H, d), 7.19 (1H, dd).

20 b)

A mixture of (*E*)-3-(3-Hydroxy-propenyl)-phenol (0.5 g, 3.33 mmol), methyl 4-(bromomethyl) benzoate (763 mg, 3.33 mmol) and potassium carbonate (1.8 g, 13.3 mmol) in acetone (40 ml) was stirred at roomtemperature over night. The reaction mixtyre was added water (30ml) and acidified with 1N HCl and extracted with ethyl acetate (90 ml). The organic phase was

25 washed with water, brine and dried with sodium sulphate and evaporated and dried *in vacuo*

yielding 954 mg (96%) (*E*)-4-[3-(3-Hydroxy-propenyl)-phenoxy-methyl]-benzoic acid methyl ester.

¹H NMR (300MHz, CDCl₃) δ: 3.8 (3H, s), 4.24 (2H, d), 5.15 (2H, s), 6.3 (1H, dt), 6.57 (1H, d), 7.0 (2H, d), 7.2 (1H, d), 7.51 (2H, d), 8.08 (2H, d).

5

c)


The title compound was prepared from (*E*)-4-[3-(3-Hydroxy-propenyl)-phenoxy-methyl]-benzoic acid methyl ester (0.298 g, 1.0 mmol) by a procedure analogous to that described in example 3c yielding 0.184 g (35%) of (*E*)-(*S*)-4-(3-[3-[4-(2-Ethoxy-2-ethoxycarbonyl-ethyl)-phenoxy]-propenyl]-phenoxy-methyl)-benzoic acid methyl ester.

10

¹H NMR (300 MHz, CDCl₃) δ: 1.15-1.35 (6H, m), 2.9 (2H, d) 3.3-3.45 (1H, m), 3.53-3.68 (1H, m), 3.89 (3H, s), 3.97 (1H, t), 4.13 (2H, q), 4.68 (2H, dd), 5.15 (2H, s), 6.35 (1H, dt), 6.62 (1H, d), 6.87 (3H, d), 7.05 (2H, d), 7.13-7.3 (3H, m), 7.5 (2H, d), 8.10 (2H, d).

15

EXAMPLE 152




20 (*E*)-(*S*)-4-(3-[3-[4-(2-Carboxy-2-ethoxy-ethyl)-phenoxy]-propenyl]-phenoxy-methyl)-benzoic acid

The title compound was prepared from (*E*)-(*S*)-4-(3-[3-[4-(2-ethoxy-2-ethoxycarbonyl-ethyl)-phenoxy]-propenyl]-phenoxy-methyl)-benzoic acid methyl ester (example 151) (0.220 g) by a
25 procedure analogous to that described in example 2 yielding 0.160 g (77%) (*E*)-(*S*)-4-(3-[3-[4-(2-carboxy-2-ethoxy-ethyl)-phenoxy]-propenyl]-phenoxy-methyl)-benzoic acid.

¹H NMR (300 MHz, CDCl₃) δ: 1.17 (3H, t), 2.9-3.15 (2H, m) 3.3-3.68 (2H, m), 4.1 (2H, q), 4.67 (2H, d), 5.17 (2H, s), 6.35 (1H, dt), 6.68 (1H, d), 6.86 (3H, d), 7.05 (2H, d), 7.12-7.32 (3H, m), 7.52 (2H, d), 8.12 (2H, d).

5

EXAMPLE 153



10

(*E*)-(S)-Ethyl 2-Ethoxy-3-{4-[3-(4'-fluoro-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionate

a)

The colourless solid (*E*)-ethyl 3-(4'-fluoro-biphenyl-4-yl)-but-2-enoate was prepared from (*E*)-ethyl 3-(4-iodophenyl)-but-2-enoate (example 91a) and 4-fluorobenzene boronic acid by a procedure analogous to that described in example 52a.

Mpt. 63.5-64.5°C. ¹H NMR (300 MHz, CDCl₃) δ: 1.33 (3H, t), 2.61 (3H, d), 4.23 (2H, q), 6.20 (1H, m), 7.14 (2H, dd), 7.52-7.62 (6H, m). MS: 284 (100%, M⁺), 255, 239, 212, 196. Microanalysis Calculated % C: 76.04, H: 6.03; found C: 76.10, H: 6.17.

20

b)

The colourless solid (*E*)-3-(4'-fluoro-biphenyl-4-yl)-but-2-en-1-ol was prepared by DIBAL-H reduction of (*E*)-ethyl 3-(4'-fluoro-biphenyl-4-yl)-but-2-enoate as described for example 52b. Mpt. 120.5-122°C (n-heptane). ¹H NMR (300 MHz, CDCl₃) δ: 1.39 (1H, br s), 2.12 (3H, d), 4.40 (2H, d), 6.05 (1H, tm), 7.12 (2H, dd), 7.42-7.60 (6H, m). MS: 242 (100%, M⁺), 227 (M-Me), 224 (M-H₂O), 203, 199. Microanalysis Calculated % C: 79.32, H: 6.24; found C: 79.34, H: 6.37.

25


c)

The title compound (849mg, 89%) was prepared as a colourless gum from (*E*)-3-(4'-fluoro-biphenyl-4-yl)-but-2-en-1-ol (500 mg, 2.06 mmol) and (*S*)-ethyl 2-ethoxy-3-(4-hydroxyphenyl)-propionate (516 mg, 2.17 mmol) by a procedure analogous to that described in example 52c.

- 5 ¹H NMR (200 MHz, CDCl₃) δ: 1.17 (3H, t), 1.23 (3H, t), 2.17 (3H, d), 2.97 (2H, d), 3.27-3.44 (1H, m), 3.52-3.69 (1H, m), 3.98 (1H, t), 4.17 (2H, q), 4.75 (2H, d), 6.12 (1H, tm), 6.88 (2H, dm), 7.05-7.22 (4H, m), 7.44-7.62 (6H, m). LCMS: 687 (M+225), 641 (687-EtOH), 485 (M+Na), 480 (M+NH₄), 225 (100%).

10

EXAMPLE 154




- 15 (*E*)-(S)-2-Ethoxy-3-{4-[3-(4'-fluoro-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionic acid

The title compound was prepared from (*E*)-(S)-ethyl 2-ethoxy-3-{4-[3-(4'-fluoro-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionate (example 153) (463 mg, 1.0 mmol) and sodium hydroxide (1M, 1.5 ml, 1.5 mmol) by a procedure analogous to that described in example 51, yielding

20 (*E*)-(S)-2-ethoxy-3-{4-[3-(4'-fluoro-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionic acid (229 mg, 53%) as a colourless solid containing a trace of water.

- ¹H NMR (300 MHz, CDCl₃) δ: 1.19 (3H, t), 2.16 (3H, d), 2.97 (1H, dd), 3.09 (1H, dd), 3.42-3.65 (2H, m), 4.07 (1H, dd), 4.75 (2H, d), 6.11 (1H, tm), 6.90 (2H, dm), 7.07-7.20 (4H, m), 7.45-7.60 (6H, m), carboxylic acid proton not observed. LCMS: 457 (M+Na), 225 (100%). Microanalysis for C₂₇H₂₇FO₄•0.05H₂O Calculated % C: 74.48, H: 6.27, H₂O: 0.21; found C: 74.25, H: 6.39, H₂O: 0.21.
- 25

EXAMPLE 1555 (E)-(S)-Ethyl 2-Ethoxy-3-{4-[3-(4-Iodophenyl)-but-2-enyloxy]-phenyl}-propionate


The title compound (398 mg, 80%) was prepared as a colourless gum, from (*E*)-3-(4-iodophenyl)-but-2-en-1-ol (example 107a) (275 mg, 1.0 mmol) and (*S*)-ethyl 2-ethoxy-3-(4-hydroxyphenyl)-propionate (256 mg, 1.07 mmol) by a procedure analogous to that described
10 in example 52c.

¹H NMR (300 MHz, CDCl₃) δ: 1.17 (3H, t), 2.10 (3H, d), 2.96 (2H, d), 3.30-3.40 (1H, m), 3.55-3.65 (1H, m), 3.97 (1H, t), 4.16 (2H, q), 4.70 (2H, d), 6.04 (1H, tm), 6.86 (2H, dm), 7.13-7.20 (4H, m), 7.64 (2H, dm). LCMS: 751 (M+257), 705 (751-EtOH), 517 (100%, M+Na), 512 (M+NH₄), 449 (M+H-EtOH), 257, 130.

15

CLAIMS:

1. A compound of formula (I)



5

wherein A is aryl or heteroaryl and wherein A is optionally substituted with one or more substituents selected from hydroxy, halogen, perhalomethyl, perhalomethoxy, acyl, cyano, amino, C₁₋₆-alkylamino, C₁₋₆-dialkylamino, methylenedioxy, aralkenyl, aralkynyl, heteroaryloxy, heteroaralkoxy, aralkyl, heteroaralkyl, arylamino, or

10 A is optionally substituted with C₁₋₆-alkyl, C₂₋₆-alkenyl or C₂₋₆-alkynyl each of which is optionally substituted with one or more substituents selected from C₁₋₆-alkoxycarbonyl or carboxy, or

A is optionally substituted with C₁₋₆-alkoxy, C₁₋₆-alkylthio or C₂₋₆-alkenyloxy each of which is optionally substituted with one or more halogens, or

15 A is optionally substituted with aryloxy, arylthio or aralkoxy each of which is optionally substituted with one or more substituents selected from C₁₋₆-alkoxy, nitro, carboxy or C₁₋₆-alkoxycarbonyl; and

X₁ and X₂ independently are

20 hydrogen,

aryl or heteroaryl each of which is optionally substituted with one or more substituents selected from hydroxy, aryloxy, arylthio, aralkoxy, heteroaryloxy, aralkoxy, C₁₋₆-alkoxy, C₁₋₆-

alkylthio, halogen, perhalomethyl, perhalomethoxy, acyl, aryl, heteroaryl, aralkyl, heteroaralkyl, cyano, amino, C₁₋₆-alkylamino, C₁₋₆-dialkylamino, arylamino or methylenedioxy, or aryl or heteroaryl each of which is optionally substituted with one or more substituents selected from C₁₋₆-alkyl, C₂₋₆-alkenyl or C₂₋₆-alkynyl each of which is optionally substituted with

5 hydroxy; or

A is selected from the ring systems consisting of

10




Figure 1

15 wherein the attachment point of A to the remaining part of the structure of formula (I) is as indicated on the chemical structures in Figure 1, and wherein A is optionally substituted with one or more substituents selected from C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, hydroxy, aryloxy, arylthio, aralkoxy, C₁₋₆-alkoxy, C₁₋₆-alkylthio, heteroaryloxy, heteroaralkoxy, halogen, perhalomethyl, perhalomethoxy, acyl, aralkyl, heteroaralkyl, cyano, amino, C₁₋₆-alkylamino, C₁₋₆-dialkylamino, arylamino or methylenedioxy; and

20 wherein X₁ and X₂ are hydrogen; and

R₅ is hydrogen or C₁₋₆-alkyl; and

Y is hydrogen, or

Y is C₁₋₁₂-alkyl, C₂₋₁₂-alkenyl, C₂₋₁₂-alkynyl, C₄₋₁₂-alkenynyl, aralkyl or heteroaralkyl each of which is optionally substituted with one or more substituents selected from halogen, C₁₋₆-alkyl, perhalomethyl, hydroxy, aryl, heteroaryl, carboxy or amino; and

Z is hydrogen, halogen, hydroxy, or

Z is C₁₋₆-alkyl or C₁₋₆-alkoxy each of which is optionally substituted with one or more substituents selected from halogen, hydroxy, carboxy, amino, cyano or C₁₋₆-alkoxy; and

Q is O, S or NR₆, wherein R₆ is hydrogen, C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, C₄₋₆-alkenynyl, aralkyl, heteroaralkyl and wherein R₆ is optionally substituted with one or more substituents selected from halogen, hydroxy, C₁₋₆-alkoxy, amino or carboxy; and

Ar is arylene, heteroarylene or a divalent heterocyclic group each of which can be optionally substituted with one or more substituents selected from C₁₋₆-alkyl, aryl or C₁₋₆-alkoxy each of which can be optionally substituted with halogen, hydroxy, carboxy, cyano or heterocyclyl; and

R₁ is hydrogen, hydroxy or halogen; or R₁ forms a bond together with R₂; and

R₂ is hydrogen or C₁₋₆-alkyl; or R₂ forms a bond together with R₁; and

R₃ is hydrogen, or

R₃ is C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, C₄₋₆-alkenynyl, aryl, aralkyl, C₁₋₆-alkoxyC₁₋₆-alkyl, acyl, heterocyclyl, heteroaryl or heteroaralkyl each of which is optionally substituted with one or more substituents selected from halogen, perhalomethyl, hydroxy or cyano; and

R₄ is hydrogen, C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, C₄₋₆-alkenynyl or aryl; and


n is an integer ranging from 0 to 3; and

m is an integer ranging from 0 to 1;

or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate thereof, or any tautomeric forms, stereoisomers, mixture of stereoisomers including a racemic mixture, or polymorphs.

2. A compound according to claim 1 of formula (I)

5



wherein A is aryl or heteroaryl and wherein A is optionally substituted with one or more substituents selected from hydroxy, halogen, perhalomethyl, perhalomethoxy, acyl, cyano, amino, C₁₋₆-alkylamino, C₁₋₆-dialkylamino, methylenedioxy, aralkenyl, aralkynyl, heteroaryloxy, heteroaralkoxy, aralkyl, heteroaralkyl, arylamino, or

A is optionally substituted with C₁₋₆-alkyl, C₂₋₆-alkenyl or C₂₋₆-alkynyl each of which is optionally substituted with one or more substituents selected from C₁₋₆-alkoxycarbonyl or carboxy, or

A is optionally substituted with C₁₋₆-alkoxy, C₁₋₆-alkylthio or C₂₋₆-alkenyloxy each of which is optionally substituted with one or more halogens, or

A is optionally substituted with aryloxy, arylthio or aralkoxy each of which is optionally substituted with one or more substituents selected from C₁₋₆-alkoxy, nitro, carboxy or C₁₋₆-alkoxycarbonyl; and

X₁ and X₂ independently are hydrogen,

aryl or heteroaryl each of which is optionally substituted with one or more substituents selected from hydroxy, aryloxy, arylthio, aralkoxy, heteroaryloxy, aralkoxy, C₁₋₆-alkoxy, C₁₋₆-

alkylthio, halogen, perhalomethyl, perhalomethoxy, acyl, aryl, heteroaryl, aralkyl, heteroaralkyl, cyano, amino, C₁₋₆-alkylamino, C₁₋₆-dialkylamino, arylamino or methylenedioxy, or aryl or heteroaryl each of which is optionally substituted with one or more substituents selected from C₁₋₆-alkyl, C₂₋₆-alkenyl or C₂₋₆-alkynyl each of which is optionally substituted with hydroxy; or

A is selected from the ring systems consisting of




Figure 1

wherein the attachment point of A to the remaining part of the structure of formula (I) is as indicated on the chemical structures in Figure 1, and wherein A is optionally substituted with one or more substituents selected from C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, hydroxy, aryloxy, arylthio, aralkoxy, C₁₋₆-alkoxy, C₁₋₆-alkylthio, heteroaryloxy, heteroaralkoxy, halogen, perhalomethyl, perhalomethoxy, acyl, aralkyl, heteroaralkyl, cyano, amino, C₁₋₆-alkylamino, C₁₋₆-dialkylamino, arylamino or methylenedioxy; and wherein X₁ and X₂ are hydrogen; and

R₅ is hydrogen or C₁₋₆-alkyl; and

Y is hydrogen, or

Y is C₁₋₁₂-alkyl, C₂₋₁₂-alkenyl, C₂₋₁₂-alkynyl, C₄₋₁₂-alkenynyl, aralkyl or heteroaralkyl each of which is optionally substituted with one or more substituents selected from halogen, C₁₋₆-alkyl, perhalomethyl, hydroxy, aryl, heteroaryl, carboxy or amino; and

Z is hydrogen, halogen, hydroxy, or

Z is C₁₋₆-alkyl or C₁₋₆-alkoxy each of which is optionally substituted with one or more substituents selected from halogen, hydroxy, carboxy, amino, cyano or C₁₋₆-alkoxy; and

Q is O, S or NR₆, wherein R₆ is hydrogen, C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, C₄₋₆-alkenynyl, aralkyl, heteroaralkyl and wherein R₆ is optionally substituted with one or more substituents selected from halogen, hydroxy, C₁₋₆-alkoxy, amino or carboxy; and

Ar is arylene, heteroarylene or a divalent heterocyclic group each of which can be optionally substituted with one or more substituents selected from C₁₋₆-alkyl, aryl or C₁₋₆-alkoxy each of which can be optionally substituted with halogen, hydroxy, carboxy, cyano or heterocyclyl; and

R₁ is hydrogen, hydroxy or halogen; or R₁ forms a bond together with R₂; and

R₂ is hydrogen or C₁₋₆-alkyl; or R₂ forms a bond together with R₁; and

R₃ is hydrogen, or

R₃ is C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, C₄₋₆-alkenynyl, aryl, aralkyl, C₁₋₆-alkoxy, C₁₋₆-alkyl, acyl, heterocyclyl, heteroaryl or heteroaralkyl each of which is optionally substituted with one or more substituents selected from halogen, perhalomethyl, hydroxy or cyano; and


R₄ is hydrogen, C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, C₄₋₆-alkenynyl or aryl; and

n is an integer ranging from 1 to 3; and

m is 1;

or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate thereof, or any tautomeric forms, stereoisomers, mixture of stereoisomers including a racemic mixture, or polymorphs.

5 3. A compound according to claim 1 of formula (I)



wherein A is aryl or heteroaryl and wherein A is optionally substituted with one or more substituents selected from C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, hydroxy, aryloxy, arylthio, aralkoxy, C₁₋₆-alkoxy, C₁₋₆-alkylthio, heteroaryloxy, heteroaralkoxy, halogen, perhalomethyl, perhalomethoxy, acyl, aralkyl, heteroaralkyl, cyano, amino, C₁₋₆-alkylamino, C₁₋₆-dialkylamino, arylamino or methylenedioxy; and

X₁ and X₂ independently are hydrogen, aryl or heteroaryl optionally substituted with one or more substituents selected from C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, hydroxy, aryloxy, arylthio, aralkoxy, heteroaryloxy, aralkoxy, C₁₋₆-alkoxy, C₁₋₆-alkylthio, halogen, perhalomethyl, perhalomethoxy, acyl, aryl, heteroaryl, aralkyl, heteroaralkyl, cyano, amino, C₁₋₆-alkylamino, C₁₋₆-dialkylamino, arylamino or methylenedioxy;

or

wherein A is selected from the ring systems consisting of




Figure 1

5

wherein the attachment point of A to the remaining part of the structure of formula (I) is as indicated on the chemical structures in Figure 1, and
 wherein A is optionally substituted with one or more substituents selected from C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, hydroxy, aryloxy, arylthio, aralkoxy, C₁₋₆-alkoxy, C₁₋₆-alkylthio, heteroaryloxy, heteroaralkoxy, halogen, perhalomethyl, perhalomethoxy, acyl, aralkyl, heteroaralkyl, cyano, amino, C₁₋₆-alkylamino, C₁₋₆-dialkylamino, arylamino or methylenedioxy, and wherein X₁ and X₂ are hydrogen; and

15

R₅ is hydrogen or C₁₋₆-alkyl; and

20

Y is hydrogen, C₁₋₁₂-alkyl, C₂₋₁₂-alkenyl, C₂₋₁₂-alkynyl, C₄₋₁₂-alkenynyl, aralkyl or heteroaralkyl optionally substituted with one or more substituents selected from halogen, C₁₋₆-alkyl, perhalomethyl, hydroxy, aryl, heteroaryl, carboxy or amino; and

Z is hydrogen, halogen, hydroxy, C₁₋₆-alkyl or C₁₋₆-alkoxy optionally substituted with one or more substituents selected from halogen, hydroxy, carboxy, amino, cyano or C₁₋₆-alkoxy; and

Q is O, S or NR₆, wherein R₆ is hydrogen, C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, C₄₋₆-alkenynyl, aralkyl, heteroaralkyl and wherein R₆ is optionally substituted with one or more substituents selected from halogen, hydroxy, C₁₋₆-alkoxy, amino or carboxy; and

- 5 Ar is arylene, heteroarylene or a divalent heterocyclic group each of which can be optionally substituted with one or more substituents selected from C₁₋₆-alkyl, aryl or C₁₋₆-alkoxy each of which can be optionally substituted with halogen, hydroxy, carboxy, cyano or heterocyclyl; and

R₁ is hydrogen, hydroxy or halogen; or R₁ forms a bond together with R₂; and

10

R₂ is hydrogen or C₁₋₆-alkyl; or R₂ forms a bond together with R₁; and

15

R₃ is hydrogen, C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, C₄₋₆-alkenynyl, aryl, aralkyl, C₁₋₆-alkoxyC₁₋₆alkyl, acyl, heterocyclyl, heteroaryl or heteroaralkyl groups optionally substituted with one or more substituents selected from halogen, perhalomethyl, hydroxy or cyano; and

R₄ is hydrogen, C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, C₄₋₆-alkenynyl or aryl; and

20


n is an integer ranging from 0 to 3; and

m is an integer ranging from 0 to 1;

25


or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate thereof, or any tautomeric forms, stereoisomers, mixture of stereoisomers including a racemic mixture, or polymorphs.

4. A compound according to any one of the preceding claims of formula (I)



wherein A is aryl or heteroaryl and wherein A is optionally substituted with one or more substituents selected from C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, hydroxy, aryloxy, arylthio, aralkoxy, C₁₋₆-alkoxy, C₁₋₆-alkylthio, heteroaryloxy, heteroaralkoxy, halogen, perhalomethyl, perhalomethoxy, acyl, aralkyl, heteroaralkyl, cyano, amino, C₁₋₆-alkylamino, C₁₋₆-dialkylamino, arylamino or methylenedioxy; or

provided X_1 and X_2 is hydrogen, A is selected from the ring systems consisting of



wherein A is optionally substituted with one or more substituents selected from C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, hydroxy, aryloxy, arylthio, aralkoxy, C₁₋₆-alkoxy, C₁₋₆-alkylthio, heteroaryloxy, heteroaralkoxy, halogen, perhalomethyl, perhalomethoxy, acyl, aralkyl, heteroaralkyl, cyano, amino, C₁₋₆-alkylamino, C₁₋₆-dialkylamino, arylamino or methylenedioxy; and

R₅ is hydrogen or C₁₋₆-alkyl; and

10

X₁ and X₂ independently are hydrogen, aryl or heteroaryl optionally substituted with one or more substituents selected from C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, hydroxy, aryloxy, arylthio, aralkoxy, heteroaryloxy, aralkoxy, C₁₋₆-alkoxy, C₁₋₆-alkylthio, halogen, perhalomethyl, perhalomethoxy, acyl, aryl, heteroaryl, aralkyl, heteroaralkyl, cyano, amino, C₁₋₆-alkylamino, C₁₋₆-dialkylamino, arylamino or methylenedioxy; and

15

Y is hydrogen, C₁₋₁₂-alkyl, C₂₋₁₂-alkenyl, C₂₋₁₂-alkynyl, C₄₋₁₂-alkenynyl, aralkyl or heteroaralkyl optionally substituted with one or more substituents selected from halogen, C₁₋₆-alkyl, perhalomethyl, hydroxy, aryl, heteroaryl, carboxy or amino; and

20

Z is hydrogen, halogen, hydroxy, C₁₋₆-alkyl or C₁₋₆-alkoxy optionally substituted with one or more substituents selected from halogen, hydroxy, carboxy, amino, cyano or C₁₋₆-alkoxy; and

Q is O, S or NR₆, wherein R₆ is hydrogen, C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, C₄₋₆-alkenynyl, aralkyl, heteroaralkyl and wherein R₆ is optionally substituted with one or more substituents selected from halogen, hydroxy, C₁₋₆-alkoxy, amino or carboxy; and

- 5 Ar is arylene, heteroarylene or a divalent heterocyclic group each of which can be optionally substituted with one or more substituents selected from C₁₋₆-alkyl, aryl or C₁₋₆-alkoxy each of which can be optionally substituted with halogen, hydroxy, carboxy, cyano or heterocyclyl; and

R₁ is hydrogen, hydroxy or halogen; or R₁ forms a bond together with R₂; and

10

R₂ is hydrogen or C₁₋₆-alkyl; or R₂ forms a bond together with R₁; and

R₃ is hydrogen, C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, C₄₋₆-alkenynyl, aryl, aralkyl, C₁₋₆-alkoxyC₁₋₆-alkyl, acyl, heterocyclyl, heteroaryl or heteroaralkyl groups optionally substituted

- 15 with one or more substituents selected from halogen, perhalomethyl, hydroxy or cyano; and

R₄ is hydrogen, C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, C₄₋₆-alkenynyl or aryl; and

n is an integer ranging from 0 to 3; and

20

m is an integer ranging from 0 to 1;

or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate thereof, or any tautomeric forms, stereoisomers, mixture of stereoisomers including a racemic mixture, or polymorphs.

25


5. A compound according to any one of the preceding claims wherein A is aryl or heteroaryl optionally substituted with one or more substituents selected from C₁₋₆-alkyl, C₂₋₆-alkenyl each of which is optionally substituted with one or more substituents selected from C₁₋₆-alkoxycarbonyl or carboxy, or
- 30 A is optionally substituted with aryloxy optionally substituted with one or more C₁₋₆-alkoxy, or A is optionally substituted with aralkoxy optionally substituted with one or more substituents selected from C₁₋₆-alkoxy, nitro, carboxy or C₁₋₆-alkoxycarbonyl, or

A is optionally substituted with C₁₋₆-alkoxy optionally substituted with one or more halogens,
or

A is optionally substituted with aralkenyl, C₂₋₆-alkenyloxy, aralkynyl, halogen, perhalomethyl,
perhalomethoxy, acyl, aralkyl or methylenedioxy.

5

6. A compound according to any one of the preceding claims wherein A is aryl, heteroaryl or



10 optionally substituted with one or more substituents selected from aryloxy, arylthio, aralkoxy,
C₁₋₆-alkoxy, C₁₋₆-alkylthio, halogen, perhalomethyl, aralkyl, heteroaralkyl, heteroaryloxy or
heteroaralkoxy.

15 7. A compound according to any one of the preceding claims wherein A is aryl optionally sub-
stituted with one or more substituents selected from aryloxy, arylthio, aralkoxy, C₁₋₆-alkoxy,
C₁₋₆-alkylthio, halogen, perhalomethyl, aralkyl, heteroaralkyl, heteroaryloxy or heteroaralkoxy.

8. A compound according to any one of the preceding claims wherein A is aryl optionally sub-
stituted with one or more substituents selected from from C₁₋₆-alkyl, C₂₋₆-alkenyl each of
20 which is optionally substituted with one or more substituents selected from C₁₋₆-
alkoxycarbonyl or carboxy, or

A is optionally substituted with aryloxy optionally substituted with one or more C₁₋₆-alkoxy, or

A is optionally substituted with aralkoxy optionally substituted with one or more substituents
selected from C₁₋₆-alkoxy, nitro, carboxy or C₁₋₆-alkoxycarbonyl, or

25 A is optionally substituted with C₁₋₆-alkoxy optionally substituted with one or more halogens,
or

A is optionally substituted with aralkenyl, C₂₋₆-alkenyloxy, aralkynyl, halogen, perhalomethyl,
perhalomethoxy, acyl, aralkyl or methylenedioxy.

30 9. A compound according to any one of the preceding claims wherein A is aryl optionally sub-
stituted with one or more substituents selected from from C₁₋₆-alkyl, or

A is optionally substituted with aryloxy optionally substituted with one or more C₁₋₆-alkoxy,

A is optionally substituted with aralkoxy optionally substituted with one or more substituents selected from C₁₋₆-alkoxy, or


A is optionally substituted with C₁₋₆-alkoxy optionally substituted with one or more halogens, or

- 5 A is optionally substituted with aralkenyl, aralkynyl, halogen, perhalomethyl, perhalomethoxy or aralkyl.

10. A compound according to any one of the preceding claims wherein A is heteroaryl optionally substituted with one or more substituents selected from aryloxy, arylthio, aralkoxy, C₁₋₆-alkoxy, C₁₋₆-alkylthio, halogen, perhalomethyl, aralkyl, heteroaralkyl, heteroaryloxy or heteroaralkoxy.



11. A compound according to any one of the preceding claims wherein A is heteroaryl.

- 15 12. A compound according to any one of the preceding claims wherein A is



- optionally substituted with one or more substituents selected from aryloxy, arylthio, aralkoxy, C₁₋₆-alkoxy, C₁₋₆-alkylthio, halogen, perhalomethyl, aralkyl, heteroaralkyl, heteroaryloxy or heteroaralkoxy.

13. A compound according to any one of the preceding claims wherein A is



- 25 optionally substituted with one or more substituents selected from C₁₋₆-alkyl, and wherein R₅ is hydrogen or C₁₋₆-alkyl.

14. A compound according to any one of the preceding claims wherein X₁ and X₂ independently are hydrogen, aryl or heteroaryl optionally substituted with one or more substituents se-

lected from aryloxy, arylthio, aralkoxy, halogen, perhalomethyl, aryl, heteroaryl, aralkyl, heteroaralkyl, heteroaryloxy or heteroaralkoxy.

15. A compound according to any one of the preceding claims wherein X_1 and X_2 independently are
5 hydrogen,
aryl or heteroaryl optionally substituted with one or more substituents selected from halogen, acyl, aryl, or
aryl or heteroaryl optionally substituted with one or more substituents selected from C_{1-6} -
10 alkyl, C_{2-6} -alkynyl each of which is optionally substituted with hydroxy.

16. A compound according to any one of the preceding claims wherein X_1 and X_2 independently are
hydrogen, or
15 aryl optionally substituted with one or more substituents selected from halogen.

17. A compound according to any one of the preceding claims wherein X_1 and X_2 independently are hydrogen or aryl optionally substituted with one or more substituents selected from
aryloxy, arylthio, aralkoxy, halogen, perhalomethyl, aryl, heteroaryl, aralkyl, heteroaralkyl,
20 heteroaryloxy or heteroaralkoxy.

18. A compound according to any one of the preceding claims wherein X_1 and X_2 independently are
hydrogen,
25 aryl optionally substituted with one or more substituents selected from halogen, acyl, aryl, or
aryl optionally substituted with one or more substituents selected from C_{1-6} -alkyl, C_{2-6} -alkynyl
each of which is optionally substituted with hydroxy.

19. A compound according to any one of the preceding claims wherein X_1 and X_2 independently are
30 hydrogen, or
phenyl optionally substituted with one or more substituents selected from halogen.

20. A compound according to any one of the preceding claims wherein X_1 and X_2 independently are hydrogen or heteroaryl optionally substituted with one or more substituents selected from aryloxy, arylthio, aralkoxy, halogen, perhalomethyl, aryl, heteroaryl, aralkyl, heteroaralkyl, heteroaryloxy or heteroaralkoxy.
- 5
21. A compound according to any one of the preceding claims wherein X_1 and X_2 independently are hydrogen or heteroaryl.
22. A compound according to any one of the preceding claims wherein X_1 and X_2 is
- 10 hydrogen.
23. A compound according to any one of the preceding claims wherein Y is hydrogen or C_{1-12} -alkyl.
- 15
24. A compound according to any one of the preceding claims wherein Y is hydrogen or methyl.
25. A compound according to any one of the preceding claims wherein Z is hydrogen or C_{1-6} -alkoxy.
- 20
26. A compound according to any one of the preceding claims wherein Z is hydrogen or C_{1-6} -alkyl.
27. A compound according to any one of the preceding claims wherein Z is hydrogen.
- 25
28. A compound according to any one of the preceding claims wherein Q is O.
29. A compound according to any one of the preceding claims wherein Ar is arylene optionally substituted with one or more substituents selected from C_{1-6} -alkyl or C_{1-6} -alkoxy each of
- 30 which can be optionally substituted with carboxy.
30. A compound according to any one of the preceding claims wherein Ar is arylene.

31. A compound according to any one of the preceding claims wherein R_1 is hydrogen or R_1 forms a bond together with R_2 .
32. A compound according to any one of the preceding claims wherein R_1 is hydrogen.
- 5 33. A compound according to any one of the preceding claims wherein R_2 is hydrogen or R_2 forms a bond together with R_1 .
34. A compound according to any one of the preceding claims wherein R_2 is hydrogen.
- 10 35. A compound according to any one of the preceding claims wherein R_3 is C_{1-6} -alkyl or aralkyl.
36. A compound according to any one of the preceding claims wherein R_3 is C_{1-6} -alkyl.
- 15 37. A compound according to any one of the preceding claims wherein R_4 is hydrogen, C_{1-3} -alkyl.
38. A compound according to any one of the preceding claims wherein R_4 is hydrogen.
- 20 39. A compound according to any one of the preceding claims wherein n is 1.
40. A compound according to any one of the preceding claims wherein m is 1.
- 25 41. A compound according to any one of the preceding claims wherein alkyl is methyl, ethyl, n-propyl, iso-propyl, butyl, tert-butyl, pentyl, hexyl, cyclopropyl or cyclopentyl.
42. A compound according to any one of the preceding claims wherein alkenyl is vinyl or 1-propenyl.
- 30 43. A compound according to any one of the preceding claims wherein alkynyl is ethynyl, 1-propynyl and 2-propynyl.

44. A compound according to any one of the preceding claims wherein alkoxy is methoxy, ethoxy, propoxy, butoxy, pentoxy, hexoxy, isopropoxy or cyclopentyloxy.
- 5 45. A compound according to any one of the preceding claims wherein alkylthio is methylthio, ethylthio, propylthio or cyclopentylthio.
46. A compound according to any one of the preceding claims wherein aryl is phenyl optionally substituted with halogen.
- 10 47. A compound according to any one of the preceding claims wherein arylene is phenylene optionally substituted with halogen.
48. A compound according to any one of the preceding claims wherein halogen is fluorine or chlorine.
- 15 49. A compound according to any one of the preceding claims wherein perhalomethyl is trifluoromethyl.
50. A compound according to any one of the preceding claims wherein acyl is acetyl.
- 20 51. A compound according to any one of the preceding claims wherein heteroaryl is furan, thiophene, pyrrole, imidazole, pyrazole, pyridine, quinoline, isoquinoline, quinazoline, quinoxaline, indole, benzimidazole or benzofuran.
- 25 52. A compound according to any one of the preceding claims wherein heteroaryl is furan, pyrrole, indole or benzofuran.
53. A compound according to any one of the preceding claims wherein heteroarylene is furan, thiophene, pyrrole, imidazole, pyrazole, triazole, pyridine, pyrazine, pyrimidine or pyridazine.
- 30 54. A compound according to any one of the preceding claims wherein aralkyl is benzyl or phenethyl.
55. A compound according to any one of the preceding claims wherein aryloxy is phenoxy.

56. A compound according to any one of the preceding claims wherein aralkoxy is benzyloxy.
57. A compound according to any one of the preceding claims wherein n is an integer
5 ranging from 1 to 3 and m is 1.
58. A compound according to any of the preceding claims wherein the substituents Z and Y are arranged in a trans-configuration.
59. A compound according to any of the preceding claims wherein the substituents Z and Y are arranged in a cis-configuration.
- 10 60. The compound according to claim 1, 2 or 3 which is:
- (E)-(S)-Ethyl 3-[4-(3-biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-ethoxy-propionate,
(E)-(S)-3-[4-(3-Biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-ethoxy-propionic acid,
15 (E)-(S)-3-[4-[3-(4'-Bromo-biphenyl-4-yl)-but-2-enyloxy]-phenyl]-2-ethoxy-propionic acid ethyl ester,
(E)-(S)-3-[4-[3-(4'-Bromo-biphenyl-4-yl)-but-2-enyloxy]-phenyl]-2-ethoxy-propionic acid,
(E)-(S)-2-Ethoxy-3-[4-[3-(4-phenoxy-phenyl)-but-2-enyloxy]-phenyl]-propionic acid ethyl ester,
20 (E)-(S)-2-Ethoxy-3-[4-[3-(4-phenoxy-phenyl)-but-2-enyloxy]-phenyl]-propionic acid,
(E)-(S)-2-Ethoxy-3-[4-[3-[4-(4-methoxy-phenoxy)-phenyl]-but-2-enyloxy]-phenyl]-propionic acid ethyl ester,
(E)-(S)-2-Ethoxy-3-[4-[3-[4-(4-methoxy-phenoxy)-phenyl]-but-2-enyloxy]-phenyl]-propionic acid,
25 (E)-(S)-2-Ethoxy-3-[4-[3-(9H-fluoren-2-yl)-but-2-enyloxy]-phenyl]-propionic acid ethyl ester,
(E)-(S)-2-Ethoxy-3-[4-[3-(9H-fluoren-2-yl)-but-2-enyloxy]-phenyl]-propionic acid,
(E)-(S)-3-[4-[3-(3,4-Dimethoxy-phenyl)-but-2-enyloxy]-phenyl]-2-ethoxy-propionic acid ethyl ester,
(E)-(S)-3-[4-[3-(3,4-Dimethoxy-phenyl)-but-2-enyloxy]-phenyl]-2-ethoxy-propionic acid,
30 (E)-(S)-3-[4-[3-(3,5-Bis-trifluoromethyl-phenyl)-but-2-enyloxy]-phenyl]-2-ethoxy-propionic acid ethyl ester,
(E)-(S)-3-[4-[3-(3,5-Bis-trifluoromethyl-phenyl)-but-2-enyloxy]-phenyl]-2-ethoxy-propionic acid,

- (E)-(S)-3-[4-(3-Biphenyl-4-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid ethyl ester,
(E)-(S)-3-[4-(3-Biphenyl-4-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid,
(E)-(S)-2-Ethoxy-3-[4-(3-naphthalen-2-yl-but-2-enyloxy)-phenyl]-propionic acid ethyl ester,
(E)-(S)-2-Ethoxy-3-[4-(3-naphthalen-2-yl-but-2-enyloxy)-phenyl]-propionic acid,
5 (E)-(S)-2-Ethoxy-3-[4-(3-pyridin-2-yl-but-2-enyloxy)-phenyl]-propionic acid ethyl ester,
(E)-(S)-2-Ethoxy-3-[4-(3-pyridin-2-yl-but-2-enyloxy)-phenyl]-propionic acid,
(E)-(S)-3-[4-[3-(3,5-Bis-benzyloxy-phenyl)-but-2-enyloxy]-phenyl]-2-ethoxy-propionic acid
ethyl ester,
(E)-(S)-3-[4-[3-(3,5-Bis-benzyloxy-phenyl)-but-2-enyloxy]-phenyl]-2-ethoxy-propionic acid,
10 (E)-(S)-2-Ethoxy-3-[4-(3-naphthalen-2-yl-allyloxy)-phenyl]-propionic acid ethyl ester,
(E)-(S)-2-Ethoxy-3-[4-(3-naphthalen-2-yl-allyloxy)-phenyl]-propionic acid,
(E)-(S)-2-Ethoxy-3-[4-[3-(3-phenoxy-phenyl)-allyloxy]-phenyl]-propionic acid ethyl ester,
(E)-(S)-2-Ethoxy-3-[4-[3-(3-phenoxy-phenyl)-allyloxy]-phenyl]-propionic acid,
(S)-3-[4-(2-Benzofuran-3-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid ethyl ester,
15 (S)-3-[4-(2-Benzofuran-3-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid;

or a pharmaceutically acceptable salt thereof.

61. The compound according to claim 4 which is:

20

- (E)-(S)-Ethyl 3-[4-(3-biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-ethoxy-propionate,
(E)-(S)-3-[4-(3-Biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-ethoxy-propionic acid,
(E)-(S)-3-[4-[3-(4'-Bromo-biphenyl-4-yl)-but-2-enyloxy]-phenyl]-2-ethoxy-propionic acid ethyl
ester,
25 (E)-(S)-3-[4-[3-(4'-Bromo-biphenyl-4-yl)-but-2-enyloxy]-phenyl]-2-ethoxy-propionic acid,
(E)-(S)-2-Ethoxy-3-[4-[3-(4-phenoxy-phenyl)-but-2-enyloxy]-phenyl]-propionic acid ethyl es-
ter,
(E)-(S)-2-Ethoxy-3-[4-[3-(4-phenoxy-phenyl)-but-2-enyloxy]-phenyl]-propionic acid,
(E)-(S)-2-Ethoxy-3-[4-[3-[4-(4-methoxy-phenoxy)-phenyl]-but-2-enyloxy]-phenyl]-propionic
acid ethyl ester,
30 (E)-(S)-2-Ethoxy-3-[4-[3-[4-(4-methoxy-phenoxy)-phenyl]-but-2-enyloxy]-phenyl]-propionic
acid,
(E)-(S)-2-Ethoxy-3-[4-[3-(9H-fluoren-2-yl)-but-2-enyloxy]-phenyl]-propionic acid ethyl ester,
(E)-(S)-2-Ethoxy-3-[4-[3-(9H-fluoren-2-yl)-but-2-enyloxy]-phenyl]-propionic acid,

(E)-(S)-3-{4-[3-(3,4-Dimethoxy-phenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester,

(E)-(S)-3-{4-[3-(3,4-Dimethoxy-phenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid,

(E)-(S)-3-{4-[3-(3,5-Bis-trifluoromethyl-phenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid
5 ethyl ester,

(E)-(S)-3-{4-[3-(3,5-Bis-trifluoromethyl-phenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid,

(E)-(S)-3-{4-(3-Biphenyl-4-yl-allyloxy)-phenyl}-2-ethoxy-propionic acid ethyl ester,

(E)-(S)-3-{4-(3-Biphenyl-4-yl-allyloxy)-phenyl}-2-ethoxy-propionic acid,

10 (E)-(S)-2-Ethoxy-3-{4-(3-naphthalen-2-yl-but-2-enyloxy)-phenyl}-propionic acid ethyl ester,

(E)-(S)-2-Ethoxy-3-{4-(3-naphthalen-2-yl-but-2-enyloxy)-phenyl}-propionic acid,

(E)-(S)-2-Ethoxy-3-{4-(3-pyridin-2-yl-but-2-enyloxy)-phenyl}-propionic acid ethyl ester,

(E)-(S)-2-Ethoxy-3-{4-(3-pyridin-2-yl-but-2-enyloxy)-phenyl}-propionic acid,

(E)-(S)-3-{4-[3-(3,5-Bis-benzyloxy-phenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid
15 ethyl ester,

(E)-(S)-3-{4-[3-(3,5-Bis-benzyloxy-phenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid,

(E)-(S)-2-Ethoxy-3-{4-(3-naphthalen-2-yl-allyloxy)-phenyl}-propionic acid ethyl ester,

(E)-(S)-2-Ethoxy-3-{4-(3-naphthalen-2-yl-allyloxy)-phenyl}-propionic acid,

(E)-(S)-2-Ethoxy-3-{4-[3-(3-phenoxy-phenyl)-allyloxy]-phenyl}-propionic acid ethyl ester,

20 (E)-(S)-2-Ethoxy-3-{4-[3-(3-phenoxy-phenyl)-allyloxy]-phenyl}-propionic acid,

(S)-3-{4-(2-Benzofuran-3-yl-allyloxy)-phenyl}-2-ethoxy-propionic acid ethyl ester,

(S)-3-{4-(2-Benzofuran-3-yl-allyloxy)-phenyl}-2-ethoxy-propionic acid;

or a pharmaceutically acceptable salt thereof.

25

62. The compound according to claim 1, 2 or 3 which is:

(E)-(S)-3-{4-[3-(4-Benzyloxy-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester,

(E)-(S)-3-{4-[3-(4-Benzyloxy-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid,

30 (E)-(S)-3-{4-(3-Benzo[1,3]dioxol-5-yl-allyloxy)-phenyl}-2-ethoxy-propionic acid ethyl ester,

(E)-(S)-3-{4-(3-Benzo[1,3]dioxol-5-yl-allyloxy)-phenyl}-2-ethoxy-propionic acid,

(E)-(S)-3-{4-[3-(4-Allyloxy-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester,

(E)-(S)-3-{4-[3-(4-Allyloxy-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid,

(E)-(S)-3-{4-(3-Benzofuran-7-yl-allyloxy)-phenyl}-2-ethoxy-propionic acid ethyl ester,

- (E)-(S)-3-[4-(3-Benzofuran-7-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid,
(S)-3-[4-(3-Benzo[1,3]dioxol-4-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid ethyl ester,
(S)-3-[4-(3-Benzo[1,3]dioxol-4-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid,
(E)-(S)-2-Ethoxy-3-[4-[3-(9H-fluoren-2-yl)-allyloxy]-phenyl]-propionic acid ethyl ester,
5 (E)-(S)-2-Ethoxy-3-[4-[3-(9H-fluoren-2-yl)-allyloxy]-phenyl]-propionic acid,
(S)-2-Ethoxy-3-[4-(3-quinolin-2-yl-allyloxy)-phenyl]-propionic acid ethyl ester,
(S)-2-Ethoxy-3-[4-(3-quinolin-2-yl-allyloxy)-phenyl]-propionic acid,
(E)-(S)-3-[4-[3-(3,5-Bis-benzyloxy-phenyl)-allyloxy]-phenyl]-2-ethoxy-propionic acid ethyl ester,
10 (E)-(S)-3-[4-[3-(3,5-Bis-benzyloxy-phenyl)-allyloxy]-phenyl]-2-ethoxy-propionic acid,
(E)-(S)-3-[4-[3-(3,5-Dimethoxy-phenyl)-allyloxy]-phenyl]-2-ethoxy-propionic acid ethyl ester,
(E)-(S)-3-[4-[3-(3,5-Dimethoxy-phenyl)-allyloxy]-phenyl]-2-ethoxy-propionic acid,
(E)-(S)-2-Ethoxy-3-[4-(3-phenanthren-9-yl-allyloxy)-phenyl]-propionic acid ethyl ester,
(E)-(S)-2-Ethoxy-3-[4-[3-(2-methoxy-naphthalen-1-yl)-allyloxy]-phenyl]-propionic acid ethyl
15 ester,
(E)-(S)-2-Ethoxy-3-[4-[3-(2-methoxy-naphthalen-1-yl)-allyloxy]-phenyl]-propionic acid,
(E)-(S)-Ethyl 3-[4-[3-(4-Bromophenyl)-but-2-enyloxy]-phenyl]-2-ethoxy-propionate,
(E)-(S)-3-[4-[3-(4-Bromophenyl)-but-2-enyloxy]-phenyl]-2-ethoxy-propionic acid,
(E)-(S)-Ethyl 3-[4-[3-(4'-Chloro-biphenyl-4-yl)-but-2-enyloxy]-phenyl]-2-ethoxy-propionate,
20 (E)-(S)-3-[4-[3-(4'-Chloro-biphenyl-4-yl)-but-2-enyloxy]-phenyl]-2-ethoxy-propionic acid,
(E)-(S)-Ethyl 2-Ethoxy-3-[4-[3-(5'-isopropyl-2'-methoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl]-
propionate,
(E)-(S)-2-Ethoxy-3-[4-[3-(5'-isopropyl-2'-methoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl]-
propionic acid,
25 (E)-(S)-Ethyl 2-Ethoxy-3-[4-[3-(5'-chloro-2'-methoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl]-
propionate,
(E)-(S)-3-[4-[3-(5'-Chloro-2'-methoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl]-2-ethoxy-
propionic acid,
(E)-(S)-Ethyl 3-[4-[3-(2',3'-Dichloro-biphenyl-4-yl)-but-2-enyloxy]-phenyl]-2-ethoxy-
30 propionate,
(E)-(S)-3-[4-[3-(2',3'-Dichloro-biphenyl-4-yl)-but-2-enyloxy]-phenyl]-2-ethoxy-propionic acid,
(E)-(S)-Ethyl 3-[4-[3-(2',6'-Dimethoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl]-2-ethoxy-
propionate,

(E)-(S)-3-{4-[3-(2',6'-Dimethoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid,

(E)-(S)-Ethyl 3-{4-[3-(4-Bromophenyl)-2-methyl-but-2-enyloxy]-phenyl}-2-ethoxy-propionate,

(E)-(S)-3-{4-[3-(4-Bromophenyl)-2-methyl-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid,

5 (Z)-(S)-Ethyl 3-{4-[3-(4-Bromophenyl)-2-methyl-but-2-enyloxy]-phenyl}-2-ethoxy-propionate,

(Z)-(S)-3-{4-[3-(4-Bromophenyl)-2-methyl-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid,

(E)-(S)-Ethyl 2-Ethoxy-3-[4-(3-[1,1';3',1'']terphenyl-4''-yl)-but-2-enyloxy]-phenyl]-propionate,

(E)-(S)-2-Ethoxy-3-[4-(3-[1,1';3',1'']terphenyl-4''-yl)-but-2-enyloxy]-phenyl]-propionic acid,

(E)-(S)-Ethyl 2-Ethoxy-3-[4-[3-(3'-methyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl]-propionate,

10 (E)-(S)-2-Ethoxy-3-[4-[3-(3'-methyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl]-propionic acid,

(E)-(S)-3-[4-[3-(3'-Acetyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl]-2-ethoxy-propionate,

(E)-(S)-2-Ethoxy-3-[4-[3-(5'-isopropyl-2'-methoxy-biphenyl-4-yl)-allyloxy]-phenyl]-propionic acid ethyl ester,

15 (E)-(S)-2-Ethoxy-3-[4-[3-(5'-isopropyl-2'-methoxy-biphenyl-4-yl)-allyloxy]-phenyl]-propionic acid,

(E)-(S)-3-[4-[3-(3,5-Distyryl-phenyl)-allyloxy]-phenyl]-2-ethoxy-propionic acid ethyl ester,

(E)-(S)-3-[4-[3-(3,5-Distyryl-phenyl)-allyloxy]-phenyl]-2-ethoxy-propionic acid;

or a pharmaceutically acceptable salt thereof.

20

63. The compound according to claim 4 which is:

(E)-(S)-3-[4-[3-(4-Benzoyloxy-phenyl)-allyloxy]-phenyl]-2-ethoxy-propionic acid ethyl ester,

(E)-(S)-3-[4-[3-(4-Benzoyloxy-phenyl)-allyloxy]-phenyl]-2-ethoxy-propionic acid,

25 (E)-(S)-3-[4-(3-Benzo[1,3]dioxol-5-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid ethyl ester,

(E)-(S)-3-[4-(3-Benzo[1,3]dioxol-5-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid,

(E)-(S)-3-[4-[3-(4-Allyloxy-phenyl)-allyloxy]-phenyl]-2-ethoxy-propionic acid ethyl ester,

(E)-(S)-3-[4-[3-(4-Allyloxy-phenyl)-allyloxy]-phenyl]-2-ethoxy-propionic acid,

(E)-(S)-3-[4-(3-Benzofuran-7-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid ethyl ester,

30 (E)-(S)-3-[4-(3-Benzofuran-7-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid,

(S)-3-[4-(3-Benzo[1,3]dioxol-4-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid ethyl ester,

(S)-3-[4-(3-Benzo[1,3]dioxol-4-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid,

(E)-(S)-2-Ethoxy-3-[4-[3-(9H-fluoren-2-yl)-allyloxy]-phenyl]-propionic acid ethyl ester,

(E)-(S)-2-Ethoxy-3-[4-[3-(9H-fluoren-2-yl)-allyloxy]-phenyl]-propionic acid,

- (S)-2-Ethoxy-3-[4-(3-quinolin-2-yl-allyloxy)-phenyl]-propionic acid ethyl ester,
(S)-2-Ethoxy-3-[4-(3-quinolin-2-yl-allyloxy)-phenyl]-propionic acid,
(E)-(S)-3-[4-[3-(3,5-Bis-benzyloxy-phenyl)-allyloxy]-phenyl]-2-ethoxy-propionic acid ethyl ester,
5 (E)-(S)-3-[4-[3-(3,5-Bis-benzyloxy-phenyl)-allyloxy]-phenyl]-2-ethoxy-propionic acid,
(E)-(S)-3-[4-[3-(3,5-Dimethoxy-phenyl)-allyloxy]-phenyl]-2-ethoxy-propionic acid ethyl ester,
(E)-(S)-3-[4-[3-(3,5-Dimethoxy-phenyl)-allyloxy]-phenyl]-2-ethoxy-propionic acid,
(E)-(S)-2-Ethoxy-3-[4-(3-phenanthren-9-yl-allyloxy)-phenyl]-propionic acid ethyl ester,
(E)-(S)-2-Ethoxy-3-[4-[3-(2-methoxy-naphthalen-1-yl)-allyloxy]-phenyl]-propionic acid ethyl
10 ester,
(E)-(S)-2-Ethoxy-3-[4-[3-(2-methoxy-naphthalen-1-yl)-allyloxy]-phenyl]-propionic acid,
(E)-(S)-Ethyl 3-[4-[3-(4-Bromophenyl)-but-2-enyloxy]-phenyl]-2-ethoxy-propionate,
(E)-(S)-3-[4-[3-(4-Bromophenyl)-but-2-enyloxy]-phenyl]-2-ethoxy-propionic acid,
(E)-(S)-Ethyl 3-[4-[3-(4'-Chloro-biphenyl-4-yl)-but-2-enyloxy]-phenyl]-2-ethoxy-propionate,
15 (E)-(S)-3-[4-[3-(4'-Chloro-biphenyl-4-yl)-but-2-enyloxy]-phenyl]-2-ethoxy-propionic acid,
(E)-(S)-Ethyl 2-Ethoxy-3-[4-[3-(5'-isopropyl-2'-methoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl]-
propionate,
(E)-(S)-2-Ethoxy-3-[4-[3-(5'-isopropyl-2'-methoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl]-
propionic acid,
20 (E)-(S)-Ethyl 2-Ethoxy-3-[4-[3-(5'-chloro-2'-methoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl]-
propionate,
(E)-(S)-3-[4-[3-(5'-Chloro-2'-methoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl]-2-ethoxy-
propionic acid,
(E)-(S)-Ethyl 3-[4-[3-(2',3'-Dichloro-biphenyl-4-yl)-but-2-enyloxy]-phenyl]-2-ethoxy-
25 propionate,
(E)-(S)-3-[4-[3-(2',3'-Dichloro-biphenyl-4-yl)-but-2-enyloxy]-phenyl]-2-ethoxy-propionic acid,
(E)-(S)-Ethyl 3-[4-[3-(2',6'-Dimethoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl]-2-ethoxy-
propionate,
(E)-(S)-3-[4-[3-(2',6'-Dimethoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl]-2-ethoxy-propionic
30 acid,
(E)-(S)-Ethyl 3-[4-[3-(4-Bromophenyl)-2-methyl-but-2-enyloxy]-phenyl]-2-ethoxy-propionate,
(E)-(S)-3-[4-[3-(4-Bromophenyl)-2-methyl-but-2-enyloxy]-phenyl]-2-ethoxy-propionic acid,
(Z)-(S)-Ethyl 3-[4-[3-(4-Bromophenyl)-2-methyl-but-2-enyloxy]-phenyl]-2-ethoxy-propionate,
(Z)-(S)-3-[4-[3-(4-Bromophenyl)-2-methyl-but-2-enyloxy]-phenyl]-2-ethoxy-propionic acid,

- (E)-(S)-Ethyl 2-Ethoxy-3-[4-(3-[1,1';3',1'']terphenyl-4''-yl-but-2-enyloxy)-phenyl]-propionate,
(E)-(S)-2-Ethoxy-3-[4-(3-[1,1';3',1'']terphenyl-4''-yl-but-2-enyloxy)-phenyl]-propionic acid,
(E)-(S)-Ethyl 2-Ethoxy-3-[4-[3-(3'-methyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl]-propionate,
(E)-(S)-2-Ethoxy-3-[4-[3-(3'-methyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl]-propionic acid,
5 (E)-(S)-3-[4-[3-(3'-Acetyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl]-2-ethoxy-propionate,
(E)-(S)-2-Ethoxy-3-[4-[3-(5'-isopropyl-2'-methoxy-biphenyl-4-yl)-allyloxy]-phenyl]-propionic
acid ethyl ester,
(E)-(S)-2-Ethoxy-3-[4-[3-(5'-isopropyl-2'-methoxy-biphenyl-4-yl)-allyloxy]-phenyl]-propionic
acid,
10 (E)-(S)-3-[4-[3-(3,5-Distyryl-phenyl)-allyloxy]-phenyl]-2-ethoxy-propionic acid ethyl ester,
(E)-(S)-3-[4-[3-(3,5-Distyryl-phenyl)-allyloxy]-phenyl]-2-ethoxy-propionic acid;

or a pharmaceutically acceptable salt thereof.

- 15 64. The compound according to claim 1, 2, 3 or 4 which is:

- (E)-(S)-3-[4-[3-(3,5-Diisopropoxy-phenyl)-allyloxy]-phenyl]-2-ethoxy-propionic acid ethyl es-
ter,
20 (E)-(S)-3-[4-[3-(3,5-Diisopropoxy-phenyl)-allyloxy]-phenyl]-2-ethoxy-propionic acid,
(S)-3-[4-[3-(3-Bromo-5-styryl-phenyl)-allyloxy]-phenyl]-2-ethoxy-propionic acid ethyl ester,
(S)-3-[4-[3-(3-Bromo-5-styryl-phenyl)-allyloxy]-phenyl]-2-ethoxy-propionic acid,
25 (E)-(S)-2-Ethoxy-3-[4-(3-phenyl-allyloxy)-phenyl]-propionic acid ethyl ester,
(E)-(S)-2-Ethoxy-3-[4-(3-phenyl-allyloxy)-phenyl]-propionic acid,
(E)-(S)-3-[4-[3-(2',3'-Dichloro-biphenyl-4-yl)-allyloxy]-phenyl]-2-ethoxy-propionic acid ethyl
30 ester,
(E)-(S)-3-[4-[3-(2',3'-Dichloro-biphenyl-4-yl)-allyloxy]-phenyl]-2-ethoxy-propionic acid,

- (E)-(S)-3-{4-[3-(3,5-Bis-phenylethynyl-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester,
- (E)-(S)-3-{4-[3-(3,5-Bis-phenylethynyl-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid,
- 5 (E)-(S)-3-{4-[3-(3,5-Diphenethyl-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester,
- (E)-(S)-3-{4-[3-(3,5-Diphenethyl-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid,
- 10 3-{4-[3-(3,5-Bis-cyclopentyloxy-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester,
- (E)-(S)-3-{4-[3-(3,5-Bis-cyclopentyloxy-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid,
- (E)-(S)-3-{4-[3-[3,5-Bis-(2,2,2-trifluoro-ethoxy)-phenyl]-allyloxy]-phenyl}-2-ethoxy-propionic
- 15 acid ethyl ester,
- (E)-(S)-3-{4-[3-[3,5-Bis-(2,2,2-trifluoro-ethoxy)-phenyl]-allyloxy]-phenyl}-2-ethoxy-propionic acid,
- 20 (E)-(S)-Ethyl 2-Ethoxy-3-{4-[3-(4-furan-2-yl-phenyl)-but-2-enyloxy]-phenyl}-propionate,
- (E)-(S)-Ethyl 2-Ethoxy-3-{4-[3-(2'-methyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionate,
- (E)-(S)-2-Ethoxy-3-{4-[3-(2'-methyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionic acid,
- 25 (E)-(S)-Ethyl 3-{4-[3-(2',5'-Dimethoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate,
- (E)-(S)-3-{4-[3-(2',5'-Dimethoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic
- 30 acid,
- (E)-(S)-Ethyl 3-{4-[3-(4-Bromophenyl)-2-ethyl-but-2-enyloxy]-phenyl}-2-ethoxy-propionate,
- (E)-(S)-3-{4-[3-(4-Bromophenyl)-2-ethyl-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid,

(Z)-(S)-Ethyl 3-{4-[3-(4-Bromophenyl)-2-ethyl-but-2-enyloxy]-phenyl}-2-ethoxy-propionate,

(Z)-(S)-3-{4-[3-(4-Bromophenyl)-2-ethyl-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid,

5

(E)-(S)-Ethyl 3-{4-[3-(4'-*tert*-Butyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate,

(E)-(S)-Ethyl 3-{4-[3-(3',5'-Bis-trifluoromethyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate,

10

(E)-(S)-3-{4-[3-(3',5'-Bis-trifluoromethyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid,

(E)-(S)-Ethyl 2-Ethoxy-3-{4-[3-(4'-isopropyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionate,

15

(E)-(S)-2-Ethoxy-3-{4-[3-(4'-isopropyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionic acid,

(E)-(S)-Ethyl 3-{4-[3-(3'-Acetyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate,

20 (E)-(S)-3-{4-[3-(3,5-Dimethoxy-phenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester,

(E)-(S)-Ethyl 3-{4-[3-(4'-Acetyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate,

25 (E)-(S)-3-{4-[3-(4'-Acetyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid,

(E)-(S)-Ethyl 2-Ethoxy-3-[4-(3-[1,1';3',1'']terphenyl-5'-yl-allyloxy)-phenyl]-propionate,

(E)-(S)-2-Ethoxy-3-[4-(3-[1,1';3',1'']terphenyl-5'-yl-allyloxy)-phenyl]-propionic acid,

30

(E,E)-(S)-Ethyl 3-(4'-{3-[4-(2-Ethoxy-2-ethoxycarbonyl-ethyl)-phenoxy]-1-methyl-propenyl}-biphenyl-3-yl)-but-2-enoate,

(*E,E*)-(S)- 3-(4'-{3-[4-(2-Carboxy-2-ethoxy-ethyl)-phenoxy]-1-methyl-propenyl}-biphenyl-3-yl)-but-2-enoic acid,

(*E*)-(S)-Ethyl 2-Ethoxy-3-{4-[3-(3'-methoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionate,

5

(*E*)-(S)-2-Ethoxy-3-{4-[3-(3'-methoxy-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionic acid,

(*E*)-(S,S/*R*)-Ethyl 2-Ethoxy-3-(4-{3-[3'-(1-hydroxy-ethyl)-biphenyl-4-yl]-but-2-enyloxy}-phenyl)-propionate,

10

(*E*)-(S,S/*R*)-2-Ethoxy-3-(4-{3-[3'-(1-hydroxy-ethyl)-biphenyl-4-yl]-but-2-enyloxy}-phenyl)-propionic acid,

(*E*)-(S)-Ethyl 3-{4-[3-(3,5-Dibromophenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate,

15

(*E*)-(S)-3-{4-[3-(3,5-Dibromophenyl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid,

(*E*)-(S)-Ethyl 3-{4-[3-(3,5-Dibromophenyl)-allyloxy]-phenyl}-2-ethoxy-propionate,

20 (*E*)-(S)-3-{4-[3-(3,5-Dibromophenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid,

(*E*)-(S)-Ethyl 3-{4-[3-(4,4''-Di-*tert*-butyl-[1,1';3',1'']terphenyl-5'-yl)-allyloxy]-phenyl}-2-ethoxy-propionate,

25 (*E*)-(S)-3-{4-[3-(4,4''-Di-*tert*-butyl-[1,1';3',1'']terphenyl-5'-yl)-allyloxy]-phenyl}-2-ethoxy-propionic acid,

(*E*)-(S)-Ethyl 3-{4-[3-(3',5'-Dibromo-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate,

30

(*E*)-(S)-3-{4-[3-(3',5'-Dibromo-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid,

(*E*)-(S)-Ethyl 3-{4-[3-(3',5'-Dichloro-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate,

(E)-(S)-3-{4-[3-(3',5'-Dichloro-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid,

(E)-(S)-Ethyl 3-{4-[3-(3',5'-Dichloro-biphenyl-4-yl)-allyloxy]-phenyl}-2-ethoxy-propionate,

5

(E)-(S)-3-{4-[3-(3',5'-Dichloro-biphenyl-4-yl)-allyloxy]-phenyl}-2-ethoxy-propionic acid,

(E)-(S)-Ethyl 3-{4-[3-(3',5'-di-*tert*-butyl-biphenyl-4-yl)-allyloxy]-phenyl}-2-ethoxy-propionate,

10 (E)-(S)-3-{4-[3-(3',5'-Di-*tert*-butyl-biphenyl-4-yl)-allyloxy]-phenyl}-2-ethoxy-propionic acid,

(E)-(S)-Ethyl 3-{4-[3-(3',5'-Di-*tert*-butyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionate,

15 (E)-(S)-3-{4-[3-(3',5'-Di-*tert*-butyl-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-2-ethoxy-propionic acid,

(E)-(S/R)-Ethyl 3-[4-(3-Biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-isopropoxy-propionate,

20 (E)-(S/R)-3-[4-(3-Biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-isopropoxy-propionic acid,

(E)-(S/R)-Ethyl 3-[4-(3-Biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-butoxy-propionate,

(E)-(S/R)-3-[4-(3-Biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-butoxy-propionic acid,

25

(E)-(S/R)-Ethyl 3-[4-(3-Biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-hexyloxy-propionate,

(E)-(S/R)-3-[4-(3-Biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-hexyloxy-propionic acid,

30 (E)-(S/R)-Ethyl 3-[4-(3-Biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-(3-phenyl-propoxy)-propionate,

(E)-(S/R)-3-[4-(3-Biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-(3-phenyl-propoxy)-propionic acid,

(E)-(S/R)-Ethyl 3-[4-(3-Biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-(4-phenyl-butoxy)-propionate,

- (E)-(S/R)-3-[4-(3-Biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-(4-phenyl-butoxy)-propionic acid,
- (E)-(S/R)-Propyl 3-[4-(3-Biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-propoxy-propionate,
- 5 (E)-(S/R)-3-[4-(3-Biphenyl-4-yl-but-2-enyloxy)-phenyl]-2-propoxy-propionic acid,
- (E)-(S)-3-{4-[3-(3,5-Diethoxyoxy-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester,
- (E)-(S)-3-{4-[3-(3,5-Diethoxyoxy-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid,
- 10 (E)-(S)-3-{4-[3-(3,5-Bis-trifluoromethyl-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid ethyl ester,
- (E)-(S)-3-{4-[3-(3,5-Bis-trifluoromethyl-phenyl)-allyloxy]-phenyl}-2-ethoxy-propionic acid,
- (E)-(R,S)-3-[4-(3-Biphenyl-4-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid ethyl ester,
- (E)-(R,S)-3-[4-(3-Biphenyl-4-yl-allyloxy)-phenyl]-2-ethoxy-propionic acid,
- 15 (E)-(S)- 4-(3-{3-[4-(2-Ethoxy-2-ethoxycarbonyl-ethyl)-phenoxy]-propenyl}-phenoxy-methyl)-benzoic acid methyl ester,
- (E)-(S)- 4-(3-{3-[4-(2-Carboxy-2-ethoxy-ethyl)-phenoxy]-propenyl}-phenoxy-methyl)-benzoic acid;
- 20 or a pharmaceutically acceptable salt thereof.
65. The compound according to claim 1, 2, 3 or 4 which is:
- (E)-(S)-Ethyl 2-Ethoxy-3-{4-[3-(4'-fluoro-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionate,
- 25 (E)-(S)-2-Ethoxy-3-{4-[3-(4'-fluoro-biphenyl-4-yl)-but-2-enyloxy]-phenyl}-propionic acid,
- (E)-(S)-Ethyl 2-Ethoxy-3-{4-[3-(4-Iodophenyl)-but-2-enyloxy]-phenyl}-propionate;
- or a pharmaceutically acceptable salt thereof.
- 30 66. A pharmaceutical composition comprising, as an active ingredient, a compound according to any one of the preceding compound claims or a pharmaceutically acceptable salt thereof together with a pharmaceutically acceptable carrier or diluent.

67. A composition according to claim 66 in unit dosage form, comprising from about 0.05 to about 100 mg, preferably from about 0.1 to about 50 mg of the compound according to any one of the preceding compound claims or a pharmaceutically acceptable salt thereof.

5 68. A pharmaceutical composition useful in the treatment and/or prevention of conditions mediated by nuclear receptors, in particular the Peroxisome Proliferator-Activated Receptors (PPAR), the composition comprising, as an active ingredient, a compound according to any one of the preceding compound claims or a pharmaceutically acceptable salt thereof together with a pharmaceutically acceptable carrier or diluent.

10

69. A pharmaceutical composition useful in the treatment and/or prevention of diabetes and/or obesity, the composition comprising, as an active ingredient, a compound according to any one of the preceding compound claims or a pharmaceutically acceptable salt thereof together with a pharmaceutically acceptable carrier or diluent.

15

70. A pharmaceutical composition according to any one of the claims 66-69 for oral, nasal, transdermal, pulmonal, or parenteral administration.

20 71. A method for the treatment of ailments, the method comprising administering to a subject in need thereof an effective amount of a compound according to any one of the preceding compound claims or a pharmaceutically acceptable salt thereof, or of a composition according to any one of the preceding claims 66-70.

25 72. A method for the treatment and/or prevention of conditions mediated by nuclear receptors, in particular the Peroxisome Proliferator-Activated Receptors (PPAR), the method comprising administering to a subject in need thereof an effective amount of a compound according to any one of the preceding compound claims or a pharmaceutically acceptable salt thereof, or of a composition according to any one of the preceding claims 66-70.

30 73. A method for the treatment and/or prevention of diabetes and/or obesity, the method comprising administering to a subject in need thereof an effective amount of a compound according to any one of the preceding compound claims or a pharmaceutically acceptable salt thereof, or of a composition according to any one of the preceding claims 66-70.

74. The method according to claims 71, 72 or 73, wherein the effective amount of the compound according to any one of the preceding compound claims or a pharmaceutically acceptable salt or ester thereof is in the range of from about 0.05 to about 100 mg per day, preferably from about 0.1 to about 50 mg per day.

5

75. Use of a compound according to any one of the preceding compound claims or a pharmaceutically acceptable salt thereof for the preparation of a medicament.

76. Use of a compound according to any one of the preceding compound claims or a pharmaceutically acceptable salt thereof for the preparation of a medicament useful in the treatment and/or prevention of conditions mediated by nuclear receptors, in particular the Peroxisome Proliferator-Activated Receptors (PPAR).

77. Use of a compound according to any one of the preceding compound claims or a pharmaceutically acceptable salt thereof for the preparation of a medicament for treatment and/or prevention of diabetes and/or obesity.

78. A pharmaceutical composition suitable for treating type I diabetes, type II diabetes, impaired glucose tolerance, insulin resistance or obesity comprising a compound according to any of the claims 1 to 65 or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate thereof, or any tautomeric forms, stereoisomers, mixture of stereoisomers including a racemic mixture, or polymorphs together with one or more pharmaceutically acceptable carriers or diluents and an ACE (angiotensin converting enzyme) inhibitor.

79. The use of a compound according to any one of the claims 1 to 65 or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate thereof, or any tautomeric forms, stereoisomers, mixture of stereoisomers including a racemic mixture, or polymorphs together with one or more pharmaceutically acceptable carriers or diluents and an ACE (angiotensin converting enzyme) inhibitor for the preparation of a medicament suitable for the treatment of type I diabetes, type II diabetes, impaired glucose tolerance, insulin resistance or obesity.

80. A method of treating type I diabetes, type II diabetes, impaired glucose tolerance, insulin resistance or obesity comprising administering to a subject in need thereof an effective

amount of a compound according to any of the claims 1 to 65 or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate thereof, or any tautomeric forms, stereoisomers, mixture of stereoisomers including a racemic mixture, or polymorphs together with one or more pharmaceutically acceptable carriers or diluents and an ACE
5 (angiotensin converting enzyme) inhibitor to said subject.

81. A pharmaceutical composition suitable for treating type I diabetes, type II diabetes, impaired glucose tolerance, insulin resistance or obesity comprising a compound according to any of the claims 1 to 65 or a pharmaceutically acceptable salt thereof, or a pharmaceutically
10 acceptable solvate thereof, or any tautomeric forms, stereoisomers, mixture of stereoisomers including a racemic mixture, or polymorphs together with one or more pharmaceutically acceptable carriers or diluents and an agent stimulating insulin release from β cells such as a meglitinide, like repaglinide or senaglinide.

82. The use of a compound according to any one of the claims 1 to 65 or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate thereof, or any tautomeric forms, stereoisomers, mixture of stereoisomers including a racemic mixture, or polymorphs together with one or more pharmaceutically acceptable carriers or diluents and an agent
15 stimulating insulin release from β cells such as a meglitinide, like repaglinide or senaglinide, for the preparation of a medicament suitable for the treatment of type I diabetes, type II
20 diabetes, impaired glucose tolerance, insulin resistance or obesity.

83. A method of treating type I diabetes, type II diabetes, impaired glucose tolerance, insulin resistance or obesity comprising administering to a subject in need thereof an effective
25 amount of a compound according to any of the claims 1 to 65 and an agent stimulating insulin release from β cells such as a meglitinide, like repaglinide or senaglinide, to said subject.

84. A pharmaceutical composition suitable for treating type I diabetes, type II diabetes, impaired glucose tolerance, insulin resistance or obesity comprising a compound according to any of the claims 1 to 65 or a pharmaceutically acceptable salt thereof, or a pharmaceutically
30 acceptable solvate thereof, or any tautomeric forms, stereoisomers, mixture of stereoisomers including a racemic mixture, or polymorphs together with one or more pharmaceutically acceptable carriers or diluents and a biguanide like metformin.

85. The use of a compound according to any one of the claims 1 to 65 or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate thereof, or any tautomeric forms, stereoisomers, mixture of stereoisomers including a racemic mixture, or polymorphs together with one or more pharmaceutically acceptable carriers or diluents and a biguanide, like metformin, for the preparation of a medicament suitable for the treatment of type I diabetes, type II diabetes, impaired glucose tolerance, insulin resistance or obesity.

86. A method of treating type I diabetes, type II diabetes, impaired glucose tolerance, insulin resistance or obesity comprising administering to a subject in need thereof an effective amount of a compound according to any of the claims 1 to 65 or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate thereof, or any tautomeric forms, stereoisomers, mixture of stereoisomers including a racemic mixture, or polymorphs together with one or more pharmaceutically acceptable carriers or diluents and a biguanide, like metformin, to said subject.

87. A pharmaceutical composition suitable for treating type I diabetes, type II diabetes, impaired glucose tolerance, insulin resistance or obesity comprising a compound according to any of the claims 1 to 65 or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate thereof, or any tautomeric forms, stereoisomers, mixture of stereoisomers including a racemic mixture, or polymorphs together with one or more pharmaceutically acceptable carriers or diluents and a HMG CoA inhibitor.

88. The use of a compound according to any one of the claims 1 to 65 or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate thereof, or any tautomeric forms, stereoisomers, mixture of stereoisomers including a racemic mixture, or polymorphs together with one or more pharmaceutically acceptable carriers or diluents and a HMG CoA inhibitor for the preparation of a medicament suitable for the treatment of type I diabetes, type II diabetes, impaired glucose tolerance, insulin resistance or obesity.

89. A method of treating type I diabetes, type II diabetes, impaired glucose tolerance, insulin resistance or obesity comprising administering to a subject in need thereof an effective amount of a compound according to any of the claims 1 to 65 or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate thereof, or any tautomeric

forms, stereoisomers, mixture of stereoisomers including a racemic mixture, or polymorphs together with one or more pharmaceutically acceptable carriers or diluents and a HMG CoA inhibitor to said subject.

INTERNATIONAL SEARCH REPORT

Int. Application No
PCT/DK 01/00058

A. CLASSIFICATION OF SUBJECT MATTER		
IPC 7	C07C69/708 C07D215/14 A61P3/04	C07C59/64 C07D307/40 A61P3/10
	C07D213/55 A61K31/435	C07D307/80 A61K31/34
	C07D317/54 A61K31/357	
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols) IPC 7 C07C C07D A61K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) EPO-Internal		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 00 63153 A (NOVO NORDISK AS) 26 October 2000 (2000-10-26) the whole document	1-89
X	WO 99 20614 A (REDDY RESEARCH FOUNDATION) 29 April 1999 (1999-04-29) the whole document	1-89
X	WO 99 19313 A (REDDY RESEARCH FOUNDATION) 22 April 1999 (1999-04-22) the whole document	1-89
X	WO 99 38850 A (REDDY RESEARCH FOUNDATION) 5 August 1999 (1999-08-05) the whole document	1-89
	--- -/-	
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the International filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "&" document member of the same patent family		
Date of the actual completion of the international search 18 May 2001		Date of mailing of the international search report 22.06.2001
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer Nebil Gecer

INTERNATIONAL SEARCH REPORT

Ints. Application No

PCT/DK 01/00058

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 99 16758 A (REDDY RESEARCH FOUNDATION) 8 April 1999 (1999-04-08) the whole document ---	1-89
X	WO 99 08501 A (REDDY RESEARCH FOUNDATION) 25 February 1999 (1999-02-25) the whole document ---	1-89
X	EP 0 903 343 A (SSP CO LTD) 24 March 1999 (1999-03-24) the whole document ---	1-89
X	CHEMICAL ABSTRACTS, vol. 65, no. 3, 1 August 1966 (1966-08-01) Columbus, Ohio, US; abstract no. 3781, ELZBIETA WYRZYKIEWICZ ET AL: "Optically active stilbene derivatives" XP002901748 abstract & ZESZYTY NAUK. UNIW. POZNANIU MAT., FIS., CHEM, no. 9, 1965, pages 29-45, -----	1-9, 14-34, 37,38, 41-56, 58,59

INTERNATIONAL SEARCH REPORT

International application No.
PCT/DK01/00058

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 71-74, 80, 83, 86, 89
because they relate to subject matter not required to be searched by this Authority, namely:
see next sheet
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/DK01/00058

Claims 71-74, 80, 83, 86, 89 relate to methods of treatment of the human or animal body by surgery or by therapy/ diagnostic methods practised on the human or animal body/Rule 39.1.(iv). Nevertheless, a search has been executed for these claims. The search has been based on the alleged effects of the compounds/compositions

INTERNATIONAL SEARCH REPORT

International Application No

PCT/DK 01/00058

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0063153 A	26-10-2000	AU 3958000 A	02-11-2000
WO 9920614 A	29-04-1999	AU 1120699 A	10-05-1999
		BR 9812770 A	12-12-2000
		CN 1281443 T	24-01-2001
		EP 1082313 A	14-03-2001
		NO 20002114 A	26-06-2000
WO 9919313 A	22-04-1999	BR 9812772 A	10-10-2000
		EP 1049684 A	08-11-2000
		NO 20002113 A	26-06-2000
		US 6054453 A	25-04-2000
WO 9938850 A	05-08-1999	AU 1887999 A	16-08-1999
		EP 1051403 A	15-11-2000
WO 9916758 A	08-04-1999	US 6130214 A	10-10-2000
		ZA 9809790 A	28-04-1999
WO 9908501 A	25-02-1999	AU 1120599 A	08-03-1999
		CN 1280571 T	17-01-2001
		EP 1073643 A	07-02-2001
EP 0903343 A	24-03-1999	CN 1215051 A	28-04-1999
		JP 11158144 A	15-06-1999

1. An inspection lamp for detecting refrigerant leaks from air conditioning systems through illumination of fluorescent materials added to the refrigerant and lubricating oil, said inspection lamp comprising an LED pen light having a housing enclosing a power source and an LED, the housing having a rear end and a front end, with a glass lens capsule at the front end, said inspection lamp further comprising:

a hollow handle adapted to receive the pen light, whereby the pen light may be retracted into the handle or extended out of the handle, and

wherein the LED emits light within a wavelength band below about 600 nm.

2. An inspection lamp as in claim 1, wherein the LED emits light within a wavelength band between about 315 nm and about 400 nm.

3. An inspection lamp as in claim 1, wherein the LED emits light within a wavelength band between about 400 nm and about 480 nm.

4. An inspection lamp as in claim 1, wherein the LED emits light within a wavelength band between about 480 nm and about 600 nm.

5. An inspection lamp as in claim 1, wherein the handle is slidably engaged within the housing and has a threaded top portion adapted to receive a locking mechanism for locking the pen light at a desired extension out of the handle.

22. An LED inspection lamp for detecting refrigerant leaks from air conditioning systems through illumination of fluorescent materials added to the refrigerant and lubricating oil, said inspection lamp comprising:

extendable and retractable handle having a grip section and telescoping sections adapted to be pulled out to extend the handle and pushed in to retract the handle;

a battery housing containing a battery, the battery housing being attached at one end thereof to the telescoping section that is furthest from the grip section when the handle is extended, and

an LED lamp assembly connected to the battery housing and containing an LED.

23. An inspection lamp as in claim 22, further comprising the lamp having controls to selectively connect the LED to the battery.

24. An inspection lamp as in claim 23, wherein the lamp assembly contains a plurality of LEDs.

25. An inspection lamp as in claim 24, further comprising the lamp having controls to selectively connect an LED to the battery.

26. An inspection lamp as in claim 22, wherein the LED emits light within a wavelength band between about 315 nm to about 400 nm.

27. An inspection lamp as in claim 22, wherein the LED emits light within a wavelength band between 400 nm to about 480 nm.

28. An inspection lamp as in claim 22, wherein the LED lamp assembly is removable and a first LED lamp assembly having an LED emitting light within a first wavelength band is interchangeable with a second LED lamp assembly having an LED emitting light within a different wavelength band.

29. An inspection lamp as in claim 22, further comprising a hinge located between the grip end of the handle and the LED lamp assembly.